

Annual patterns of presence and activity of marine bacteria monitored by 16S rDNA–16S rRNA fingerprints in the coastal NW Mediterranean Sea

Raphaël Lami^{1,2,4,*}, Jean-François Ghiglione^{1,2}, Yves Desdevises^{1,3}, Nyree J. West^{1,2}, Philippe Lebaron^{1,2}

¹Université Pierre Marie Curie – Paris 6, Laboratoire ARAGO, Avenue du Fontaule, BP44, 66650 Banyuls-sur-Mer, France

²CNRS, UMR 7621, Laboratoire d'Océanographie Biologique de Banyuls, Avenue du Fontaule, BP44, 66650 Banyuls-sur-Mer, France

³CNRS, UMR 7628, Modèles en Biologie Cellulaire et Evolutive, Avenue du Fontaule, BP44, 66650 Banyuls-sur-Mer, France

⁴Present address: College of Marine and Earth Studies, University of Delaware, Lewes, Delaware 19958, USA

ABSTRACT: The annual dynamics of the bacterial community structure and its activity remain poorly studied in marine environments. Our goal was to gain new insights into the year-long bacterial dynamics and activity in correlation with environmental changes in the NW Mediterranean Sea. To accomplish this goal, we combined 16S rDNA versus 16S rRNA fingerprints and environmental variables using multivariate analyses. A clone library was constructed to determine to which bacterial taxa the major ribotypes were related. Our results revealed similar environmental controls of bacterial community structure at both the DNA and RNA levels. The 16S rRNA signal of several ribotypes differed relative to their 16S rDNA counterpart, suggesting that the members of the community have either a higher or lower activity than their relative abundance would suggest. Such differences included the dominant ribotypes observed in the fingerprints, such as *Roseobacter*, *Synechococcus*, SAR11 and SAR116. All these results give support to the 16S rDNA and 16S rRNA fingerprinting approach to monitor bacterial community structure and activity in marine environments.

KEY WORDS: Bacteria · 16S rDNA – 16S rRNA · Bacterial activity · CE-SSCP · Canonical correspondence analysis · Mediterranean Sea · Time series

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Despite the importance of bacteria in marine biogeochemical cycles (Azam et al. 1983), little is known about the dynamics of bacterial populations and of their activities (Fuhrman et al. 2006). In particular, the response of the bacterial community to environmental changes over mid- and long-term scales remains poorly characterized. The short-term variations of bacterial populations have been frequently studied in the marine environment (i.e. Zubkov et al. 2001, Fandino et al. 2005, Pinhassi et al. 2005), but only a few studies

focused on the annual dynamics of the bacterial community composition throughout the year (Murray et al. 1998, Ghiglione et al. 2005, Mary et al. 2006). In addition, only a few studies have examined the activity of these populations (Alonso-Saez & Gasol 2007). Bulk measurements of bacterial production or respiration have been conducted over a year (Lemée et al. 2002, Reinthaler & Herndl 2005), but these studies did not provide any information regarding the distribution of activity among different members of the community.

The distribution of activity among different members of the bacterial community has been investigated

*Email: lami@udel.edu

using micro-autoradiography combined with fluorescence *in situ* hybridization (micro-FISH) to examine uptake of radiolabeled substrates by specific phylogenetic groups (Alonso-Saez & Gasol 2007). However, this technique is too laborious for intensive samplings and is limited to only a few substrates that can be radiolabeled. Capillary-based fingerprinting techniques are well adapted to study bacterial dynamics in marine environments at a high throughput (Brown et al. 2005). The 16S rDNA profiles describe the relative composition of the bacterioplankton community (i.e. Muyzer et al. 1993, Lee et al. 1996), and previous studies used the 16S rRNA profiles to characterize the active fraction of bacterial communities in different environments (i.e. Moeseneder et al. 2001, Troussellier et al. 2002, Gentile et al. 2006), based on the observation that the number of ribosomes per cell is proportional to the growth rate (DeLong et al. 1989, Kemp et al. 1993, Kerkhof & Ward. 1993, Lee & Kemp 1994). To our knowledge, little is known on how the activity of individual members of a bacterial community varies in response to environmental changes.

In this study, we used an integrated approach combining capillary electrophoresis–single strand conformation polymorphism (CE-SSCP) of 16S rDNA and 16S rRNA profiles to obtain insights into the dynamics of the total and active communities during a 1 yr survey in the NW Mediterranean Sea. For text simplicity, we use the expression '16S rDNA' to refer to the 16S rRNA genes, as did several authors in recently published papers (i.e. Gentile et al. 2006, West et al. 2008). Multivariate analyses were performed to determine the fraction of the variance in both present and active bacterial communities that could be explained by the local environmental factors and time. In addition, a clone library was constructed to identify the ribotypes dominating the fingerprints.

MATERIALS AND METHODS

Sample collection and environmental measurements. Sampling was done from March 2003 and April 2004 at the SOLA station (42° 31' N, 03° 11' E), in the Bay of Banyuls-sur-Mer, on the French Mediterranean coast. Seawater was collected in 5 l Niskin bottles at 5 m depth. Environmental variables were measured by the French coastal monitoring program Service d'Observation en Milieu Littoral (SOMLIT) (www.domino.u-bordeaux.fr/somlit_national) according to protocols previously described (i.e. Joux et al. 2006) and available on the SOMLIT website. All samples were processed after sampling within 30 min.

Microbiological variables. Bacterial abundances were measured using the flow cytometry protocol de-

scribed by Lebaron et al. (1998). Briefly, 2 ml of seawater was pre-filtered at 5 µm before being fixed (2% formaldehyde, 1 h, 4°C) and incubated with SYBR Green II (final conc. 0.05% [v/v] of the commercial solution, ≥15 min, at 20°C in the dark). A 10 to 15 µl volume of this solution was used to enumerate and characterize cells with their right angle light scatter and their green fluorescence collected through a 530 ± 30 nm bandpass filter on a FACS Calibur cytometer (Becton Dickinson, San Jose, CA), equipped with an air-cooled argon laser (488 nm). Polycarbonate beads (1.002 µm, Polyscience Europe) were used to normalize cell fluorescence and light scattering values. Data were obtained and analyzed with CellQuest software (Becton Dickinson). Acquisition was triggered by green fluorescence. The volume analyzed (and thus cell concentrations) was calculated by measuring the remaining volume and subtracting it from the initial volume. The sheath fluid was seawater filtered at 0.22 µm.

Nucleic acid extraction. A total of 5 l of seawater was pre-filtered onto 3 µm pore size filters (47 mm Nucleopore) before being filtered onto a 0.2 µm Sterivex-GV system (Millipore). The Sterivex filters were frozen at –80°C after addition of 1.8 ml of lysis buffer (EDTA 40 mM; Tris 50 mM pH = 8.3, sucrose 0.75 M) until extraction of nucleic acids. First, cells were lysed by adding 50 µl of lysozyme (1 mg ml⁻¹, final conc.) to the lysis buffer on the filter and incubating at 37°C for 1 h. Next, 100 µl of sodium dodecyl sulfate (10% v/v) and 10 µl of proteinase K were added, and then the preparation was incubated a second time at 55°C for 1 h. The lysate was examined under the microscope to check for complete lysis. Total DNA was extracted with the DNeasy tissue kit (Qiagen) with a modified protocol for marine samples. In brief, an 800 µl volume of lysate was extracted from the Sterivex, to which 800 µl of ethanol (95%) was added to clear the cell lysate before adding 400 µl of AW2 buffer. The final steps of the extraction were conducted by following the instructions of the manufacturer. Total RNA was extracted with an SV Total RNA kit (Promega) modified protocol. A total of 600 µl of lysate was extracted from the Sterivex and cleared with 240 µl of ethanol (95%). The final steps of RNA extraction were conducted using the manufacturer protocol. Molecular size and purity of the DNA and RNA were analyzed by agarose gel and spectrophotometry (GeneQuant II, Pharmacia Biotech). DNA removal was controlled by PCR amplification of 1 µl of each RNA sample (PCR conditions described below). In these controls, no amplifications were detected.

16S cDNA synthesis. 16S cDNA was synthesized just after total RNA extraction. RNA (2 µg) was incubated for 5 min at 90°C before cooling in a water/ice bath. Twenty units of M-MLV (Roche) was added, with 5 µl of buffer, and 50 µM of primers (Eurogentec). By using

the same protocols, Moeseneder et al. (2001) did not find a chimeric structure after a cloning-sequencing step of obtained cDNA.

PCR, RT (reverse transcription)-PCR and CE-SSCP.

Specific bacterial primers were used for PCR and RT-PCR: w49dir (5'-CGGTCCAGACTCCTACGGG-3'; Delbès et al. 1998) and w34rev (5'-TTACCGCGGCT-GCTGGCAC-3'; Lee et al. 1996). Both of these were synthesized commercially (Eurogentec). Primer w34 was fluorescently 5'-labeled with phosphoramidite (TET, Eurogentec). Primers were designed to target a short sequence (around 200 bp) for the best resolution of the CE-SSCP signal (V3 region of 16S rDNA, *Escherichia coli* positions 329 to 533; Brosius et al. 1981). Amplifications were performed according to a previously described protocol (Ghiglione et al. 2005). The CE-SSCP analysis used a Genetic Analyzer (Applied Biosystems) and protocols described in Ghiglione et al. (2005). Briefly, each sample was diluted between 2- and 40-fold in sterile water (molecular biology grade, Sigma) to obtain 10 ng μl^{-1} of PCR products. From the diluted solution, 1 μl of PCR product was mixed with 0.1 μl of size standard (GeneScan 400 ROX, Applied Biosystems) and 18.9 μl of deionized formamide (Aplera), before heating (94°C, 10 min) followed by rapid cooling in a water/ice bath for 10 min. Samples were electrokinetically injected (5 s, 12 kV) into a capillary (47 cm \times 50 μm) filled with polymer solution (5.6% GeneScan polymer gel, 10% autoclaved glycerol and 1 \times EDTA buffer; Applied Biosystems) and were electrophoresed at 12 kV and 30°C for 30 min. The fluorescently labeled fragments were detected by laser with the virtual filter C (detection wavelengths 532, 537 and 584 nm). Data on the reproducibility of our methods on samples collected at the same station have been previously published (Ghiglione et al. 2005). Similar observations have also been collected by Zumstein et al. (2000). Variation of peak area between normalized similar profiles remained below 5% (data not shown).

Data were collected using ABI Prism 310 collection software (Applied Biosystems). Electrophoregrams were aligned with GeneScan analysis software by fixing positions of the internal standard (GeneScan 400 ROX) with a second-order, least-squares curve fit. Peak fingerprints were based on the first derivative of a polynomial curve fitted to the data within a window centered on each data point (GeneScan analysis software). As some of the peaks overlapped in the environmental fingerprints we obtained, the polynomial degree was increased to 10 to avoid artifacts in peak enumeration or in peak area estimates and to maximize the sensitivity of peak detection. The peak amplitude threshold applied for both ROX and TET fluorescent dyes was fixed at 50. Peak areas were calculated by the GeneScan analysis software, based on the half-height peak width.

Total area of profiles was normalized to 100, and then the area of each identified peak was expressed as a percentage of the total area of each scan.

Clone library and CE-SSCP peak assignments. CE-SSCP peaks were assigned using a clone library constructed from samples taken at the same location on April 1, 2003. Cloning and restriction fragment length polymorphism (RFLP) procedures were performed as previously described (Ferrera et al. 2004), using bacterial primers SAdir (5'-AGAGTTTGATCATGGCTC-AGA-3', *Escherichia coli* positions 8 to 28) and S17rev (5'-GTTACCTTGTTACGACTT-3', *E. coli* positions 1492 to 1508). Among 184 clones, 84 different bacterial phylotypes were determined by RFLP, and 1 representative from each phylotype was sequenced and identified in the GenBank data base using BLAST (www.ncbi.nlm.nih.gov/blast) to obtain their closest phylogenetic neighbors. The phylotypes identified as chimeras were removed from the data set using the RDP-II Chimera Check program (Cole et al. 2005). These sequences were submitted to GenBank under the accession numbers DQ436518 to DQ436545, DQ436638 to DQ436663, DQ436718 to DQ436733 and DQ436775 to DQ436778 (see Table 2). Phylogenetic groups or operational taxonomic units (OTUs) were defined at 99% similarity from a ClustalW alignment (500 bp) using the Clusterer software with the single linkage setting (Klepac-Ceraj et al. 2006). We assigned an environmental fingerprint obtained at the sampling point for the same date. Each clone was prepared according to the previously described protocols for CE-SSCP samples, and then aligned with each other and the environmental profiles using the same size standard as that used for all our environmental fingerprints.

Data analyses. Data were analyzed using 2 software packages: XLSTAT 2006 (Addinsoft, Paris) and CANOCO (ter Braak 1987) version 4.0. Fingerprints were analyzed by their peak areas, generating an intensity matrix. The similarity matrix was computed using Steinhaus's similarity index. A canonical correspondence analysis (CCA) was used to investigate the variations in the intensity matrix under the constraint of our set of environmental variables. Significant variables (i.e. variables that significantly explained changes in 16S rDNA and 16S rRNA signals) in our data set were chosen using a forward-selection procedure and 999 permutations. In our analysis, we also introduced time as a variable to determine which proportion of the total variance in the 16S rDNA and 16S rRNA could be thus explained, following the variation partitioning approach described in Borcard et al. (1992). We assumed in this work a unimodal response of species to environmental variations. When a linear response was assumed, the percentage of variation explained was lower (data not shown).

RESULTS

Physical, chemical and biological characteristics of the sampling site

Environmental factors measured over the study period are reported in Table 1 and are expressed as mean values \pm SD. Temperature varied between 11.6 and 21.7°C in spring ($15.0 \pm 3.5^\circ\text{C}$), 21.4 and 25.2°C in summer ($23.0 \pm 1.6^\circ\text{C}$), 13.7 and 19.7°C in autumn ($16.9 \pm 2.3^\circ\text{C}$), and 10.5 and 12.3°C in winter ($11.5 \pm 0.8^\circ\text{C}$). Salinity varied seasonally with mean \pm SD values of 37.0 ± 0.5 in spring, 37.9 ± 0.1 in summer, 37.4 ± 1.1 in autumn and 37.7 ± 0.2 in winter. Chl *a* was relatively low in summer ($0.3 \pm 0.3 \text{ g l}^{-1}$) and highest in winter ($1.3 \pm 0.4 \text{ }\mu\text{g l}^{-1}$). Average values in spring and autumn were $0.6 \pm 0.4 \text{ }\mu\text{g l}^{-1}$ and $0.7 \pm 0.2 \text{ }\mu\text{g l}^{-1}$, respectively. Nutrients were present at either low or undetectable concentrations in summer ($\text{NH}_4^+ = 0.1 \pm 0.1 \text{ }\mu\text{M}$, $\text{NO}_3^- = 0.1 \pm 0.1 \text{ }\mu\text{M}$, $\text{NO}_2^- =$ undetectable levels, $\text{PO}_4^{3-} = 0.1 \pm 0.1 \text{ }\mu\text{M}$, $\text{Si(OH)}_4 = 1.3 \pm 0.4 \text{ }\mu\text{M}$). Nitrates, nitrites and silicates reached maximum concentrations in winter

(1.7 ± 0.8 , 0.4 ± 0.1 and $2.0 \pm 0.7 \text{ }\mu\text{M}$, respectively). Concentrations of ammonium were higher, on average, in spring and autumn (mean $0.4 \pm 0.2 \text{ }\mu\text{M}$). Phosphate concentrations remained low year round, the values being highest in autumn ($0.04 \text{ }\mu\text{M}$ on September 23, 2003) and in spring ($0.03 \text{ }\mu\text{M}$ on April 15, 2003). Bacterial abundances varied between 6.23×10^5 and $1.31 \times 10^6 \text{ cells ml}^{-1}$ ($8.93 \pm 1.78 \times 10^5 \text{ cells ml}^{-1}$).

Clone library and assignment of the corresponding CE-SSCP profile

CE-SSCP peaks were assigned using a clone library constructed from samples taken at the same location on April 1, 2003. The library coverage was 61% and the total number of phylotypes was estimated to be 139 using the Chao-1 estimator (Chao 1987). Clustering of sequences (99% similarity) revealed 7 major clusters of *Alphaproteobacteria* (*Roseobacter* RCA, *Roseobacter* CHAB-I-5, *Roseobacter* PLY-P3-48, SAR11 S1, SAR11 S3, SAR116, LA1-B32N), 2 major clusters of *Bac-*

Table 1. Environmental variables and bacterial abundances collected at the SOLA station (3 m water depth). nd = undetectable concentration, nm = not measured. Dates given as mo/d/yr

| Sampling date | Temp. (°C) | Salinity (PSU) | Chl <i>a</i> (g l ⁻¹) | NH ₄ ⁺ (μM) | NO ₃ ⁻ (μM) | NO ₂ ⁻ (μM) | PO ₄ ³⁻ (μM) | Si(OH) ₄ (μM) | Bacteria (×10 ⁵ ml ⁻¹) |
|---------------|------------|----------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|------------------------------------|--------------------------|---|
| Spring | | | | | | | | | |
| 04/15/03 | 12.5 | 37.5 | 0.33 | 0.59 | 1.24 | 0.14 | 0.03 | 2.8 | 11.1 |
| 04/29/03 | 14.0 | 37.2 | 0.91 | 0.74 | 0.64 | 0.07 | 0.02 | 0.65 | 9.67 |
| 05/13/03 | 16.2 | 37.2 | 0.66 | 0.20 | 0.08 | 0.02 | 0.02 | 0.85 | 7.37 |
| 05/27/03 | 15.9 | 37.6 | 0.72 | 0.26 | 0.26 | 0.03 | 0.02 | 1.45 | 8.23 |
| 06/11/03 | 21.7 | 36.5 | 0.58 | 0.15 | 0.42 | 0.03 | 0.03 | 1.32 | 8.24 |
| Summer | | | | | | | | | |
| 06/24/03 | nm | 37.7 | 0.29 | 0.09 | 0.24 | nd | 0.15 | 1.84 | 7.19 |
| 07/08/03 | 21.3 | 37.8 | 0.12 | 0.16 | 0.20 | nd | 0.12 | 1.00 | 9.57 |
| 07/22/03 | 23.0 | 37.9 | 0.26 | 0.13 | 0.04 | nd | nd | 1.35 | 13.4 |
| 08/05/03 | 23.6 | 38.0 | 0.24 | 0.22 | 0.02 | 0.02 | 0.02 | 1.23 | 7.46 |
| 08/19/03 | 25.2 | 37.7 | 0.28 | 0.03 | 0.08 | 0.06 | 0.02 | 1.69 | 7.29 |
| 09/09/03 | 21.8 | 37.7 | 0.38 | 0.08 | 0.02 | nd | 0.03 | 0.95 | 9.93 |
| Autumn | | | | | | | | | |
| 09/23/03 | 19.7 | 37.9 | 0.29 | 0.18 | 0.01 | 0.01 | 0.04 | 1.82 | 8.77 |
| 10/07/03 | 19.2 | 37.9 | 0.40 | 0.34 | 0.21 | 0.02 | 0.03 | 1.09 | 9.61 |
| 10/14/03 | 18.1 | 38.0 | 0.58 | nm | nm | nm | nm | nm | 9.84 |
| 10/21/03 | 17.0 | 37.8 | 0.76 | nm | nm | nm | nm | nm | 11.1 |
| 11/03/03 | 15.8 | 38.0 | 0.66 | 0.24 | 0.20 | 0.15 | 0.02 | 1.83 | 9.38 |
| 11/18/03 | 14.6 | 36.5 | 1.39 | 0.76 | 2.54 | 0.43 | 0.02 | 1.33 | 13.1 |
| 12/09/03 | 13.7 | 35.2 | 0.39 | 0.35 | 5.15 | 0.55 | 0.16 | 8.83 | 9.92 |
| Winter | | | | | | | | | |
| 01/06/04 | 12.3 | 37.8 | 0.75 | 0.40 | 1.08 | 0.47 | 0.01 | 2.65 | 7.81 |
| 01/27/04 | 11.9 | 37.8 | 1.34 | 0.29 | 1.24 | 0.31 | 0.02 | 1.98 | 8.03 |
| 02/17/04 | 11.0 | 37.5 | 1.77 | nm | 1.85 | 0.29 | nd | 1.05 | 9.8 |
| 03/09/04 | 10.5 | 37.5 | 1.18 | 0.26 | 2.78 | 0.4 | nd | 2.34 | 6.95 |
| Spring | | | | | | | | | |
| 03/23/04 | 11.6 | 36.7 | 0.97 | 0.43 | 2.66 | 0.35 | 0.02 | 2.01 | 8.76 |
| 04/06/04 | 12.6 | 36.2 | 0.79 | 0.41 | 2.42 | 0.08 | 0.01 | 1.72 | 6.23 |

teroidetes sequences (Agg58, PLY-P3-79) and 5 major clusters of *Gammaproteobacteria* (*Alteromonas*, OM60 [HTCC2080], KTc1119, HTCC2149 and SAR86-II [EBAC31A08]) (Table 2).

The bacterial community structure from April 1, 2003 was determined by combining the corresponding CE-SSCP fingerprint with the clone library (Fig. 1). Based on the retention times of the cloned 16S rDNA, we observed that each phylogenetic group had a different migration pattern. *Alphaproteobacteria* clones had the shortest retention times, *Gammaproteobacteria* clones were detected with the largest retention times and *Bacteroidetes* clones were detected in between (except the 2 *Gammaproteobacteria* clones related to the OM60 cluster, which co-migrated with the *Bacteroidetes*) (Fig. 1). For the

minor clone groups, 1 *Epsilonproteobacteria* and 3 *Betaproteobacteria* clones migrated with retention times comparable to those observed for *Bacteroidetes* and *Gammaproteobacteria* clones, respectively. At a lower phylogenetic rank, all 10 *Roseobacter* clones and very close relatives migrated closely together. A similar observation was made for the 5 clones affiliated with *Synechococcus*. Two other ribotypes which appeared dominant on various profiles collected during the year are composed of different populations of *Alphaproteobacteria*, mostly SAR11 and SAR116. Interestingly, the 4 clones belonging to the *Bacteroidetes* Agg58 cluster and the 3 belonging to the *Gammaproteobacteria* cluster OM60 also co-migrated, allowing us to assign 2 other peaks in the fingerprints (Fig. 1).

Table 2. Clustering of sequences done with Clusterer at 99 % similarity after a ClustalW alignment. Alignment was trimmed to 500 bp before cluster analysis

| Group | Accession number |
|---|--------------------------------|
| <i>Alphaproteobacteria</i> | |
| <i>Roseobacter</i> RCA (DC11-80-2) | DQ436523 / DQ436544 / DQ436535 |
| <i>Roseobacter</i> (CHAB-I-5) | DQ436538 / DQ436524 / DQ436527 |
| | DQ436520 / DQ436533 |
| <i>Roseobacter</i> (PLY-P3-48) | DQ436526 / DQ436532 |
| SAR11 S1 (MB11F01) | DQ436518 / DQ436540 / DQ436545 |
| | DQ436531 |
| SAR11 S3 (Z0410) | DQ436522 / DQ436530 |
| SAR116 (ZD0209) | DQ436519 / DQ436528 |
| LA1-B32N | DQ436537 / DQ436529 |
| Other <i>Alphaproteobacteria</i> | DQ436536 / DQ436539 / DQ436541 |
| | DQ436542 / DQ436543 / DQ436521 |
| | DQ436534 / DQ436545 |
| <i>Bacteroidetes</i> | |
| Agg58 | DQ436719 / DQ436723 / DQ436725 |
| | DQ436726 |
| PLY-P3-79 | DQ436733 / DQ436727 |
| Unidentified <i>Bacteroidetes</i> cluster | DQ436731 / DQ436718 / DQ436721 |
| Other <i>Bacteroidetes</i> | DQ436720 / DQ436722 / DQ436724 |
| | DQ436728 / DQ436729 / DQ436730 |
| | DQ436732 |
| <i>Gammaproteobacteria</i> | |
| <i>Alteromonas</i> | DQ436640 / DQ436643 |
| | DQ436645 / DQ436651 |
| OM60 (HTCC2080) | DQ436653 / DQ436644 / DQ436638 |
| KTc1119 | DQ436639 / DQ436660 |
| HTCC2149 | DQ436650 |
| SAR86-II (EBAC31A08) | DQ436641 / DQ436646 |
| Other <i>Gammaproteobacteria</i> | DQ436642 / DQ436654 / DQ436655 |
| | DQ436656 / DQ436659 / DQ436662 |
| | DQ436663 / DQ436647 / DQ436648 |
| | DQ436649 |
| Other groups | |
| <i>Betaproteobacteria</i> | DQ436658 / DQ436661 / DQ436652 |
| <i>Epsilonproteobacteria</i> | DQ436657 |
| <i>Synechococcus</i> IV (MB11E09) | DQ436775 / DQ436776 / DQ436777 |
| | DQ436778 / DQ436775 |

Year-long monitoring of presence/absence and relative abundance of 16S rDNA and 16S rRNA ribotypes

A total of 54 different ribotypes were observed during the year. Each 16S rDNA fingerprint had between 22 and 35 peaks (29.4 ± 4.0), and between 11 and 30 ribotypes were observed in each 16S rRNA fingerprint (24.4 ± 4.8) (Fig. 2). For both the 16S rDNA and 16S rRNA data sets, the total number of ribotypes varied between 31 and 46 for each sampling date (38.4 ± 4.8 ribotypes) (Fig. 2). Most of the ribotypes were detected in both 16S rDNA and 16S rRNA fingerprints, between 30 and 90% of the ribotypes per sampling date ($66.7 \pm 13.7\%$). A ribotype was detected in a 16S rDNA fingerprint but not in the corresponding 16S rRNA fingerprint on average for only 7.6 ± 5.5 ribotypes per fingerprint. A ribotype was detected on a 16S rRNA fingerprint but not in the corresponding 16S rDNA profile for only 2.6 ± 1.9 ribotypes per fingerprint, on average (Fig. 2).

Interestingly, both 16S rDNA and 16S rRNA fingerprints were dominated all year by 4 ribotypes (Figs. 3 & 4). When considering a specific sampling date, these 4 dominant ribotypes represented between 20 and 80% ($48.7 \pm 14.9\%$) of the 16S rDNA fingerprint area, and between 30 and 80% ($53.8 \pm 17.1\%$) of the 16S rRNA fingerprints area. There were some large differences between the area of a 16S rDNA ribotype and its area on the cor-





Fig. 1. Assignment of an environmental capillary electrophoresis–single strand conformation polymorphism (CE-SSCP) fingerprint obtained from the April 1, 2003 sample with a clone library created in parallel. The superimposition of the clones on the environmental fingerprint is represented by vertical bars. Clones include A (= *Epsilonproteobacteria* clone) and B,C (= 2 *Betaproteobacteria* clones). The number of clones from each bacterial group is indicated in parentheses. Retention time is in arbitrary units



responding 16S rRNA fingerprints, including the 4 dominant ribotypes. The ratio of 16S rDNA to 16S rRNA area for these major CE-SSCP peaks ranged from 0 to 3 (1.2 ± 0.9) (*Roseobacter*), 0 to 7 (1.8 ± 1.5) (*Synechococcus*), 0 to 2 (1.0 ± 0.3) (SAR11) and 0 to 4 (1.0 ± 0.9) (SAR116) (Figs. 3 & 4). For these 4 dominant ribotypes, some differences were detected between 16S rDNA and 16S rRNA signals during the year-long sampling. For example, in autumn 2003, *Synechococcus* was detected on 16S rRNA profiles, but had a relatively low area on 16S rDNA fingerprints. The same phenomenon occurred, but in lower proportions, for *Roseobacter* and SAR11 in spring and autumn 2003 (Figs. 3 & 4).

Analysis of 16S rDNA data set

Variation partitioning indicated that environmental variables explained 49% of the total variation observed in the 16S rDNA data set (Fig. 5). Seasonal groups were well defined by CCA on the basis of calendar dates (Fig. 6). The forward selection of environmental data indicated that 6 of the 9 variables (Table 1) were significantly related to the variation of 16S rDNA ribotypes important in the bacterioplankton community ($p < 0.05$, vectors in Fig. 6). Bacterial abundances, inorganic phosphate and silicate, did not significantly influence the 16S rDNA signal. The signal was highly correlated to the selected environmental variables (Fig. 5). These variables were divided into 2 groups on axis 1 of the CCA (Fig. 6): temperature and salinity versus NH_4^+ , NO_3^- , chl *a* and NO_2^- , reflecting a strong seasonal signal. In our data set, 36% of the observed

Fig. 2. Percentage of total ribotypes unique to the 16S rDNA (DNA) or the 16S rRNA (RNA) fingerprints, or common to both (DNA+RNA). Total number of ribotypes indicated at the top of each bar (mean 30.4 ± 4.9 SD). Dates given as mo/d/yr




Fig. 3. Dynamics of the 4 dominant CE-SSCP ribotypes (*Roseobacter*, *Synechococcus*, SAR11 and SAR 116) during the year-long sampling of 16S rDNA and 16S rRNA fingerprints




Fig. 4. Variation of the 16S rRNA/16S rDNA area ratio for the 4 dominant ribotypes observed in the fingerprints (see Fig. 3)




Fig. 5. Variation partitioning analysis of 16S rDNA and 16S rRNA data sets. Significant variables, determined following a forward-selection procedure, were temperature, salinity, chl *a*, ammonium, nitrate and nitrite. For each data set, $p < 0.01$

variance remain unexplained. When time was coded as a variable it explained 27% of the total variance (pure time 15%) (Fig. 5).

Analysis of 16S rRNA data set

Variation partitioning indicated that environmental variables explained 45.7% of total variance present in the 16S rRNA data set (Fig. 5). Patterns similar to those observed for the 16S rDNA dataset were found (Fig. 6). The CANOCO forward-selection process selected the same significant variables ($p < 0.05$, vectors on Fig. 6) as in the 16S rDNA data set, and the CCA also showed the same separation of environmental variables on the canonical axis 1 (Fig. 6), with temperature and salinity versus NH_4^+ , NO_3^- , chl *a* and NO_2^- reflecting a seasonal control of the bacterial activity. The 16S rRNA signal was also highly correlated to the environmental variables (Fig. 5). Variation partitioning indicated that 40% of the observed variance in the 16S rRNA data set remained unexplained. Interestingly, time explained 24% of this variance (pure time 14%) (Fig. 5).

Projection of particular OTUs

CCA allows the projection of sampling dates but also of bacterial ribotypes in the constructed 2-dimensional




Fig. 6. Canonical correspondence analysis on (A) 16S rDNA fingerprints and (B) 16S rRNA fingerprints. Fingerprints are clustered according to the calendar dates of the seasons. (▲) Position of the dominant and of noteworthy ribotypes in the 2-dimensional space of the plot. *Roseo.* = *Roseobacter*, *Syn.* = *Synechococcus*. Dates are given as mo/d/yr

space. *Roseobacter* presence (16S rDNA) and activity (16S rRNA) were linked to spring, and were independent of temperature (Fig. 6). *Synechococcus* presence and activity were linked to temperature. The dominant ribotypes, including mainly SAR11 and SAR116 populations, were associated with autumn, in terms of both presence and activity. When considering the other assigned ribotypes, we noticed that the Agg58 and OM60 clades were linked with winter and chl *a*.

DISCUSSION

In this work, the dynamics of marine bacterial populations and their activity were examined during a year-long study in NW Mediterranean coastal waters. For each sampling date, a 16S rDNA CE-SSCP fingerprint (reflecting the bacterial community structure) was compared to the corresponding 16S rRNA fingerprint (reflecting the active component of the community) and a set of environmental factors. A complementary clone library allowed the identification of some of the members of the studied community.

Each 16S rDNA and 16S rRNA date-ribotype area matrix constitutes a multidimensional data set. Multivariate analyses were used to represent these data by projecting the fingerprints in a 2-dimensional space using CCA, which calculates axes by maximizing the variance between each fingerprint (Legendre & Legendre 1983). In CCA, the ordination of samples in the 2-dimensional space is constrained by the selected environmental variables (ter Braak 1987). This analysis revealed seasonal patterns in both 16S rDNA and 16S rRNA signals, suggesting a seasonal dynamic of bacterial populations and their activities. Similar trends were previously observed with 16S rDNA signals (i.e. Ghiglione et al. 2005, Mary et al. 2006) but no previous studies examined a year-long variation of the 16S rRNA signal. This similar seasonal trend on a 16S rRNA-based level was not surprising because most of the ribotypes (mean 67%) were common to both 16S rRNA and 16S rDNA fingerprints.

The use of variation partitioning analysis in microbial ecology to link bacterial community changes with environmental factors is rare (Ramette 2007). Both biotic (chl *a*) and abiotic (temperature, nutrients) variables had a marked influence on the bacterial community composition (bottom-up effects), consistent with other studies (Schauer et al. 2003, Brown et al. 2005, Fuhrman et al. 2006). Our data indicate that the same set of environmental variables similarly influenced bacterial populations and their activities. Nevertheless, 40% of the variance in the 16S rDNA and in the 16S rRNA data sets remains unexplained. Some unrecorded environmental variables like dissolved

organic matter (bottom-up effects) (Schauer et al. 2003, Mary et al. 2006), grazer community composition (top-down effects) or viral activity may have affected the bacterial populations and their activities. Interestingly, our partitioning analysis indicated that time also explained a noticeable fraction (pure time, 15%) of the total variance. This result suggests that time alone may partially explain part of the bacterial community structure.

In this study, 4 dominant ribotypes were detected on 16S rDNA fingerprints when considering the average area of each ribotype over the year. The same dominant ribotypes were also observed in 16S rRNA fingerprints, which implies that the dominant bacterial populations were also the most active. It is well established that marine bacterial populations are dominated by a few taxa (Giovannoni et al. 1990, Giovannoni & Rappé 2000). However, recent studies have indicated that some of the most abundant and ubiquitous taxa of bacterioplankton are also very micro-diverse (Acinas et al. 2004, Brown & Fuhrman 2005). Thus, the dominant ribotypes we detected are likely to be a result of the complex signal created by a variety of micro-diverse populations, which could also be physiologically distinct between seasons.

We identified the bacterial populations detected by the CE-SSCP peaks by constructing a clone library that was used to assign peaks to the profiles. Two of the dominant ribotypes detected on our fingerprints were putatively assigned to *Synechococcus* and *Roseobacter*. The 2 other dominant ribotypes were composed of *Alphaproteobacteria* populations, mostly SAR11 and SAR116. *Gammaproteobacteria* and *Bacteroidetes* also represented a large proportion of the bacterial community, but were distributed within a high number of ribotypes in the fingerprints, suggesting that these bacterial populations were diverse and present in low abundances. We are also aware that primers used in the CE-SSCP study can introduce biases (as is the case for any PCR-based methods). It is well established that the available Bacteria primers are not truly universal, as there are some mismatches for certain phylogenetic groups (Baker et al. 2003). Indeed, there are some mismatches between the w34 primer used in our CE-SSCP protocols and some *Bacteroidetes* sequences (data not shown). Nevertheless, relative abundances of *Synechococcus* deduced from the fingerprints were highly correlated to those determined by flow cytometry ($p < 0.05$, $n = 24$), giving strong support to the assignation of the corresponding CE-SSCP peak (data not shown). Such observations also support our assignment procedures and reinforce the quantitative analysis of CE-SSCP peaks conducted in this study.

According to the CCA analysis, the *Synechococcus* ribotype (presence and activity) was correlated to

temperature, as previously reported (Agawin et al. 1998). The *Roseobacter* ribotype (presence and activity) was detected mostly during spring, as observed by Brown et al. (2005) using an automated ribosomal intergenic spacer analysis-based approach in Californian coastal waters. Interestingly, the presence and activity of the OM60 clade, known to harbor representatives of photoheterotrophic species (Fuchs et al. 2007), was closely correlated with chl *a*.

Common seasonal trends in the 16S rDNA and 16S rRNA data were found in our observations. However, some discrepancies between 16S rDNA and 16S rRNA signals were also detected. On the one hand, bacterial populations were sometimes present but not active. That is, a ribotype detected in 16S rDNA fingerprints was absent or had a small area for the corresponding ribotype on the 16S rRNA profile, suggesting low abundances of ribosomes per bacterial cell. Some authors who made similar observations have linked these patterns with periods of starvation (Davis et al. 1986, Moyer & Mortita 1989, Fegatella et al. 1998). On the other hand, bacterial populations were sometimes absent or present in low abundances, while their activity clearly appeared in the 16S rRNA fingerprint. That is, a specific ribotype was absent or had a small area on 16S rDNA fingerprints, but appeared on 16S rRNA fingerprints. Such patterns have been previously described in the marine environment, and are identified by the presence of a few abundant cells with a high metabolic potential (Moeseneder et al. 2001, Moeseneder et al. 2005).

Cottrell & Kirchman (2004) observed in Delaware River estuary waters that only 50% of the contribution of bacterial taxa to biomass production could be explained by their relative abundance. Thus, the activities of bacterial populations might be higher or lower than the abundance would predict. Such a result is congruent with the fact that the relationship between 16S rRNA content and growth rate is not linear. During spring 2003, a period of high bacterial production (I. Obernosterer pers. comm.), the ribotype assigned as *Roseobacter* was relatively constant and had high 16S rRNA/16S rDNA ratios. Thus, in spring 2003, the contribution of *Roseobacter* and SAR11 to the total bacterial production was higher than what could be predicted from their respective abundances. We also observed a similar situation for the functional group of the oxygenic photosynthetic *Synechococcus*, which is the most abundant cyanobacteria in our coastal system, and which had a high 16S rRNA/16S rDNA ratio in October 2003, suggesting high activity of this population at this time of the year. Interestingly, this peak of activity did not correlate with the period when the 16S rDNA and 16S rRNA ribotypes had the largest relative area, which was in summer.

Collectively, these data point out the interest in both 16S rDNA and 16S rRNA fingerprints for monitoring bacterial dynamics in marine environments. The 16S rDNA fingerprints describe reasonably well the dynamics of a bacterial community structure, while 16S rRNA fingerprints lead to the detection of the largest variations in bacterial activity, in response to environmental changes. Such results also reinforce the use of CE-SSCP as a tool to describe bacterial community structure and activity in long-term monitoring approaches in the marine environment, which is being done more and more at microbial observatories. However, the taxonomic resolution of fingerprints does not allow a fine identification of bacterial taxa which are responsible for the variations, even if a few ribotypes could be putatively assigned to widespread bacterial players (*Synechococcus*, *Roseobacter*). Nevertheless, the detection of significant variations in 16S rRNA relative to 16S rDNA fingerprints suggests that all the members of the bacterial community do not participate equally in the flow of carbon in coastal environments. These results highlight the need to re-evaluate the assumptions in biogeochemical cycles models, which implicitly hypothesize that all members of the prokaryotic community contribute equally to the flux of matter and energy (Alonso-Saez & Gasol 2007) whatever the season of the year.

Acknowledgements. We thank the captain and the crew of the RV 'Nereis II' for the sample collection. This work was supported by the European program BASICS (Bacterial Single Cell Analysis in the Sea, EVK3-CT-2002-00078). The project was also carried out in the context of the MarBEF Network of Excellence (Marine Biodiversity and Ecosystem Functioning), which is funded by the European Community's Sixth Framework Programme (contract no. GOCECT-2003-505446). R.L.'s research was supported by a doctoral fellowship from the French Research and Education Ministry and the post-doctoral fellowship Lavoisier from the French foreign office. We thank P. Catala and N. Batailler for technical help. We are also grateful to D. L. Kirchman for helpful comments on an earlier version of this manuscript.

LITERATURE CITED

- Acinas SG, Klepac-Ceraj V, Hunt DE, Pharino C, Ceraj I, Distel DL, Polz MF (2004) Fine-scale phylogenetic architecture of a complex bacterial community. *Nature* 430: 551–554
- Agawin NSR, Duarte CM, Agustí S (1998) Growth and abundance of *Synechococcus* sp. in a Mediterranean Bay: seasonality and relationship with temperature. *Mar Ecol Prog Ser* 170:45–53
- Alonso-Saez L, Gasol JM (2007) Seasonal variations in the contributions of different bacterial groups to the uptake of low-molecular-weight compounds in Northwestern Mediterranean coastal waters. *Appl Environ Microbiol* 73:3528–3535

- Azam F, Fenchel T, Field JG, Gray JS, Meyer-Reil LA, Thingstad F (1983) The ecological role of water-column microbes in the sea. *Mar Ecol Prog Ser* 10:257–263
- Baker GC, Smith JJ, Cowan DA (2003) Review and re-analysis of domain-specific 16S primers. *J Microbiol Methods* 55:541–555
- Borcard D, Legendre P, Drapeau P (1992) Partialling out the spatial component of ecological variation. *Ecology* 73:1045–1055
- Brosius J, Dull TJ, Sleeter DD, Noller HF (1981) Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*. *J Mol Biol* 148:107–127
- Brown MV, Fuhrman JA (2005) Marine bacterial microdiversity as revealed by internal transcribed spacer analysis. *Aquat Microb Ecol* 41:15–23
- Brown MV, Schwalbach MS, Hewson I, Fuhrman JA (2005) Coupling 16S-ITS rDNA clone libraries and automated ribosomal intergenic spacer analysis to show marine microbial diversity: development and application to a time series. *Environ Microbiol* 7:1466–1479
- Chao A (1987) Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* 43:783–791
- Cole JR, Chai B, Farris RJ, Wang Q and others (2005) The Ribosomal Database Project (RDP-II): sequences and tools for high-throughput rRNA analysis. *Nucleic Acids Res* 33:D294–D296
- Cottrell MT, Kirchman DL (2004) Single-cell analysis of bacterial growth, cell size, and community structure in the Delaware estuary. *Aquat Microb Ecol* 34:139–149
- Davis BD, Luger SM, Tai PC (1986) Role of ribosome degradation in the death of starved *Escherichia coli* cells. *J Bacteriol* 166:439–445
- Delbès C, Godon JJ, Moletta R (1998) 16S rDNA sequence diversity of a culture-accessible part of an anaerobic digester bacterial community. *Anaerobe* 4:267–275
- DeLong EF, Wickham GS, Pace NR (1989) Phylogenetic stains: ribosomal RNA-based probes for the identification of single cells. *Science* 243:1360–1363
- Fandino LB, Riemann L, Steward GF, Azam F (2005) Population dynamics of *Cytophaga-Flavobacteria* during marine phytoplankton blooms analyzed by real-time quantitative PCR. *Aquat Microb Ecol* 40:251–257
- Fegatella F, Lim J, Kjelleberg S, Cavicchioli R (1998) Implications of rRNA operon copy number and ribosome content in the marine oligotrophic ultramicrobacterium *Sphingomonas* sp. strain RB2256. *Appl Environ Microbiol* 64:4433–4438
- Ferrera I, Massana R, Casamayor EO, Balagué V, Sánchez O, Pedrós-Alió C, Mas J (2004) High-diversity biofilm for the oxidation of sulfide-containing effluents. *Appl Microbiol Biotechnol* 64:726–734
- Fuchs BM, Spring S, Teeling H, Quast C and others (2007) Characterization of a marine Gammaproteobacterium capable of aerobic anoxygenic photosynthesis. *Proc Natl Acad Sci USA* 104:2891–2896
- Fuhrman JA, Hewson I, Schwalbach MS, Steele JA, Brown MV, Naheem S (2006) Annually reoccurring bacterial communities are predictable from ocean conditions. *Proc Natl Acad Sci USA* 103:13104–13109
- Gentile G, Giuliano L, D'Auria G, Smedile F, Azzaro M, De Domenico M, Yakimov MM (2006) Study of bacterial communities in Antarctic coastal waters by a combination of 16S rRNA and 16S rDNA sequencing. *Environ Microbiol* 8:2150–2161
- Ghiglione JF, Larcher M, Lebaron P (2005) Spatial and temporal scales of variation in bacterioplankton community structure in the NW Mediterranean Sea. *Aquat Microb Ecol* 40:229–240
- Giovannoni SJ, Rappé M (2000) Evolution, diversity and molecular ecology of marine prokaryotes. In: Kirchman DL (ed) *Microbial ecology of the oceans*, Wiley Liss, New York, p 47–48
- Giovannoni SJ, Britschgi TB, Moyer CL, Field KG (1990) Genetic diversity in Sargasso Sea bacterioplankton. *Nature* 345:60–63
- Joux F, Agogue H, Obernosterer I, Dupuy C, Reinthaler T, Herndl GJ, Lebaron P (2006) Microbial community structure in the sea surface microlayer at two contrasting coastal sites in the northwestern Mediterranean Sea. *Aquat Microb Ecol* 42:91–104
- Kemp PF, Lee S, LaRoche J (1993) Estimating the growth rate of slowly growing marine bacteria from RNA content. *Appl Environ Microbiol* 59:2594–2601
- Kerkhof L, Ward BB (1993) Comparison of nucleic acid hybridization and fluorometry for measurement of the relationship between RNA/DNA ratio and growth rate in a marine bacterium. *Appl Environ Microbiol* 59:1303–1309
- Klepac-Ceraj V, Ceraj I, Polz MF (2006). Clusterer: extendable java application for sequence grouping and cluster analyses. *Online J Bioinform* 7:15–21
- Lebaron P, Parthuisot N, Catala P (1998) Comparison of blue nucleic acid dyes for flow cytometric enumeration of bacteria in aquatic systems. *Appl Environ Microbiol* 64:1725–1730
- Lee S, Kemp PF (1994) Single-cell RNA content of natural marine planktonic bacteria measured by hybridization with multiple 16S rRNA-targeted fluorescent probes. *Limnol Oceanogr* 39:869–879
- Lee DH, Zo YG, Kim SJ (1996) Nonradioactive method to study genetic profiles of natural bacterial communities by PCR-single-strand-conformation polymorphism. *Appl Environ Microbiol* 62:3112–3120
- Legendre L, Legendre P (1983) *Numerical ecology*. Elsevier Scientific Publishing, Amsterdam
- Lemée R, Rochelle-Newall E, Van Wambeke F, Pizay MD, Rinaldi P, Gattuso JP (2002) Seasonal variation of bacterial production, respiration and growth efficiency in the open NW Mediterranean Sea. *Aquat Microb Ecol* 29:227–237
- Mary I, Cummings DG, Biegala IC, Burkill PH, Archer SD, Zubkov MV (2006) Seasonal dynamics of bacterioplankton community structure at a coastal station in the western English channel. *Aquat Microb Ecol* 42:119–126
- Moeseneder MM, Winter C, Herndl GJ (2001) Horizontal and vertical complexity of attached and free-living bacteria of the eastern Mediterranean Sea, determined by 16S rDNA and 16S rRNA fingerprints. *Limnol Oceanogr* 46:95–107
- Moeseneder MM, Arrieta JM, Herndl GJ (2005) A comparison of DNA- and RNA-based clone libraries from the same marine bacterioplankton community. *FEMS Microbiol Ecol* 51:341–352
- Moyer CL, Mortita RY (1989) Effect of growth rate and starvation-survival on cellular DNA, RNA, and protein of a psychrophilic marine bacterium. *Appl Environ Microbiol* 55:2710–2716
- Murray AE, Preston CM, Massana R, Taylor LT, Blakis A, Wu K, DeLong EF (1998) Seasonal and spatial variability of bacterial and archaeal assemblages in coastal waters near Anvers Island, Antarctica. *Appl Environ Microbiol* 64:2585–2595
- Muyzer G, de Waal EC, Uitterlinden AG (1993) Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes encoding for 16S rRNA. *Appl Environ Microbiol* 59:695–700

- Pinhassi J, Simó R, González JM, Vila M and others (2005) Dimethylsulfoniopropionate turnover is linked to the composition and dynamics of the bacterioplankton assemblage during a microcosm phytoplankton bloom. *Appl Environ Microbiol* 71:7650–7660
- Ramette A (2007) Multivariate analyses in microbial ecology. *FEMS Microbiol Ecol* 62:142–160
- Reinthal T, Herndl GJ (2005) Seasonal dynamics of bacterial growth efficiencies in relation to phytoplankton in the southern North Sea. *Aquat Microb Ecol* 39:7–16
- Schauer M, Balagué V, Pedrós-Alió C, Massana R (2003) Seasonal changes in the taxonomic composition of bacterioplankton in a coastal oligotrophic system. *Aquat Microb Ecol* 31:163–174
- ter Braak CJF (1987) CANOCO: a FORTRAN program for canonical community ordination by [partial] [detrended] [canonical] correspondence analysis, principal components analysis and redundancy analysis. Agricultural Mathematics Group, Wageningen
- Troussellier M, Schäfer H, Batailler N, Bernard L and others (2002) Bacterial activity and genetic richness along an estuarine gradient (Rhône River plume, France). *Aquat Microb Ecol* 28:13–24
- West NJ, Obernosterer I, Zemb O, Lebaron P (2008) Major differences of bacterial diversity and activity inside and outside of a natural iron-fertilized phytoplankton bloom in the Southern Ocean. *Environ Microbiol* 10:738–756
- Zubkov MV, Fuchs BM, Archer SD, Kiene RP, Amann R, Burkill PH (2001) Linking the composition of bacterioplankton to rapid turnover of dissolved dimethylsulfoniopropionate in an algal bloom in the North Sea. *Environ Microbiol* 3:304–311
- Zumstein E, Moletta R, Godon JJ (2000) Examination of two years of community dynamics in an anaerobic bioreactor using fluorescence polymerase chain reaction (PCR) single-strand conformation polymorphism analysis. *Environ Microbiol* 2:69–78

*Editorial responsibility: Jed Fuhrman,
Los Angeles, California, USA*

*Submitted: May 6, 2008; Accepted: October 23, 2008
Proofs received from author(s): February 9, 2009*