

# Ribosomal DNA sequences of *Glugea anomala*, *G. stephani*, *G. americanus* and *Spraguea lophii* (Microsporidia): phylogenetic reconstruction

Cecile Pomport-Castillon<sup>1,\*</sup>, Johan F. De Jonckheere<sup>2</sup>, B. Romestand<sup>1</sup>

<sup>1</sup>Laboratoire de Parasitologie et Immunologie, Université Montpellier II, C.C. 096, Place Eugène Bataillon, 34095 Montpellier Cédex, France

<sup>2</sup>Protozoology Laboratory, Biosafety and Biotechnology, Institute of Hygiene and Epidemiology, 1050 Brussels, Belgium

**ABSTRACT:** The microsporidian species *Glugea anomala*, *G. stephani*, *G. americanus* and *Spraguea lophii* were compared by using sequence data derived from their small subunit rDNA genes which were amplified by polymerase chain reaction and directly sequenced. These sequence data and published data of *G. atherinae* were analyzed and were used to infer a phylogenetic tree. The 5 microsporidian fish parasites appeared to be closely related. The higher sequence similarities demonstrated among *G. anomala*, *G. stephani* and *G. atherinae* suggest that these 3 parasites are in fact only 1 species of *Glugea*. Moreover, the higher sequence similarities between *S. lophii* and *G. americanus* support the transfer of the latter *Glugea* species into the genus *Spraguea*.

**KEY WORDS:** Sequence · SSUrDNA · Microsporidia · *Glugea* · *Spraguea* · Taxonomy

## INTRODUCTION

Microsporidia are obligately intracellular parasites and are found in many vertebrates and invertebrates (Sprague & Vavra 1977). However, most Microsporidia are found in insects and fishes. Many species belonging to the genus *Glugea* were described in several species of teleost fishes (Canning & Lom 1986). They have been identified and described mainly on the basis of their tissue preference, morphology and developmental life cycle as demonstrated by electron microscopy (Canning & Lom 1986). Molecular analyses based on rDNA sequence data are increasingly applied to the phylum Microsporidia. The small subunit (SSU) rRNA or rDNA sequences have already been determined for many Microsporidia including *Vairimorpha necatrix* (Vossbrinck et al. 1987), *Ameson michaelis* (Zhu et al. 1993a), *Encephalitozoon cuniculi* (Zhu et al. 1993b), *Septata intestinalis* (Zhu et al. 1993c), *Enterocytozoon bieneusi* (Zhu et al. 1994) and *Glugea atherinae* (Genbank: U15987).

Recently, a phylogenetic analysis based on riboprinting of the SSU and LSU rDNA of 7 microsporidian species has shown that *Glugea anomala*, *G. stephani* and *G. atherinae* are very closely related while *G. americanus* may be more closely related to *Spraguea lophii* than to the other *Glugea* spp. (Pomport-Castillon et al. 1997a). When we consider the species description, their taxonomic status is questionable for several reasons. (1) Experimental infection of stickleback with *G. hertwigi* obtained from the smelt *Osmerus eperlanus*, carried out successfully by Weissenberg (1968), led Canning & Lom (1986) to question the validity of the species *G. anomala* and *G. hertwigi*. (2) Successful experimental transmission of *G. atherinae*, a parasite of the sand smelt *Atherina boyeri*, to the flounder *Platichthys flesus* (Mathieu-Daudé et al. 1992) and to the turbot *Scophthalmus maximus* (Leiro et al. 1993) calls the validity of this species and of *G. stephani* into question. (3) In spite of a very similar parasitism of angler fish (little white cysts in grapes in the nervous system), genus *Lophius*, 2 species have been described: *Spraguea lophii* for the European angler fish (*L. budegassa* and *L. piscatorius*) and *G. americanus*

\*E-mail: pomport@crit.univ-montp2.fr

for the American angler fish (*L. americanus*) (Loubés et al. 1979, Takvorian & Cali 1986).

In order to improve the taxonomy of microsporidian parasites of fish, in particular of the genera *Glugea* and *Spraguea*, we have sequenced the SSU rDNA of *G. anomala*, *G. stephani*, *G. americanus* and *S. lophii*. We report here the SSU rDNA sequences of these 4 Microsporidia and a phylogenetic reconstruction based on them.

## MATERIALS AND METHODS

**Fishes and Microsporidia.** The sand smelt *Atherina boyeri* Risso, 1810, host of *Glugea atherinae* (Berrebi, 1979), the stickleback *Gasterosteus aculeatus* L., 1758, host of *Glugea anomala* (Moniez, 1887), the flounder *Platichthys flesus* L., 1758, host of *Glugea stephani* (Hagenmuller, 1899), the American angler fish *Lophius americanus* Valenciennes, 1837, host of *Glugea americanus* (Takvorian and Cali, 1986) and the European angler fish *Lophius piscatorius* L., 1758 and *L. budegassa* Spinosa, 1807, hosts of *Spraguea lophii* (Doflein, 1898) were collected directly from fishermen or from a dealer. Fishes were dissected and xenomas were isolated from parasitized host organs. The spores were recovered from the cysts in a phosphate-buffered saline (PBS) solution (pH 7.2) and purified on a Percoll gradient (25, 50, 75 and 100%) (Jouvenaz 1981, Pomport-Castillon et al. 1997b). The spore suspensions obtained were stored at  $-20^{\circ}\text{C}$  in PBS.

**Microsporidian DNA isolation.** After centrifugation ( $3000 \times g$ ), the spore pellet was resuspended in a solution containing 0.4 ml buffer (0.5 M NaCl, 0.2 M Tris HCl pH 7.5, 0.01 M ethylenediamine tetraacetic acid and 1% sodium n-lauroyl sarcosine) and 0.4 g glass beads. The DNA was liberated from the spores by shaking in a Mini Bead Beater (Biospec. Product, Bartlesville, OK, USA). The aqueous phase was extracted with Tris-saturated phenol and precipitated with ethanol (Pomport-Castillon et al. 1997a).

**PCR amplification and sequencing.** The SSU rDNA was amplified using the forward primer 5'-GGTTG-

ATTCTGCCTGACGT-3' (Baker et al. 1994) and one of the reverse primers 5'-GACGGGCGGTGTGTACA-AAG-3' or 5'-GGTCCGTGTTTCAAGACGG-3' (Vossbrinck et al. 1993). Amplification conditions were 1 min at  $94^{\circ}\text{C}$ , 1.5 min at  $55^{\circ}\text{C}$  and 2 min at  $72^{\circ}\text{C}$  for 30 cycles using Taq DNA polymerase (Boehringer Mannheim, Germany) (Pomport-Castillon et al. 1997a). The initial primer and additional primers were used to sequence the SSU rDNA. Direct sequencing of a purified fragment was accomplished using the 70170 Sequenase<sup>TM</sup> PCR Product Sequencing kit (Amersham, United States Biochemicals, Cleveland, Ohio, USA). PCR and sequencing were done 4 times with different isolates to take any variability into account.

GenBank/NCBI accession numbers are: *Spraguea lophii* AF056013, *Glugea americanus* AF056014, *G. stephani* AF056015 and *G. anomala* AF056016. The *G. atherinae* sequence was obtained from GenBank (accession no. U15987).

**Phylogenetic analysis.** Base compositions of the sequenced fragments were analysed by a DNA statistics program (ESGS, France). The alignments were done using the computer program Multalign (INRA, France). Using the Clustal algorithm percent similarities and divergents were calculated and allowed to find the shortest tree built with the Neighbour Joining method (Saitou & Nei 1987, Higgins & Sharp 1988). Moreover, a phylogenetic tree was constructed using DOLLOP program (parsimony) of the PHYLIP package (Felsenstein 1989).

## RESULTS

The base composition and GC contents of the SSU rDNA sequences of *Glugea anomala*, *G. stephani*, *G. atherinae*, *G. americanus* and *Spraguea lophii* sequences are reported the Table 1. The base composition appears to be nearly identical for *G. anomala*, *G. stephani* and *G. atherinae* with a small bias to G+C bases, whereas the 2 other species have a bias to A+T bases. The SSU rDNA shows little variation among the taxa analyzed. The *G. anomala* and *G. stephani* sequences are identical and only 1 base was different between these 2 and the *G. atherinae* sequence. By contrast, about 160 bases were different between these 3 sequences and the *G. americanus* and *S. lophii* sequences. The *G. americanus* and *S. lophii* sequences were very similar with only 7 base differences between the 2 sequences (Table 2).

Table 2 shows the percent divergence among the 5 species. The dis-

Table 1. Base composition of the SSU rDNA sequences of 5 species of Microsporidia. S: *Spraguea*; G: *Glugea*

	<i>S. lophii</i>	<i>G. americanus</i>	<i>G. anomala</i>	<i>G. stephani</i>	<i>G. atherinae</i>
No. of bases	1175	1174	1165	1165	1165
A (%)	30.6	30.5	29.4	29.4	29.5
G (%)	31.6	31.8	32.4	32.4	32.3
T (%)	20.1	20.1	19.0	19.0	19.0
C (%)	17.7	17.6	19.2	19.2	19.2
C + G (%)	49.3	49.4	51.6	51.6	51.5

Table 2. Percent sequence divergence among 5 species of Microsporidia. S: *Spraguea*; G: *Glugea*

	<i>S. lophii</i>	<i>G. americanus</i>	<i>G. anomala</i>	<i>G. stephani</i>
<i>S. lophii</i>	0.0			
<i>G. americanus</i>	0.6			
<i>G. anomala</i>	16.0	15.8		
<i>G. stephani</i>	16.0	15.8	0.0	
<i>G. atherinae</i>	15.9	15.7	0.1	0.1

tances between *Glugea atherinae* and *G. anomala*/*G. stephani* and between *G. americanus* and *Spraguea lophii* are smaller (0.1 and 0.6%, respectively) than that between *G. atherinae* and *G. americanus*/*S. lophii* (15.7/15.9%).

Fig. 1 illustrates a phylogenetic tree of the 5 species showing 2 clusters, i.e. the first grouping: *Glugea anomala*, *G. stephani* and *G. atherinae*, and the second grouping: *G. americanus* and *Spraguea lophii*. Such a tree supports the exclusion of *G. americanus* from the genus *Glugea*. Moreover, the analysis of the SSU rDNA sequences using the DOLLOP programs of the PHYLIP package generated a tree showing the phylogenetic relationship between the different fish Microsporidia (Fig. 2).

## DISCUSSION

Using conserved primers in the SSU rDNA, we obtained sequences of 4 microsporidian species belonging to the genera *Glugea* and *Spraguea* (*G. anomala*, *G. stephani*, *G. americanus* and *S. lophii*) and have compared these with a sequence of another species of the *Glugea* genus, *G. atherinae*.




Fig. 1. Phylogenetic tree of the 5 species of Microsporidia based on the percent divergence. G: *Glugea*; S: *Spraguea*

Our study and some recent riboprinting experiments on several microsporidian species (Pomport-Castillon et al. 1997a) show the following.

The SSU rDNA sequences of the 5 species are closely related since the lowest similarity is about 80.3% as observed between *Glugea anomala*, *G. stephani* and *G. americanus*. Microsporidian fish parasites constitute a homogeneous group in comparison with other species. Malone & McIvor (1995) have

shown a great sequence variability between species of the *Nosema* genus parasitising different classes of arthropods: 62.4% similarity was observed between *N. bombycis* and *N. apis*.

The analysis of the base composition outlines a linking of some of these species. Pieniazek et al. (1995) have shown a variation of the GC content of 15 microsporidian species with, at the extreme, 34% for *Nosema bombycis* and 56% for *Vavraia oncoperae*. However, in the same genus the GC content is very similar, e.g. 34 to 38% for *Nosema* spp. Malone & McIvor (1995) have demonstrated that, mainly in Protozoa, the ratio G+C/A+T is less than or equal to 1. This is consistent for *Glugea americanus* and *Spraguea lophii*, but not for the 3 other species which show a majority of G+C bases.

*Glugea anomala*, *G. stephani* and *G. atherinae* are very closely related while *G. americanus* appears to be more closely related to *Spraguea lophii* than to the other 3 *Glugea* species. Indeed, the *G. anomala* and *G. stephani* sequences are identical or almost identical (99.9% similarity) to the *G. atherinae* sequence. This result is in accordance, for these 3 species, with a well established homogenous life cycle inducing voluminous xenomas encapsulated in connective tissues of their host fishes. In spite of some morphological differ-




Fig. 2. Phylogenetic tree of the 5 species of Microsporidia based on SSU rDNA sequence using the Dollop program. Bootstrap values are 100% or close to this for all branching node. G: *Glugea*; S: *Spraguea*

ences in the spores, experimental infections have been carried out successfully between some of these species (Mathieu-Daudé et al. 1992, Leiro et al. 1993). These observations and our results suggest that *G. anomala*, *G. stephani* and *G. atherinae* could be only 1 species. However, there is evidence that some Microsporidia are cross-infective between hosts and that the morphology can be altered in different hosts (Canning & Lom 1986). Moreover, Docker et al. (1997) have demonstrated the existence of genetic variability in the rDNA sequences of another microsporidian parasites in the same host species. Indeed, the authors have sequenced 2 samples of *Loma salmonae* and have shown differences of 2 nucleotides. Such intra-species variability can explain the difference observed between *G. anomala*, *G. stephani* and *G. atherinae* by riboprinting and sequencing, and could be an argument in favour of the existence of only one species.

*Glugea americanus* and *Spraguea lophii*, in spite of causing a very similar parasitism of the American and the European angler fishes' nervous systems respectively, produce different spores. *G. americanus* has oval and uninucleated spores of  $2.8 \times 1.5 \mu\text{m}$  while *S. lophii* has 2 types of spores: oval, uninucleated ones and curved, binucleated ones with a size of  $4.2 \times 2.5 \mu\text{m}$  and  $3.7 \times 1.4 \mu\text{m}$ , respectively (Loubés et al. 1979, Lom & Dykova 1992). However, it is important to mention that the early proliferative forms and the complete development of *G. americanus* are unknown and only spores and xenomas have been described (Takvorian & Cali 1986). In our study, these 2 angler fish parasite species have a higher SSU rDNA sequence similarity (98.6%) which is due to only 2 transitions, 2 transversions and 1 deletion. The closely related sequences determined here confirm the result obtained by riboprinting and could support the transfer of *G. americanus* into the genus *Spraguea*. Indeed, percent similarities and divergence have already been used to rename species. Zhu et al. (1993c) and Hartskeel et al. (1995) have justified the transfer of *Septata intestinalis* to the genus *Encephalitozoon* by a lower divergence (21.9%) and a high similarity (70%) between *S. intestinalis* and *Encephalitozoon cuniculi*. However, it would be of interest to know how many nucleotide substitutions constitute differences between species and genera. Molecular systematics are clearly helpful when there are many nucleotide differences but slight differences create difficulties in interpretation.

The low numbers of differences between gene sequences of these species give little information. The present work is the second step, after riboprinting, in the study of the phylogeny relationships of these 5 microsporidian species and more sequence data have to be obtained. Moreover, further steps should be taken to confirm the hypotheses inferred from the mol-

ecular data: (1) an examination of the morphology of *Glugea americanus* and description of the life cycle to see whether it exhibits sporal dimorphism as is present in *Spraguea lophii*; (2) the separation of the 2 spore types of *S. lophii* and analysis of both sequences to confirm that they belong to the same species; (3) cross-transmission experiments if possible among the *G. anomala*, *G. atherinae*, *G. stephani* group; and (4) sequencing of the ITS and the LSU rDNA or another gene more variable among the 5 species under study.

#### LITERATURE CITED

- Baker MD, Vossbrinck CR, Didier ES, Maddox JV, Undeen AH (1994) Phylogenetic relationships among *Vairimorpha* and *Nosema* species (Microspora) based on ribosomal RNA sequence data. *J Invertebr Pathol* 64:100–106
- Canning EU, Lom J (1986) *The Microsporidia of vertebrates*. Academic Press, New York
- Docker MF, Devlin RH, Richard J, Khattra J, Kent ML (1997) Sensitive and specific polymerase chain reaction assay for the detection of *Loma salmonae* (Microsporea). *Dis Aquat Org* 29:41–48
- Felsenstein J (1989) PHYLIP-phylogeny inference package (version 3.2). *Cladistics* 5:164–166
- Hartskeel RA, Van Gool T, Schuitema ARJ, Didier ES, Terpstra WJ (1995) Genetic and immunological characterization of the microsporidian *Septata intestinalis* Cali. Kotler and Orenstein. 1993: reclassification to *Encephalitozoon intestinalis*. *Parasitology* 110:277–285
- Higgins DG, Sharp PM (1988) CLUSTAL: a package for performing multiple sequence alignment on a microcomputer. *Gene* 73:237–244
- Jouvenaz DP (1981) Percoll: an effective medium for cleaning microsporidian spores. *J Invertebr Pathol* 37:219
- Leiro J, Bos J, Iglesias R, Estevez J, Fernandez J, Sanmartin ML (1993) Experimental infection of turbot (*Scophthalmus maximus* L.) with a microsporean parasite (*Glugea atherinae* Berrebi, 1979) of the sand smelt (*Atherina boyeri*). *Aquaculture* 55:235–242
- Lom J, Dykova I (1992) Protozoan parasites of fishes. *Dev Aquac Fish Sci* 26:125–157
- Loubés C, Maurand J, Ormières R (1979) Étude ultrastructurale de *Spraguea lophii* (Doflein, 1898), microsporidie parasite de la baudroie: essai d'interprétation du dimorphisme sporal. *Protistologia* 15:43–54
- Malone LA, McIvor CA (1995) DNA probes for two Microsporidia. *Nosema bombycis* and *Nosema costelytrae*. *J Invertebr Pathol* 65:269–273
- Mathieu-Daudé F, Faye N, Coste F, Manier F, Marques A, Bouix G (1992) Occurrence of microsporidiosis in marine cultured gilt-head sea bream from the Languedoc coast: a problem of specificity in the genus *Glugea* (Protozoa, Microspora). *Bull Eur Assoc Fish Pathol* 12:67
- Pieniazek NJ, Da Silva AJ, Slemenda SB, Vivesvara GS, Kurtti TJ, Fries I, Yasunaga C, Ditrich O, Lom J (1995) Molecular taxonomy of microsporidia based on complete small subunit ribosomal RNA sequences and secondary structures. *Soc Invertebr Pathol 28th Annual meeting*, University Ithaca, New York
- Pomport-Castillon C, Romestand B, De Jonckheere JF, (1997a) Identification and phylogenetic relationships of Microsporidia by riboprinting. *J Eukaryot Microbiol* 44: 540–544

- Pomport-Castillon C, Gasc C, Romestand B (1997b) Development and characterization of monoclonal antibodies against spores of *Glugea atherinae* (Protozoa-Microsporidia-Fish parasites) and future prospects. *J Eukaryot Microbiol* 44:643–648
- Saitou N, Nei M (1987) The neighbour-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sprague V, Vavra J (1977) Classification and phylogeny of the Microsporidia. In: Bulla LA, Cheng TC (ed) *Comparative pathology II*. Plenum Press, New York
- Takvorian PM, Cali A (1986) The ultrastructure of spores (Protozoa: Microsporoda) from *Lophius americanus*, the angler fish. *J Protozool* 33:570–575
- Vossbrinck CR, Maddox JV, Friedman S, Debrunner-Vossbrinck BA, Woese CR (1987) Ribosomal RNA sequence suggests microsporidia are extremely ancient eukaryotes. *Nature* 326:411–414
- Vossbrinck CR, Baker MD, Didier ES, Debrunner-Vossbrinck BA, Shaddock JA (1993) Ribosomal DNA sequences of *Encephalitozoon hellem* and *Encephalitozoon cuniculi*: species identification and phylogenetic construction. *J Eukaryot Microbiol* 40:354–362
- Weissenberg R (1968) Intracellular development of the Microsporidian *Glugea anomala* Moniez in hypertrophying migratory cells of the fish *Gasterosteus aculeatus*, an example of the formation of the xenoma tumors. *J Protozool* 15:44–57
- Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM (1993a) Nucleotide sequence of the small subunit rRNA of *Ameeson michaelis*. *Nucleic Acids Res* 21:3895
- Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM (1993b) Nucleotide sequence of the small subunit rRNA of *Encephalitozoon cuniculi*. *Nucleic Acids Res* 21:1315
- Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM (1993c) Nucleotide sequence of the small subunit rRNA of *Septata intestinalis*. *Nucleic Acids Res* 21:4846
- Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM (1994) Ribosomal RNA sequences of *Enterocytozoon bieneusi*, *Septata intestinalis*, *Ameeson michaelis*: phylogenetic construction and structural correspondence. *J Eukaryot Microbiol* 41:204–209

Editorial responsibility: Wolfgang Körting,  
Hannover, Germany

Submitted: April 10, 1999; Accepted: December 9, 1999  
Proofs received from author(s): February 14, 2000