First isolation of *Brucella pinnipedialis* and detection of *Brucella* antibodies from bearded seals *Erignathus barbatus*

Geoffrey Foster^{1,*}, Ingebjørg H. Nymo^{2,7}, Kit M. Kovacs³, Kimberlee B. Beckmen⁴, Andrew C. Brownlow¹, Johanna L. Baily^{5,8}, Mark P. Dagleish⁵, Jakub Muchowski⁶, Lorraine L. Perrett⁶, Morten Tryland², Christian Lydersen³, Jacques Godfroid², Barry McGovern^{1,9}, Adrian M. Whatmore⁶

¹SAC Consulting Veterinary Services, Drummondhill, Stratherrick Road, Inverness IV2 4JZ, UK
²Arctic Infection Biology, Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Postboks 6050 Langnes, 9037 Tromsø, Norway
³Norwegian Polar Institute, Fram Centre, 9296 Tromsø, Norway
⁴Alaska Department of Fish and Game, 1300 College Road, Fairbanks, AK 99701-1551, USA
⁵Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik, EH26 0PZ, UK
⁶APHA Weybridge, Woodham Lane, Addlestone, Surrey KT15 3NB, UK

⁷Present address: Norwegian Veterinary Institute, Stakkevollveien 23, 9010 Tromsø, Norway

⁸Present address: Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK

⁹Present address: Cetacean Ecology and Acoustics Laboratory, School of Veterinary Science, University of Queensland, QLD 4343, Australia

ABSTRACT: *Brucella* species infecting marine mammals was first reported in 1994 and in the years since has been documented in various species of pinnipeds and cetaceans. While these reports have included species that inhabit Arctic waters, the few available studies on bearded seals *Erignathus barbatus* have failed to detect *Brucella* infection to date. We report the first isolation of *Brucella pinnipedialis* from a bearded seal. The isolate was recovered from the mesenteric lymph node of a bearded seal that stranded in Scotland and typed as ST24, a sequence type associated typically with pinnipeds. Furthermore, serological studies of free-ranging bearded seals in their native waters detected antibodies to *Brucella* in seals from the Chukchi Sea (1990–2011; 19%) and Svalbard (1995–2007; 8%), whereas no antibodies were detected in bearded seals from the Bering Sea or Bering Strait or from captive bearded seals.

KEY WORDS: Antibodies \cdot Bearded seal \cdot Brucella pinnipedialis \cdot Isolation \cdot Multilocus sequence typing \cdot MLST

- Resale or republication not permitted without written consent of the publisher

INTRODUCTION

The isolation of *Brucella* from marine mammals was first reported in 1994 from 4 free-ranging harbour seals *Phoca vitulina*, 2 harbour porpoises *Phocoena phocoena* and a common dolphin *Delphinus delphis*, all inhabiting Scottish coastal waters (Ross et al. 1994), and from an aborted foetus born to a captive bottlenose dolphin *Tursiops truncatus* in the USA (Ewalt et al. 1994). Since these initial reports, *Brucella* infection has become recognised in cetaceans and pinnipeds inhabiting many of the world's oceans (Foster et al. 2002, Nymo et al. 2011), and 2 species, *Brucella ceti* and *B. pinnipedialis*, have been described for isolates with cetaceans and seals as preferred hosts, respectively (Foster et al. 2007). These species are genetically distinct from *Brucella* associated with terrestrial mammals (Whatmore et al. 2016).

With respect to Scottish coastal waters, B. pinnipedialis has been recovered from the other resident seal species, grey seals Halichoerus grypus, as well as from hooded seals Cystophora cristata, which are occasional visitors to the region (Foster et al. 1996, 2002). The isolation of *B. pinnipedialis* has also been reported from hooded seals in their native Arctic waters and from harbour and grey seals elsewhere in Europe (Nymo et al. 2011). Further afield, B. pinnipedialis has been cultured from other pinniped species including Pacific harbour seal Phoca vitulina richardsi (Garner et al. 1997), ringed seal Pusa hispida, harp seal Pagophilus groenlandica (Forbes et al. 2000) and California sea lion Zalophus californianus (Goldstein et al. 2009). Serological studies provide further presumptive evidence that Brucella infections are widespread among other pinniped species, including some resident in the Southern Hemisphere (Nymo et al. 2011). Taken together, culture and serological evidence (Foster et al. 2002, Nymo et al. 2011) indicate that Brucella is endemic in many of the marine mammals that inhabit the world's open oceans and seas. Seropositive animals, however, can be due to immunological cross-reactions, in particular Yersinia enterocolitica serotype O9, although this strain has not been recovered from marine mammals to date (G. Foster pers. obs.) or an organism from a different genus; thus, the isolation of Brucella by cultural methods remains the gold standard of definitive proof of infection in different hosts and discrete populations of marine mammals.

There have been few reports on studies of Bru*cella* infection in bearded seals *Erignathus barbatus* to date, but where performed, no evidence of exposure was found (Tryland et al. 1999, Calle et al. 2008). Bearded seals are members of the Phocidae family and represent the only species within the genus *Erignathus*. They have a patchy circumpolar distribution throughout the Arctic and Subarctic between 45° and 85° N. Two subspecies are recognised, E. barbatus barbatus, which ranges from the central Canadian Arctic eastwards to the central Eurasian Arctic, and E. barbatus nauticus, which ranges from the central Canadian Arctic westwards to the Laptev Sea, Russia. The availability of sea ice to breed, moult and rest on in shallow water areas is thought to be an important factor governing the distribution of this benthic-feeding seal (Kovacs 2016).

In a review of their extralimital records, bearded seals have been reported from the Netherlands, France and Spain in the eastern Atlantic and the island of Rügen in the Baltic Sea (van Bree 2000). Sightings in the UK are rare, with most modern reports occurring around the Scottish coast, including the Shetland and Orkney islands and single sightings from the Isle of Mull, Aberdeenshire and Fife (JNCC/Defra 2013).

This paper documents the first recovery and characterisation of *B. pinnipedialis* from a bearded seal. The results of a serological study of free-living bearded seals in Arctic waters and captive members of the species kept at the Polaria aquarium in Tromsø, Norway, are also presented.

MATERIALS AND METHODS

Bearded seal necropsy

In early February 2012, a stranded bearded seal (M61/12) was reported to the Scottish Marine Animal Strandings Scheme. The juvenile male animal had stranded dead at Annachie Lagoon, St. Fergus, on the Aberdeenshire coast of the northeastern Scottish mainland (57° 34′ 10.74″ N, 1° 49′ 22.02″ W) and represented the first report of a stranded bearded seal in Scotland since records began in 1992. The carcase was transported to SAC Consulting Veterinary Services, Inverness, for a postmortem examination performed according to a standard protocol (Dierauf 1994). Samples of brain, lung, liver, spleen, kidney, mesenteric lymph node, urinary bladder and small intestine were cultured on Columbia sheep blood agar (CSBA) (Oxoid) and Farrell's medium (FM) (Farrell 1974) and incubated at 37°C in air with 5% added CO_2 as described previously (Foster et al. 2002). Plates were examined for growth daily for 4 d and at frequent intervals thereafter up to 14 d. Isolates with colonial appearance typical of Brucella were tested initially for Gram reaction, cellular morphology, acid fastness with the modified Ziehl-Neelsen stain, agglutination with Brucella abortus antiserum (Remel) and ability to grow in air without added CO₂. Further testing included urea hydrolysis, H₂S production, inhibition by basic fuchsin at 1:50000 and 1:100 000, agglutination with monospecific antisera A and M and lysis by phages TB, Wb, BK2, Fi, Iz and R/C all at routine testing dose. Multilocus sequence typing (MLST) using a 9 locus scheme was performed as described previously (Whatmore

et al. 2007). Tissue samples for histological examination (whole brain, trigeminal ganglion, skin, thyroid gland, adrenal gland, urinary bladder, spleen, lung, kidney, heart and pancreas) were collected, trimmed and processed routinely through graded alcohols and embedded in paraffin wax prior to sectioning (5 μ m), mounting on glass microscope slides and staining with haematoxylin and eosin. Blood was collected from the left ventricle of the heart using a sterile needle into heparinised and plain 6 ml vacutainers (BD) for serology, and urine analysis was performed using the Combur 9 test (Roche).

Serology

The Alaska Department of Fish and Game ice seal program recovered serum from heart blood samples from subsistence-harvested bearded seals. Seals were shot on sea ice by Alaska Native hunters, as allowed under the Marine Mammal Protection Act of 1972, in the Chukchi Sea and Bering Strait off the northern and northwestern coasts of Alaska during May, June, July and October. In addition, 17 seals were sampled immediately post mortem during 1978–1979 scientific collections conducted April to June by the Outer Continental Shelf Environmental Assessment Program during NOAA cruises in the Bering Sea under National Marine Fisheries permit no. 194. Samples from bearded seals from Svalbard were obtained from both dead and live animals. Sixteen animals were shot on the ice as part of Norwegian scientific catches to address stocks and diets for different seal species and their role in the marine ecosystems and interactions with fisheries (1992-1995). Blood was collected on the ice when cutting the main blood vessels to the foreflippers during standard bleeding-out procedures of seals (Tryland et al. 1999). From live bearded seals (pups; 1995–2007), blood was obtained from the extradural intra-vertebral vein using an 80 mm needle (14 gauge, 2.1 mm) mounted on a 50 ml syringe. Blood was transferred into blood-collecting tubes (Venoject, Terumo), and serum was prepared ($3000 \times g$, 15 min) and stored at -20° C until analysis. Sex and age category (pup: <1 yr, juvenile: <3 yr, adult: >3 yr) were known for some or all of the seals at each location (Table 1).

Furthermore, blood samples were obtained from 5 bearded seals kept in captivity at the Polaria aquarium in Tromsø. Blood was drawn from the plantar venous plexus of the hind flippers, using a 0.8×50 mm needle and blood collecting tubes (Venoject). Serum was prepared by centrifugation at $3000 \times g$ for 15 min and stored at -20° C until analysis. These animals, initially captured in the wild in Svalbard, had been kept in captivity since they were approximately 5 wk of age; the seals interact extensively with humans through training and feeding (Stokke 2010). They were 9 to 10 yr of age at the time of sampling and had been trained to tolerate handling and blood sampling (Table 1).

Serum samples (n = 205) were analyzed for anti-Brucella antibodies with a protein A/G indirect enzyme-linked immunosorbent assay as described previously (Nymo et al. 2013a). The mean optical density (OD) of duplicate wells was expressed as a percentage of the reactivity of a seal positive control: (OD sample/OD positive control) × 100 = percent positivity (%P). The cut-off was 73.6 %P.

Statistical analysis

All statistical analysis was performed in JMP 11 Pro (SAS Institute).

Table 1. Bearded seals investigated for anti-*Brucella* antibodies with a protein A/G indirect enzyme-linked immunosorbent assay. Sampling location, sampling years, sex and age category are given. Data are number of seropositives from total tested for each category, with percentages in parentheses. Pup: <1 yr; juvenile: <3 yr; adult: >3 yr; (–) no data

Location	Years		Sex			Age category —				
		Female	Male	Unknown	Pup	Juvenile	Adult	Unknown	Total	
Bering Sea	1978-2005	0/8 (0)	0/15 (0)	0/1 (0)	0/3 (0)	0/15 (0)	0/5 (0)	0/1 (0)	0/24 (0)	
Bering Strait	2001-2005	0/6 (0)	0/7 (0)	0/1 (0)	0/6 (0)	0/1 (0)	0/6 (0)	0/1 (0)	0/14 (0)	
Chukchi Sea	1990-2011	10/49 (20)	4/31 (13)	2/6 (33)	0/7 (0)	3/12 (25)	5/24 (21)	8/43 (19)	16/86 (19)	
Svalbard	1995-2008	4/29 (14)	2/47 (4)	_	5/67 (7)	_	1/9 (11)	_	6/76 (8)	
Captive	2006-2008	0/3 (0)	0/2 (0)	_	_	_	0/5 (0)	_	0/5 (0)	
Total	1978-2011	14/95 (15)	6/102 (6)	2/8 (25)	5/83 (6)	3/28 (11)	6/49 (12)	8/45 (18)	22/205 (11)	

RESULTS

Bearded seal necropsy

The carcase of M61/12 was fresh and had been chilled, but not frozen, prior to necropsy 2 d after notification. The animal was 149 cm in total length, 79 cm in girth behind the front flippers and in moderate to poor body condition, with a mid-sternal blubber thickness of 16 mm.

Hair loss ranged from complete over the ventral surface of the animal to partial coverage over the flank, with bilateral symmetrical zones of alopecia, back to almost complete hair loss over the perilumbar region. The head and neck exhibited almost full coverage, excepting significant alopecia periocularly and over the dorsal muzzle. In addition, the foreflippers exhibited partial alopecia over the carpal and phylangeal regions. No regions showed evidence for hair regrowth.

The oesophagus and stomach contained a notable amount of sand, and marine debris comprising a fragment of worn black plastic sheeting 45 mm long and a single round pebble 1 cm in diameter were recovered from the stomach. No prey items were found. Thyroid glands were grossly unremarkable. The lungs and cerebral vessels were markedly congested, and the bladder mucosa was grossly reddened; the urine was turbid and dark red in colour, and a high level of haemoglobin (ca 50 erythrocytes per μ l) was detected with the Combur 9 test. The brain showed diffuse dilation of cerebral vessels, but the cerebrospinal fluid was unremarkable.

Bacteriology

Small numbers of colonies typical of Brucella were recovered from the mesenteric lymph node on CSBA and FM after 4 d. In addition, Vibrio algi*nolyticus* was recovered from multiple tissues. Cells of suspect Brucella colonies were tiny Gram-negative coccobacilli, which were acid fast when tested in the modified Ziehl-Neelsen stain. Agglutination was obtained in slide tests with Brucella abortus antiserum. The strain required CO₂ for growth and was urease positive, H₂S negative and A dominant. Growth was inhibited by basic fuchsin at 1:50000 and 1:100000, and cultures were completely lysed by Tb phage and partially lysed by Wb, BK2 and Iz, with no lytic effect with Fi and R/C. The strain was identified by MLST as B. pinnipedialis sequence type (ST) 24.

Histopathology

The most significant histological change in M61/12 consisted of moderate multifocal granulomatous and eosinophilic meningoencephalitis within the brain, often centred on degenerate or intact nematode parasite larvae, with perivascular cuffing and multifocal haemorrhages. The nematode larvae were not identified, but gross morphology of worms seen in the stomach was consistent with anisakid nematodes Pseudoterranova bulbosa or Contracaecum osculatum. It is plausible that the granulomatous foci in the brain were the result of aberrant tissue migration of L4 larval stages from these species. Mild multifocal histiocytic and eosinophilic pneumonia (likely parasitic) was also noted along with moderate splenic histiocytosis with mild lymphodepletion. The skin lesions consisted of mild epidermal hyperplasia with follicular atrophy with no evidence of vasculitis or dermal necrosis. Moderate to marked thyroid follicular hyperplasia and moderate to marked bilateral adrenocortical hyperplasia were present. The most significant lesions, and likely cause of death, were multiple granulomatous foci in many regions of the brain consistent with migrating nematode larvae. Overall, the seal appeared to have indications of chronic morbidity and malnutrition/pica which, given the extralimital nature of this case, could be due to pathogen exposure and/or inadequate feeding capacity.

Serology

Antibodies to Brucella were detected in 22 of 200 (11%) serum samples collected from wild bearded seals in Alaska and Svalbard (Table 1). Sixteen of the seropositive seals came from 86 (19%) animals that were subsistence harvested in the Chukchi Sea between 1990 and 2011: 1 juvenile female, 2 juvenile males, 4 adult females, 1 adult male, 5 females of unknown age, 1 male of unknown age and 2 animals of unknown sex and age (Table 1). The other seropositive bearded seals, 6 of 76 (8%), were all captured in the Svalbard archipelago during the period 1995 to 2007. The positive animals were 3 female and 2 male pups and the mother of one of the female seropositive pups. It is not known whether the mothers of the other seropositive pups were among the animals sampled. Antibodies to Brucella were not detected from any of the 38 bearded seals subsistence harvested in the Bering Strait or collected in the Bering Sea or from the 5 animals kept in captivity

at Polaria (Table 1). *Brucella* antibodies were detected in the blood collected from the necropsied animal (M61/12).

The average %P of the seropositive seals was 87.7 %P (SD 9.4), and the average %P of the seronegative seals was 25.9 %P (SD 21.6). The material included 6 mother–pup pairs from Svalbard sampled between 6 and 25 May 1995. One mother–pup pair was seropositive (mother: 93.2 %P, pup: 88.8 %P). The remaining pairs had %P values below the cutoff; however, the %P of the pup could be predicted from the %P of the mother with the following formula: pup = $-75.8 + (1.7 \times \text{mother})$, r² = 0.84.

DISCUSSION

This study documents for the first time the recovery of Brucella from a bearded seal as well as the first serological evidence of Brucella exposure in this host. Antibodies were detected in sera from 2 of the 4 groups of free-ranging bearded seals sampled, the Chukchi Sea (19%) and the Svalbard archipelago (8%); however, they were not detected from 38 bearded seals from the Bering Strait region or the Bering Sea (Table 1). A previous small-scale study also failed to detect Brucella antibodies from 6 bearded seals taken during a subsistence hunt at St. Lawrence Island in the Bering Sea (Calle et al. 2008), so evidence of exposure to *Brucella* in this region remains lacking (Fig. 1). The Pacific bearded seals are not distinct populations; they move from the Bering Sea through the Bering Strait with the advancing and retreating ice edges. The detection of seropositive bearded seals from the Chukchi Sea therefore may be significant for *Erignathus barbatus* nauticus across their entire area. Another serological study for Brucella in bearded seals did not detect antibodies from 2 locations in the North Atlantic, while antibodies were detected in the other 3 sympatric species sampled, hooded, harp and ringed seals (Tryland et al. 1999).

Typing of the *Brucella* isolate by MLST (Whatmore et al. 2007) demonstrated that it belonged to the ST24 lineage of *Brucella pinnipedialis*. ST24 is the less common of 2 STs isolated predominantly from pinnipeds (Groussaud et al. 2007) and has previously been found associated with harbour seals, grey seals and a minke whale *Balaenoptera acutorostrata* which stranded in Scotland and from harbour seals and a beluga whale *Delphinapterus leucas* from North America (Groussaud et al. 2007, Whatmore et al. 2017).

Brucella-associated pathology was not found either grossly or histologically, although histology was not performed on the lymph node, and an association of *B. pinnipedialis* with the death of this animal was not established. This is in line with previous findings, which have revealed a paucity of pathologies following Brucella isolation from pinnipeds, including several apparently healthy harbour seals which had been shot by fishermen (Foster et al. 2002). In contrast, a broad range of pathologies have been reported for Brucella infection of various cetacean species which include lymphocytic meningoencephalitis, subcutaneous lesions, blubber abscessation, liver abscess, hepatic and splenic necrosis, macrophage infiltration in liver and spleen, lymph node inflammation, pneumonia, peritonitis, mastitis, osteomyelitis, spinal discospondylitis, diseased atlanto-occipital joint, endocarditis, epididymitis and abortion (Foster et al. 2002, Nymo et al. 2011, Guzman-Verri et al. 2012).

In vitro work has revealed differences between the classical terrestrial Brucella strains and B. pinnipedialis. The B. pinnipedialis reference strain NCTC 12890 and B. pinnipedialis hooded seal strains were eliminated from murine and human macrophage cell lines and a human epithelial cell line within 72 to 96 h (Larsen et al. 2013b). Even more rapid elimination patterns were observed in hooded seal primary alveolar macrophages (Larsen et al. 2013a) and epithelial cells (Larsen et al. 2016). B. pinnipedialis NCTC 12890 was also found to be attenuated in the BALB/c Brucella mouse model (Nymo et al. 2016). The reduced virulence in these models, when compared to the terrestrial virulent strain Brucella suis 1330 (Larsen et al. 2013b, Nymo et al. 2016), is in line with the limited virulence of the *B. pinnipedialis* strains in their natural hosts (Foster et al. 2002). Five seropositive pups were detected in the present study, all from Svalbard. The sampling took place in May, and peak birthing for bearded seals at Svalbard is in early May. The pups are thereafter weaned in approximately 24 d (Kovacs 2016). The seropositive pups were hence of very young age. At least one of the seropositive pups in the present study was the pup of a seropositive mother, and a strong relationship was identified between the titres (i.e. antibody levels) in the mothers and the pups in the 6 motherpup pairs from Svalbard. These findings suggest a transfer of maternal antibodies between mother and pup. Seals have an endotheliochorial placenta (Stewart & Stewart 2009) where solely 5 to 10% of the maternal antibodies are transferred to the fetus in utero. Passive immunity via the colostrum is therefore essential in species in which the type of placentation impedes contact between maternal and foetal circulation systems, hindering the transfer of antibodies (Tizard 2000). Indeed, when evaluating total IgG levels in harbour seal mothers and pups, the mothers showed a decreasing trend during lactation, while the total IgG levels in the pups were low at birth and higher at the end of lactation (Ross et al. 1993). In dogs, which also have an endotheliochorial placenta (Stewart & Stewart 2009), pups from Brucella canis-infected bitches have antibodies against B. canis (Carmichael & Kenney 1970). In animals having an epitheliochorial placenta, the young are born virtually agammaglobulinemic; however, after receiving colostrum, calves from dams with high and low Brucella antibody levels had corresponding high and low Brucella antibody levels (Sutherland & Searson 1990). Maternal transfer of antibodies against phocine distemper virus has been shown in harbour seals (Garnier et al. 2014), and maternal and pup antibody titres were shown to be strongly correlated in Scottish grey seals (Pomeroy et al. 2005). Even though the number of mother-pup pairs investigated in the present study is low, our findings and the literature support that the Brucella antibody levels of the pups are likely a reflection of the Brucella antibody levels of the mothers and the result of maternal transfer of antibodies rather than a vertical infection of the pups. However, further bacteriology work on organ samples from mother and pup pairs is needed to draw any conclusions on this matter. For hooded seals, however, no relation was found between Brucella serostatus and ovulation rate or neonatal body condition (Nymo et al. 2013b).

Bearded seals are largely solitary animals (Kovacs 2016). Ringed seals and hooded seals, from which B. pinnipedialis has been isolated and anti-Brucella antibodies detected (Forbes et al. 2000, Nymo et al. 2013b), are also generally described as being largely solitary (Kovacs 2002, Miyazaki 2002), though all 3 of these species do gather in the same areas where habitat is suitable for breeding, moulting and foraging. Contrary to cetaceans, where vertical transmission of Brucella was suggested (Ohishi et al. 2016), no evidence of vertical transmission of Brucella in true seals has been reported. Furthermore, the solitary behaviour of bearded seals suggests that opportunities for Brucella transmission between conspecifics are restricted. Altogether, this reinforces the possibility that Brucella infection may be acquired from the environment, possibly via diet, as suggested previously (Lambourn et al. 2013, Nymo et al. 2013b). In contrast, harp seals have also been shown to harbour infections with *B. pinnipedialis* (Tryland et al. 1999, Forbes et al. 2000), but this species demonstrates a much stronger tendency to congregate (Lavigne 2002), and transmission between conspecifics cannot be excluded.

Brucellosis is a significant zoonotic infection which causes a broad range of manifestations; it is especially associated with farmed animals and their products infected with Brucella melitensis, B. abortus and B. suis but also with B. canis contracted from dogs. While there have been 3 reports of human infections with marine mammal Brucella, none have involved B. pinnipedialis. Human infection has been reported in a laboratory infection scenario with ST23, a clade predominantly associated with porpoises, while naturally occurring infections have been reported only with ST27 (Whatmore et al. 2008), only isolated thus far from bottlenose dolphins Tursiops truncatus and California sea lions in the USA (Whatmore et al. 2017) and recently from a single bottlenose dolphin in the Mediterranean (Cvetnić et al. 2016).

While the lack of human infections with *B. pinnipedialis* is in contrast to the findings with *B. ceti* and the classical *Brucella* spp. mentioned in the previous paragraph, the zoonotic potential of *B. pinnipedialis* remains unknown at present. It is advisable, therefore, that those working with bearded seals and other pinniped species consider the infectious nature of the genus and follow appropriate safety procedures (Dierauf & Gulland 2001).

In conclusion, we report the first isolation of *B. pinnipedialis* from a stranded extralimital juvenile male bearded seal. In contrast to cetaceans, reports of *Brucella*-associated pathology in pinnipeds are lacking. Our study also provides novel serological evidence for *Brucella* spp. exposure in free-ranging bearded seal subpopulations. Future serological surveys and the isolation and characterization of *Brucella* isolates from stranded and free-ranging bearded seals, as well as other ice seals (ringed seal, ribbon seal, spotted seal), are needed to better understand the significance of *Brucella* infection in these northern pinnipeds.

Acknowledgements. The authors thank Eva Marie Breines and Ellinor Hareide at UiTThe Arctic University of Norway, Research Group for Arctic Infection Biology, for their excellent laboratory work. We acknowledge the contributions of the Alaska Native hunters who participated in the biosampling programs. Thanks are also given to the many Alaska Department of Fish and Game biologists, John J. Burns and Frances Fay (deceased), who collected and archived the sera samples and associated data, and Nathan Pamperin for the map. We also thank Espen Rafter and other staff members at Polaria for their interest and patience regarding sampling of the seals. The Scottish Marine Animals Strandings Scheme receives financial support from the Scottish Government Marine Directorate and the UK Department of Environment, Farming and Rural Affairs (Defra). The Norwegian Research Council and the Norwegian Polar Institute financed Norwegian collections.

LITERATURE CITED

- Calle PP, Seagars DJ, McClave C, Senne D, House C, House JA (2008) Viral and bacterial serology of six free-ranging bearded seals *Erignathus barbatus*. Dis Aquat Org 81: 77–80
- Carmichael LE, Kenney RM (1970) Canine brucellosis: the clinical disease, pathogenesis, and immune response. J Am Vet Med Assoc 156:1726–1734
- Cvetnić Ž, Duvnjak S, Đuras M, Gomerčić T, Reila I, Zdelar-Tuk M, Špičić S (2016) Evidence of *Brucella* strain ST27 in bottlenose dolphin (*Tursiops truncatus*) in Europe. Vet Microbiol 196:93–97
 - Dierauf LA (1994) Pinniped forensic necropsy and tissue collection guide. US Dept Commerce, NOAA Tech Memo NMFS-OPR-94-3
 - Dierauf L, Gulland FMD (eds) (2001) CRC handbook of marine mammal medicine. CRC Press, Boca Raton, FL
- Ewalt DR, Payeur JB, Martin BM, Cummins DR, Miller GM (1994) Characteristics of a *Brucella* species from a bottlenose dolphin (*Tursiops truncatus*). J Vet Diagn Invest 6: 448–452
- Farrell ID (1974) The development of a new selective medium for the isolation of *Brucella abortus* from contaminated sources. Res Vet Sci 16:280–286
- Forbes LB, Nielsen O, Measures L, Ewalt DR (2000) Brucellosis in ringed seals and harp seals from Canada. J Wildl Dis 36:595–598
- Foster G, Jahans KL, Reid RJ, Ross HM (1996) Isolation of Brucella species from cetaceans, seals, and an otter. Vet Rec 138:583–586
- Foster G, MacMillan AP, Godfroid J, Howie F and others (2002) A review of *Brucella* sp. infection of sea mammals with particular emphasis on isolates from Scotland. Vet Microbiol 90:563–580
- Foster G, Osterman B, Godfroid JG, Jacques I, Cloeckaert A (2007) Brucella ceti sp. nov. and Brucella pinnipedialis sp. nov. for Brucella strains with cetaceans and seals as their preferred hosts. Int J Syst Evol Microbiol 57: 2688–2693
- Garner MM, Lambourn DM, Jeffries SJ, Hall PB and others (1997) Evidence of *Brucella* infection in *Parafilaroides* lungworms in a Pacific harbor seal (*Phoca vitulina richardsii*). J Vet Diagn Invest 9:298–303
- Garnier R, Gandon S, Harding KC, Boulinier T (2014) Length of intervals between epidemics: evaluating the influence of maternal transfer of immunity. Ecol Evol 4: 568–575
- Goldstein T, Zabka TS, Delong RL, Wheeler EA and others (2009) The role of domoic acid in abortion and premature parturition of Californian sea lions (*Zalophus californianus*) on San Miguel Island. J Wildl Dis 45: 91–108
- Groussaud P, Shankster SJ, Koylass MS, Whatmore AM (2007) Molecular typing divides marine mammal strains of *Brucella* into at least three groups with distinct host

preferences. J Med Microbiol 56:1512-1518

- Guzman-Verri C, Gonzales-Barrientos R, Hernandez-Mora G, Morales JA, Baquero-Calvo J, Chaves-Olarte E, Moreno E (2012) *Brucella ceti* and brucellosis in cetaceans. Front Cell Infect Microbiol 2:3
- JNCC/Defra (Joint Nature Conservation Commission/UK Department of Environment, Farming and Rural Affairs) (2013) European Community Directive on the conservation of natural habitats and of wild flora and fauna (92/42/EEC), third report by the United Kingdom under article 17. https://onedrive.live.com/?authkey=%21AI9z EzfDLarYkjU&cid=2EBEF263871972EE&id=2EBEF2638 71972EE%21153&parId=2EBEF263871972EE%21149&o =OneUp (accessed 11 November 2016)
 - Kovacs KM (2002) Hooded seal (*Cystophora cristata*). In: Perrin WF, Würsig B, Thewissen JGM (eds) Encyclopedia of marine mammals. Academic Press, San Diego, CA, p 580–582
- Kovacs KM (2016) Erignathus barbatus. In: The IUCN Red List of Threatened Species 2016:e.T8010A45225428. http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T8010A 45225428.en (accessed 5 August 2016)
- Lambourn DM, Garner M, Ewalt D, Raverty S and others (2013) Brucella pinnipedialis infections in Pacific harbor seals (Phoca vitulina richardsi) from Washington state, USA. J Wildl Dis 49:802–815
- Larsen AK, Nymo IH, Boysen P, Tryland M, Godfroid J (2013a) Entry and elimination of marine mammal Brucella spp. by hooded seal (Cystophora cristata) alveolar macrophages in vitro. PLOS ONE 8:e70186
- Larsen AK, Nymo IH, Briquemont B, Sørensen K, Godfroid J (2013b) Entrance and survival of *Brucella pinnipedialis* by hooded seal strain in human macrophages and epithelial cells. PLOS ONE 8:e84861
- Larsen AK, Godfroid J, Nymo IH (2016) Brucella pinnipedialis in hooded seal (Cystophora cristata) primary epithelial cells. Acta Vet Scand 58:9
 - Lavigne DM (2002) Harp seal (*Pagophilus groenlandicus*). In: Perrin WF, Würsig B, Thewissen JGM (eds) Encyclopedia of marine mammals. Academic Press, San Diego, CA, p 560–562
 - Miyazaki N (2002) Ringed, Caspian, and Baikal seals *Pusa hispida, P. caspica*, and *P. sibirica*. In: Perrin WF, Würsig B, Thewissen JGM (eds) Encyclopedia of marine mammals. Academic Press, San Diego, CA, p 1033–1037
- Nymo IH, Tryland M, Godfroid J (2011) A review of *Brucella* infection in marine mammals with special emphasis on *Brucella pinnipedialis* in the hooded seal (*Cystophara cristata*). Vet Res 42:93
- Nymo IH, Godfroid JG, Åsbakk K, Larsen AK, das Neves CG, Rødven R, Tryland M (2013a) A protein A/G indirect enzyme-linked immunosorbent assay for the detection of anti-*Brucella* antibodies in Arctic wildlife. J Vet Diagn Invest 25:369–375
- Nymo IH, Tryland M, Frie AK, Haug T, Foster G, Rødven R, Godfroid J (2013b) Age-dependent prevalence of anti-Brucella antibodies in hooded seals Cystophora cristata. Dis Aquat Org 106:187–196
 - Nymo IH, Arias MA, Godfroid J, Pardo J, Alvarez MP, Alcaraz A, Jiménez de Bagüés MP (2016) Marine mammal *Brucella* reference strains are attenuated in a BALB/c mouse model. PLOS ONE 11:e0150432
- Chishi K, Bando T, Abe E, Kawai Y, Fulise Y, Maruyama T (2016) Long-term and large-scale epidemiology of *Bru*cella infection in baleen whales and sperm whales in the

western North Pacific and Antarctic oceans. J Vet Med Sci $78{:}1457{-}1464$

- Pomeroy PP, Hammond JA, Hall AJ, Lonergan M, Duck CD, Smith VJ, Thompson H (2005) Morbillivirus neutralising antibodies in Scottish grey seals *Halichoerus grypus*: assessing the effects of the 1988 and 2002 PDV epizootics. Mar Ecol Prog Ser 287:241–250
- *Ross PS, Pohajdak B, Bowen WD, Addison RF (1993) Immune function in free-ranging harbour seal (*Phoca vit-ulina*) mothers and their pups during lactation. J Wildl Dis 29:21–29
- Ross HM, Foster G, Reid RJ, Jahans KL, MacMillan AP (1994) Brucella species infection in marine mammals. Vet Rec 134:359
 - Stewart REA, Stewart BE (2009) Female reproductive systems. In: Perrin WF, Würsig B, Thewissen JGM (eds) Encyclopedia of marine mammals. Academic Press, San Diego, CA, p 423–429
- Stokke G (2010) Our seals in Polaria, an Arctic adventure. www.polaria.no/our-seals.155311.en.html
 - Sutherland SS, Searson J (1990) The immune response to *Brucella abortus*: the humoral response. In: Nielsen K, Duncan JR (eds) Animal brucellosis, 1st edn. CRC Press, Boca Raton, FL, p 65–81

Tizard I (2000) Immunity in the fetus and newborn animal.

Editorial responsibility: Stephen Raverty, Abbotsford, British Columbia, Canada

In: Tizard I (ed) Veterinary immunology: an introduction. Saunders, Philadelphia, PA, p 210–221

- Tryland M, Kleivane L, Alfredsson A, Kjeld M, Arnason A, Stuen S, Godfroid J (1999) Evidence of *Brucella* infection in marine mammals in the North Atlantic Ocean. Vet Rec 144:588–592
- van Bree PJH (2000) A review of recent extralimital records of the bearded seal (*Erignathus barbatus*) on the west European continental coast. Mar Mamm Sci 16:261–263
- Whatmore AM, Perrett LL, MacMillan AP (2007) Characterisation of the genetic diversity of *Brucella* by multilocus sequencing. BMC Microbiol 7:34
- Whatmore AM, Dawson CE, Groussaud P, Koylass MS and others (2008) Marine mammal *Brucella* genotype associated with zoonotic infection. Emerg Infect Dis 14: 517–518
 - Whatmore AM, Koylass MS, Muchowski J, Edwards-Smallbone J, Gopaul KK, Perrett LL (2016) Extended multilocus sequence analysis to describe the global population structure of the genus *Brucella*: phylogeography and relationship to biovars. Front Microbiol 7:2049
- Whatmore AM, Dawson C, Muchowski J, Perrett LL and others (2017) Characterisation of North American *Brucella* isolates from marine mammals. PLOS ONE 12: e0184758

Submitted: August 28, 2017; Accepted: January 11, 2018 Proofs received from author(s): March 7, 2018