

The following appendices accompany the article

Prey landscapes help identify potential foraging habitats for leatherback turtles in the NE Atlantic

Matthew J. Witt¹, Annette C. Broderick¹, David J. Johns², Corinne Martin³,
Rod Penrose⁴, Marinus S. Hoogmoed⁵, Brendan J. Godley^{1,*}

¹Marine Turtle Research Group, Centre for Ecology and Conservation, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

²Sir Alister Hardy Foundation for Ocean Science, The Laboratory, Citadel Hill, Plymouth PL1 2PB, UK

³Department of Geographical and Life Sciences, Canterbury Christ Church University, Canterbury CT1 1QU, UK

⁴Marine Environmental Monitoring, Penwalk, Llechryd, Cardigan, Ceredigion SA43 2PS, UK


⁵Museu Paraense Emilio Goeldi/CZO/Herpetologia, Caixa Postal 399, CEP 66017-970 Belém, PA, Brasil

*Corresponding author. Email: b.j.godley@exeter.ac.uk


Marine Ecology Progress Series 337:231–244 (2007)

Appendix 1. Monthly distribution of CPR sampling locations for the NE and central North Atlantic


January, n = 12 098




April, n = 14 823




February, n = 12 725




May, n = 14 868



March, n = 15 343




June, n = 14 592




Appendix 1 (continued)


July, n = 15 098




October, n = 14 576




August, n = 15 297




November, n = 13 865



September n = 14 851



December, n = 12 750



Appendix 2. Seasonal mean long-term distribution of siphonophores recorded by the CPR Survey mapped using (a–d) abundance and (e–i) proportion of positive records. Note similarity in results for the 2 methods

