

# Diet of invasive lionfish on hard bottom reefs of the Southeast USA: insights from stomach contents and stable isotopes

Roldan C. Muñoz<sup>1,\*</sup>, Carolyn A. Currin<sup>2</sup>, Paula E. Whitfield<sup>2</sup>

<sup>1</sup>National Marine Fisheries Service, Beaufort Laboratory, National Oceanic and Atmospheric Administration, Beaufort, North Carolina 28516, USA

<sup>2</sup>National Ocean Service, Center for Coastal Fisheries and Habitat Research, National Oceanic and Atmospheric Administration, Beaufort, North Carolina 28516, USA

\*Email: roldan.munoz@noaa.gov

*Marine Ecology Progress Series 432:181–193 (2011)*

**Supplements 1–5.** Distribution of sample sizes, additional methodological details and analyses, and selection of U.S. South Atlantic fishes with potential dietary overlap with *Pterois volitans*.

**Supplement 1.** Table S1. Distribution of sample sizes of *Pterois volitans* by study site and by year, study site characteristics, and distribution of prey surveys. Stomach samples do not include empty stomachs

| Site              | 2004 Stomachs | 2006 Stomachs | 2004 Isotopes | Depth (m) | Habitat <sup>a</sup> | Prey Survey <sup>b</sup> |
|-------------------|---------------|---------------|---------------|-----------|----------------------|--------------------------|
| 18 Fathom         | 31            | 3             | 24            | 38.1      | wreck                | X                        |
| Bobby6            | 2             | 0             | 2             | 43.9      | low                  |                          |
| Capdan99          | 2             | 0             | 2             | 39.6      | medium               |                          |
| CF001             | 7             | 2             | 10            | 41.2      | high                 | X                        |
| CF14              | 6             | 0             | 6             | 39.6      | medium               |                          |
| City of Houston   | 3             | 0             | 4             | 30.5      | wreck                | X                        |
| Dan's spot        | 2             | 8             | 2             | 39.6      | high                 | X                        |
| Gerry's wreck     | 11            | 5             | 12            | 45.4      | low                  | X                        |
| Kenny1            | 2             | 10            | 1             | 42.7      | medium               | X                        |
| Kenny3            | 0             | 0             | 1             | 42.7      | medium               |                          |
| Lobsterrocksth3   | 9             | 23            | 12            | 42.7      | high                 | X                        |
| Lobsterrocknorth1 | 5             | 7             | 4             | 36.6      | low                  | X                        |
| Lobsterrocksth1   | 5             | 0             | 10            | 41.2      | high                 |                          |
| Lobsterwreck      | 3             | 0             | 4             | 38.1      | wreck                |                          |
| Naecobow          | 6             | 0             | 7             | 42.7      | wreck                |                          |
| Naecostern        | 3             | 19            | 6             | 42.7      | wreck                | X                        |
| SETower           | 7             | 0             | 8             | 41.2      | medium               | X                        |
| Woo2              | 0             | 2             | 0             | 39        | high                 | X                        |

|       |     |    |     |
|-------|-----|----|-----|
| Total | 104 | 79 | 115 |
|-------|-----|----|-----|

<sup>a</sup>Wreck sites were characterized by vertical relief >1 m heavily colonized by sessile invertebrates and macroalgae, with the wreck structure providing ample overhangs and undercuts for shelter purposes. Low relief hard bottom sites were characterized by vertical relief <0.25 m sparsely colonized by sessile invertebrates and macroalgae, with no overhangs. Available shelter consisted of holes eroded/scoured into the mostly flat surfaces. Medium relief hard bottom sites consisted of vertical relief >0.25 m but < 1 m. Sessile invertebrates and macroalgae tended to occupy greater percent cover than on low relief sites and shelter consisted of eroded/scoured holes. High relief hard bottom sites provided vertical relief >1 m in the form of ledges or rocky outcrops, were densely colonized by sessile invertebrates and macroalgae, and contained overhangs or undercuts as shelter. Riggs et al. (1998) provides further details of benthic community processes on hard bottoms and Kendall et al. (2009) provides justification for ledge height classifications.

<sup>b</sup>X indicates a site was surveyed with a visual prey census.

**Supplement 2.** *Pterois volitans* — cumulative prey analysis. We constructed cumulative prey curves for each year separately and combined to determine the adequacy of sample sizes of *Pterois volitans* and plotted the average number of new prey items found in each stomach versus the total number of stomachs analyzed (Ferry & Caillet 1996). We used PRIMER to permute the order of stomachs analyzed 999 times and considered that an adequate number of stomachs had been analyzed if an asymptote was reached (Ver. 6; Warwick 1993, Warwick & Clarke 2001). We also calculated five different estimates of prey items (Chao 1, Chao 2, jackknife 1, jackknife 2, bootstrap) to determine how many prey types were potentially missed by sampling. These additional estimators differ in exact method of computation but tend to utilize rare prey items to estimate the true number of prey in the diet (Colwell & Coddington 1994). To provide a standard measure of sample size precision and for comparison among diet studies, we calculated the mean coefficient of variation ( $CV = [\text{standard deviation} / \text{mean}] \times 100$ ) of the mean cumulative number of prey taxa generated for the final four stomach samples (Bizzarro et al. 2007). Values of < 10% CV are a general benchmark for adequate precision in age and growth studies (Campana 2001, Bizzarro et al. 2007). For both years combined, the cumulative number of prey categories (18) recorded from these specimens had neared an asymptote and additional estimates of prey items indicated that only 1.5–3 prey categories may have been missed by sampling. In addition, the mean CV of the mean cumulative number of prey taxa generated for the final four stomach samples was 0.59%, suggesting that characterizing the diet of lionfish with these samples was sufficiently precise. For 2004, 14 prey categories were recorded, only 0.4–2 prey categories may have been missed by sampling, and the mean CV for the final four stomach samples was 1.1%. For 2006, 15 prey categories were recorded, only 2–4 prey categories may have been missed by sampling, and the mean CV for the final four stomach samples was 1.5%. Figure S1 shows the mean cumulative number of prey categories per stomach sample for *Pterois volitans*. Prey categories are from Table 1 in the main article with the addition of unidentified stomach contents. Mullidae and Echinodermata (items regurgitated from multiple fish while in a common holding tank) were not included because they could not be assigned to a single or multiple stomachs.




Fig. S1. Mean cumulative number of prey categories per stomach sample for *Pterois volitans*

**Supplement 3.** Methodological details of 1-way Analysis of Similarities (ANOSIM) for *Pterois volitans* stomach contents.

For ANOSIM of prey categories, each prey category was treated as a variable (species) and the percent weight of each category per stomach examined between years. For prey lengths, we generated a frequency distribution of all prey lengths across years and prey length bins were then treated as variables and the number of prey in each length bin examined between years. For prey volume, we generated a frequency distribution of total prey volume per stomach across years and total prey volume per stomach bins were then treated as variables and the distribution of total stomach volume examined between years. For prey number, we generated a frequency distribution of total number of prey per stomach (without regards to prey category) across years and total number of prey per stomach bins were then treated as variables and the distribution of total number of prey per stomach examined between years. Note that the ANOSIM of prey categories examines differences between years by accounting for the identity and number of prey consumed per category, while the ANOSIM of prey number differs by examining differences between years in the physical prey processing characteristics of lionfish (i.e., the total number of prey per stomach without regards to prey category).

**Supplement 4.** Two-way ANOSIM pairwise tests (Table S2) and similarity percentages (SIMPER) analyses (Table S3) comparing changes in *Pterois volitans* diet with size while accounting for any differences between years.

Table S2. Two-way ANOSIM pairwise tests

| Size class (TL, cm)    | R statistic | p     |
|------------------------|-------------|-------|
| 15 – 19.9 vs 20 – 24.9 | 0.133       | 0.045 |
| 15 – 19.9 vs 25 – 29.9 | 0.212       | 0.094 |
| 15 – 19.9 vs 30 – 34.9 | 0.345       | 0.001 |
| 15 – 19.9 vs ≥35       | 0.223       | 0.028 |
| 20 – 24.9 vs 25 – 29.9 | 0.094       | 0.195 |
| 20 – 24.9 vs 30 – 34.9 | 0.120       | 0.099 |
| 20 – 24.9 vs ≥35       | 0.072       | 0.192 |
| 25 – 29.9 vs 30 – 34.9 | -0.0290     | 0.766 |
| 25 – 29.9 vs ≥35       | -0.0250     | 0.655 |
| 30 – 34.9 vs ≥35       | 0.022       | 0.326 |

Table S3. Two-way similarity percentages (SIMPER) analyses comparing changes in *Pterois volitans* diet with size while accounting for any differences between years. SIMPER results between years were similar to one-way results already reported so those data are not presented. Results presented following two-way ANOSIM significant pairwise tests for size classes (S3.1) 15 – 19.9 cm TL vs 20 – 24.9 cm TL, (S3.2) 15 – 19.9 vs 30 – 34.9, and (S3.3) 15 – 19.9 vs ≥35. Abundance<sup>a</sup> is average percent volume of recognizable prey in each size class. Percent contribution is the contribution of each prey category to the observed dissimilarity. Prey categories contributing < 3% to the observed dissimilarity are not listed

Table S3.1

| Prey Category | Size Class Abundance <sup>a</sup> |                | Percent Contribution |
|---------------|-----------------------------------|----------------|----------------------|
|               | 15 – 19.9 cm TL                   | 20 – 24.9cm TL |                      |
| Crustacea     | 5.73                              | 0.83           | 28.28                |
| Haemulidae    | 0.91                              | 2.29           | 17.42                |
| Pomacentridae | 0.91                              | 1.42           | 12.92                |
| Carangidae    | 0                                 | 1.63           | 11.29                |
| Blennidae     | 0.91                              | 1.80           | 10.03                |
| Acanthuridae  | 0                                 | 0.83           | 6.24                 |
| Serranidae    | 0.87                              | 0.75           | 5.58                 |

<sup>a</sup>Values presented are square root transformed as in SIMPER analysis.

Table S3.2

| Prey Category | Size Class Abundance <sup>a</sup> |                | Percent Contribution |
|---------------|-----------------------------------|----------------|----------------------|
|               | 15 – 19.9 cm TL                   | 30 – 34.9cm TL |                      |
| Crustacea     | 5.73                              | 0.32           | 29.92                |
| Haemulidae    | 0.91                              | 2.35           | 20.2                 |
| Serranidae    | 0.87                              | 3.14           | 10.0                 |
| Carangidae    | 0                                 | 1.26           | 6.75                 |
| Gobiidae      | 0.91                              | 0.3            | 6.51                 |
| Pomacentridae | 0.91                              | 0.12           | 5.06                 |
| Blennidae     | 0.91                              | 0.38           | 5.05                 |
| Scaridae      | 0                                 | 1.65           | 3.2                  |
| Labridae      | 0.02                              | 0.53           | 3.02                 |
| Bothidae      | 0                                 | 0.34           | 3.01                 |

<sup>a</sup>Values presented are square root transformed as in SIMPER analysis.

Table S3.3

| Prey Category | Size Class Abundance <sup>a</sup> |           | Percent Contribution |
|---------------|-----------------------------------|-----------|----------------------|
|               | 15 – 19.9 cm TL                   | ≥35 cm TL |                      |
| Crustacea     | 5.73                              | 0.88      | 31.31                |
| Haemulidae    | 0.91                              | 1.62      | 17.9                 |
| Serranidae    | 0.87                              | 3.2       | 11.18                |
| Scaridae      | 0                                 | 1.75      | 9.45                 |
| Bothidae      | 0                                 | 0.96      | 7.67                 |
| Pomacentridae | 0.91                              | 0.59      | 5.34                 |
| Carangidae    | 0                                 | 0.39      | 3.77                 |
| Gobiidae      | 0.91                              | 0         | 3.77                 |

<sup>a</sup>Values presented are square root transformed as in SIMPER analysis.

**Supplement 5.** Table S4. Selection of U.S. South Atlantic fishes with potential dietary overlap with *Pterois volitans*.<sup>a</sup>

| Species   | Fishes as % of diet | Sample size | Prey taxa overlap <sup>b</sup> |
|---|---------------------|-------------|--------------------------------|
| sand diver, <i>Synodus intermedius</i>                | 94.5                | 18          | Y                              |
| inshore lizardfish, <i>S. foetens</i> <sup>d</sup>    | 76 – 100            | 3 – 603     | Y                              |
| red lizardfish, <i>S. synodus</i>                     | 100                 | 2           | ?                              |
| green moray, <i>Gymnothorax funebris</i> <sup>e</sup> | ? <sup>c</sup>      | 3           | ?                              |
| spotted moray, <i>G. moringa</i>                      | 100                 | 6           | Y                              |
| purplemouth moray, <i>G. vicinus</i>                  | 62.5                | 4           | Y                              |
| spotted snake eel, <i>Ophichthus ophis</i>            | 50                  | 4           | Y                              |
| coronetfish, <i>Fistularia tabacaria</i>              | 100                 | 2           | ?                              |
| trumpetfish, <i>Aulostomus maculatus</i>              | 73.5                | 69          | Y                              |
| scamp, <i>Mycteroperca phenax</i> <sup>f</sup>        | “most important”    | 91          | Y                              |
| gag, <i>M. microlepis</i> <sup>g</sup>                | 95.1 – 99.8         | 979         | Y                              |
| black grouper, <i>M. bonaci</i> <sup>h</sup>          | 76.7 – 100          | 4 – 72      | Y                              |
| yellowmouth grouper, <i>M. interstitialis</i>         | 100                 | 5           | Y                              |
| tiger grouper, <i>M. tigris</i>                       | 100                 | 34          | Y                              |
| yellowfin grouper, <i>M. venenosa</i>                 | 95.3                | 51          | Y                              |
| graysby, <i>C. cruentata</i>                          | 66.2                | 26          | Y                              |
| Nassau grouper, <i>Epinephelus striatus</i>           | 54                  | 153         | Y                              |
| schoolmaster, <i>Lutjanus apodus</i>                  | 60.7                | 58          | Y                              |
| club snapper, <i>L. cyanopterus</i>                   | 100                 | 11          | Y                              |
| dog snapper, <i>L. jocu</i>                           | 60.7                | 56          | Y                              |
| mahogany snapper, <i>L. mahogoni</i>                  | 75                  | 8           | Y                              |
| peacock flounder, <i>Bothus lunatus</i>               | 85.7                | 7           | Y                              |
| longlure frogfish, <i>Antennarius multiocellatus</i>  | 75                  | 5           | Y                              |
| splitlure frogfish, <i>A. scaber</i>                  | 100                 | 5           | Y                              |

<sup>a</sup>Taxa are listed if they were determined to have at least 50% fish prey in stomach contents and if they are believed to have a benthic feeding strategy while occupying the mesocarnivore trophic level.

<sup>b</sup>Prey taxa overlap with Y indicates that species listed had consumed the same species or genus of prey as lionfish; ? = unknown.

<sup>c</sup>Green moray were included because they are the largest western Atlantic moray and because Gudger (1929) (cited in Randall 1967) reported fish in the stomachs of three specimens, although their % contribution to the diet is not known.

References: Dietary data from Randall (1967) except as indicated; <sup>d</sup>Randall (1967) and Cruz-Escalona et al. (2005);

<sup>e</sup>Randall (1967) and Gudger (1929); <sup>f</sup>Matheson et al. (1986); <sup>g</sup>Naughton & Saloman (1985); <sup>h</sup>Randall (1967) and Brulé et al. (2005).

## LITERATURE CITED

- Bizzarro JJ, Robinson HJ, Rinewalt CS, Ebert DA (2007) Comparative feeding ecology of four sympatric skate species off central California, USA. *Environ Biol Fishes* 80:197–220
- Brulé T, Puerto-Novelo E, Pérez-Díaz E, Renán-Galindo X (2005) Diet composition of juvenile black grouper (*Mycteroperca bonaci*) from coastal nursery areas of the Yucatan Peninsula, Mexico. *Bull Mar Sci* 77:441–452
- Campana SE (2001) Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *J Fish Biol* 59:197–242
- Colwell RK, Coddington JA (1994) Estimating terrestrial biodiversity through extrapolation. *Philos Trans R Soc B* 345:101–118
- Cruz-Escalona VH, Peterson MS, Campos-Davila L, Zetina-Rejon M (2005) Feeding habits and trophic morphology of inshore lizardfish (*Synodus foetens*) on the central continental shelf off Veracruz, Gulf of Mexico. *J Appl Ichthyol* 21:525–530
- Ferry LA, Caillet GM (1996) Sample size and data analysis: Are we characterizing and comparing diet properly? In: MacKinley D, Shearer K (eds) *Gutshop '96. Feeding ecology and nutrition in fish: symposium proceedings*. American Fisheries Society, San Francisco, p 71–80
- Gudger EW (1929) On the morphology, coloration and behavior of seventy teleostean fishes of Tortugas, Florida. *Papers of the Tortugas Lab. Carnegie Inst Wash Publ* 391:149–204
- Kendall MS, Bauer LJ, Jeffrey CFG (2009) Influence of hard bottom morphology on fish assemblages of the continental shelf off georgia, southeastern USA. *Bull Mar Sci* 84:265–286
- Matheson RH, Huntsman GR, Manooch CS (1986) Age, growth, mortality, food and reproduction of the scamp, *Mycteroperca phenax*, collected off North Carolina and South Carolina. *Bull Mar Sci* 38:300–312
- Naughton SP, Saloman CH (1985) Food of gag (*Mycteroperca microlepis*) from North Carolina and three areas of Florida. US Dept Commerce, NOAA Tech Memo NMFS-SEFC-160
- Randall JE (1967) Food habits of reef fishes of the West Indies. *Stud Trop Oceanogr* 5:665–847
- Riggs SR, Ambrose WG, Cook JW, Snyder SW, Snyder SW (1998) Sediment production on sediment-starved continental margins; the interrelationship between hardbottoms, sedimentological and benthic community processes, and storm dynamics. *J Sediment Res* 68:155–168
- Warwick RM (1993) Environmental impact studies on marine communities: pragmatical considerations. *Aust J Ecol* 18:63–80
- Warwick RM, Clarke KR (2001) Practical measures of marine biodiversity based on relatedness of species. *Oceanogr Mar Biol Annu Rev* 39:207–231