

Effects of temperature on life history set the sensitivity to fishing in Atlantic cod *Gadus morhua*

Hui-Yu Wang¹, Louis W. Botsford^{2,*}, J. Wilson White³, Michael J. Fogarty⁴, Francis Juanes^{5,6}, Alan Hastings⁷, Matthew D. Holland², Keith Brander⁸

¹Institute of Oceanography, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd, Taipei 106, Taiwan

²Department of Wildlife, Fish and Conservation Biology, University of California, One Shields Avenue, Davis, CA 95616, USA

³Department of Biology and Marine Biology, University of North Carolina Wilmington, 601 S. College Rd., Wilmington, NC 28403, USA

⁴NOAA Fisheries Service, Northeast Fisheries Science Center, 166 Water St, Woods Hole, MA 02543, USA

⁵Department of Biology, University of Victoria, Victoria, BC V8P 5C2, Canada

⁶Department of Environmental Conservation, University of Massachusetts, 160 Holdsworth Way, Amherst, MA 01003, USA

⁷Department of Environmental Science and Policy, University of California, One Shields Avenue, Davis, CA 95616, USA

⁸Technical University of Denmark, Charlottenlund Slot, Jægersborg Allé 1, 2920 Charlottenlund, Denmark

*Corresponding author: lwbotsford@ucdavis.edu

Marine Ecology Progress Series 514: 217–229 (2014)

Supplement. This supplement shows the figures displaying the modeled distributions of spawning biomass (Fig. S1) and deviation of A_{50S} from S_{50S} (Fig. S2) for the 20 Atlantic cod populations




Fig. S1. Normalized spawning biomass from the model, at 3 different levels of fishing mortality (F) rates for 14 cod *Gadus morhua* populations. These are calculated from the parameter values in Tables 2 & 3 in the main article




Fig. S2. Scatterplot of A_{50} vs. S_{50} for the 20 Atlantic cod *Gadus morhua* populations. The straight line is a 1:1 line. The population abbreviations are shown in Table 1 in the main article