

Erratum

Alphaproteobacterial dominance in a large mesotrophic lake (Lake Biwa, Japan)

Yoko Nishimura, Toshi Nagata*

Aquatic Microbial Ecology 48: 231–240, 2007

- On page 235, in the last line of Table 1, the 2 values were displaced one column to the left. The corrected table is shown here.

Table 1. Summary of community structures of total and filamentous forms of prokaryotes determined by FISH in Lake Biwa. Negative control was not subtracted from the other results. CF cluster: *Cytophaga-Flavobacteria* cluster; CV: coefficient of variation

Phylogenetic group	Probe	— % total prokaryotic abundance —					
		Mean	SD	CV	Min.	Max.	n
Total							
<i>Bacteria</i>	Eub338	59.1	8.8	14.9	34.5	73.0	40
<i>Alphaproteobacteria</i>	Alfa968	41.7	12.4	29.7	20.1	66.9	40
<i>Betaproteobacteria</i>	Beta42	11.8	6.4	54.2	5.4	36.7	40
<i>Gammaproteobacteria</i>	Gam42a	2.2	1.6	72.7	0.0	6.8	40
CF cluster	CF319	5.8	3.5	60.3	1.6	18.1	40
<i>Actinobacteria</i>	HGC69a	14.3	9.6	67.1	2.7	38.4	40
<i>Archaea</i>	Arch915	8.3	7.8	94.0	1.6	36.4	40
Negative control	NON338	2.8	1.3	46.4	0.5	5.9	39
Sum of the major groups ^a		73.6	21.8				
Filament							
<i>Bacteria</i>	Eub338	87.6	6.2	7.1	69.7	94.5	44
<i>Alphaproteobacteria</i>	Alfa968	64.1	14.8	23.1	13.6	86.3	44
CF cluster	CF319	28.3	16.0	56.5	1.5	65.0	44
LD2 clade	LD2	17.0	14.1	82.9	0.6	53.6	44
Sum of <i>Alphaproteobacteria</i> + CF cluster		92.4	15.0				

^a*Alphaproteobacteria*, *Betaproteobacteria*, CF cluster and *Actinobacteria*

*Corresponding author. Email: nagata@ecology.kyoto-u.ac.jp