

NOTE

First record of chytridiomycosis in Bolivia (*Rhinella quechua*; Anura: Bufonidae)

J. Sebastián Barrionuevo^{1,*}, Rodrigo Aguayo², Esteban O. Lavilla¹

¹Instituto de Herpetología, Fundación Miguel Lillo, Miguel Lillo 251 (4000) Tucumán, Argentina

²Centro de Biodiversidad y Genética, Universidad Mayor de San Simón, Cochabamba, Bolivia

ABSTRACT: The finding of tadpoles of *Rhinella quechua* (Huayramayu River, Carrasco National Park, Cochabamba, Bolivia) with oral abnormalities caused by *Batrachochytrium dendrobatidis* constitutes the first record of this fungal infection reported for Bolivian amphibians.

KEY WORDS: Chytridiomycosis · *Rhinella quechua* · Bolivia

—Resale or republication not permitted without written consent of the publisher—

INTRODUCTION

Chytridiomycosis is an amphibian emerging disease caused by the fungal pathogen *Batrachochytrium dendrobatidis*. This mortal disease has been linked to the decline of amphibian species worldwide, especially in Australia and Latin America (Berger et al. 1998, Lips et al. 2006), although there is debate about its epidemiology (Alford et al. 2007, Di Rosa et al. 2007, Pounds et al. 2007). Two hypotheses on emerging diseases have been proposed. The novel pathogen hypothesis states that the pathogen has been dispersed recently, affecting naive populations, while the endemic pathogen hypothesis claims that cofactors raise the virulence of the pathogen already present in the environment (Rachowicz et al. 2005, Skerratt et al. 2007).

Bolivia is ranked among the 10 most diverse countries for amphibians in the world, containing around 258 described species (Reichle 2003, R. Aguayo unpubl. data). Even though no large numbers of amphibian deaths have been reported in the country (Reichle 2006), some amphibian populations have experienced declines (De la Riva 2005). The causes of decline are not clear, leaving chytridiomycosis as a possible cause. Although to date, the disease has not been diagnosed in Bolivia, it has been found infecting amphibians in southern Peru (Seimon et al. 2005, 2006) and north-western Argentina (Barrionuevo & Mangione 2006).

Here we report the presence of chytridiomycosis in torrent-dwelling tadpoles of *Rhinella quechua*, whose population status is listed as vulnerable (IUCN 2006, Reichle 2006).

MATERIALS AND METHODS

Twelve tadpoles of *Rhinella quechua* were collected on 10 June 2007 from Huayramayu River, Sehuencas Forest, Carrasco National Park (Cochabamba Department, Bolivia, 17° 56' S, 65° 29' W) at 2530 m a.s.l. Tadpoles were identified by rearing some of them to metamorphosis. Specimens were anesthetized and fixed in 10% neutral buffered formaldehyde, and intact voucher specimens are housed in the collection of the Centro de Biodiversidad y Genética, Cochabamba, Bolivia (CBG 1502). Keratinized mouthparts of 3 tadpoles were excised with a clean, sharp scalpel. Standard histological techniques for light microscopy were applied. The samples were sectioned at 6 µm, stained with hematoxylin and eosin, and examined at 63 and 100× magnification. We analyzed 48 histological sections of tadpole CBG 1502-1, 18 sections of CBG 1502-2, and 18 sections of CBG 1502-3. To identify chytrids, we followed Berger et al. (1999) and Pessier et al. (1999).

RESULTS

Of the *Rhinella quechua* tadpoles, 9 had normal mouthparts (Fig. 1a) and 3 tadpoles showed anomalies in the keratinized mouthparts of the oral disc (Fig. 1b). The anomalies included lost, interrupted, or misshapen tooththrows, and lack of pigmentation in the tooththrows and upper and lower jaw sheaths (Fig. 1b). One of these tadpoles (CBG 1502-1) was positive for *Batrachochytrium dendrobatidis* under histological exami-

*Email: pretorum@hotmail.com

nation. Zoosporangia at different stages were evident in the stratum corneum of keratinized oral disc skin. Mature zoosporangia are usually round, ranging from 7.5 to 10 μm in diameter, with a thin wall (Fig. 2), and contain rounded basophilic zoospores approximately 1 μm in diameter. In some zoosporangia, a discharge tube formed by an evagination of the zoosporangium wall was evident. The empty post-discharge zoosporangia were the most frequently observed stages in the stratum corneum. The zoosporangial morphology matches the descriptions provided in previous works (Berger et al. 1999, Pessier et al. 1999). Hyperkeratosis was not observed in any of the examined specimens. In the other 2 tadpoles (CBG 1502-2 and 1502-3), we failed to identify the chytrid fungus, probably due to the clustered distribution of zoosporangia.

DISCUSSION AND CONCLUSIONS

This is the first report of chytridiomycosis in Bolivia. Although some amphibian population declines have been reported in this country (De la Riva 2005) and the presence of chytrid fungus was highly suspected (De la Riva et al. 2005) and predicted (Ron 2005), until now, the disease had not been diagnosed in any amphibian in the country.




Fig. 1. *Rhinella quechua*. (a) Normal and (b) abnormal keratinized mouthparts in stage 35 tadpoles (Gosner 1960). (Scale bar = 1 mm)




Fig. 2. *Rhinella quechua*. (a) Skin section of the oral disc of a tadpole. Note mature zoosporangium in the stratum corneum (arrow). Zoospores can be distinguished within the zoosporangia. The rest of zoosporangia are empty (scale bar = 10 μm). (b) An empty zoosporangium with projecting discharge tube (arrow; scale bar = 10 μm)

The oral disc anomalies found in tadpoles of *Rhinella quechua* are coincident with those described as indicative of chytridiomycosis (Fellers et al. 2001, Knapp & Morgan 2006). Although oral disc anomalies caused by chytridiomycosis differ from those caused by DDT or corticosterone contamination (Fellers et al. 2001) factors other than chytridiomycosis can affect mouthpart pigmentation (e.g. temperature; Rachowicz 2002) or alter tooththrow formation (e.g. development, diet; Altig 2007). For these reasons, histological analysis was performed to confirm the presence of chytridiomycosis.

Besides *Rhinella quechua*, chytridiomycosis has been reported in other closely related new world Bufonidae (*R. marina*, Daszak et al. 2003; *Anaxyrus boreas*, Muths et al. 2003; and *A. canorus*, Green & Kagarise Sherman 2001). Wild populations of *R. marina* are not declining, even though chytrid fungi can infect and kill individuals (Daszak et al. 2003), but the declines of the 2 species of *Anaxyrus* in western North America (Carey 1993, Kagarise Sherman & Morton 1993) were related to the fungus (Green & Kagarise Sherman 2001, Muths et al. 2003, Carey et al. 2006). The differential effect that *Batrachochytrium dendrobatidis* has on these species seems to depend on the life history and distribution of the toads (Daszak et al. 2003). While *R. marina* is a lowland, prolific species that has been introduced worldwide, the 2 *Anaxyrus*

species are mountain-dwellers and endemic species. *R. quechua* is also a mountain-dweller occurring in humid montane forests (De la Riva et al. 2000), and its presence has been confirmed in only 2 areas of Bolivia (Santa Cruz and Cochabamba).

At present, no field data are available on the population status of *Rhinella quechua* or its susceptibility to chytridiomycosis, although its habitat and distribution patterns suggest that this species could be negatively affected by chytridiomycosis. In Bolivia, this disease is a potential threat, especially to some species that have been demonstrated to be highly susceptible to chytridiomycosis, like montane aquatic frogs of the genus *Telmatobius* (Merino-Viteri et al. 2005, Seimon et al. 2005, 2006, Barrionuevo & Mangione 2006) as well as in other species from the mountain regions of this megadiverse country.

Acknowledgements. We are indebted to Marisa Alcaide and Ana and Franco Pucci, Instituto de Morfología Animal, Fundación Miguel Lillo, for their help in the preparation of histological slides. This project and the collection of samples were supported by a DAPTF seed grant 2006 and a WWF Russell E. Train grant to R.A. for monitoring endangered species in Carasco National Park, Cochabamba, Bolivia.

LITERATURE CITED

- Alford RA, Bradfield KS, Richards SJ (2007) Global warming and amphibian losses. *Nature* 447:E3–E4
- Altig R (2007) Comments on the descriptions and evaluations of tadpole mouthpart anomalies. *Herpetol Conserv Biol* 2: 1–4
- Barrionuevo S, Mangione S (2006) Chytridiomycosis in two species of *Telmatobius* (Anura: Leptodactylidae) from Argentina. *Dis Aquat Org* 73:171–174
- Berger L, Speare R, Daszak P, Green DE and others (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proc Natl Acad Sci USA* 95: 9031–9036
- Berger L, Speare R, Kent A (1999) Diagnosis of chytridiomycosis in amphibians by histologic examination. www.jcu.edu.au/school/phtm/PHTM/frogs/histo/chhisto.htm
- Carey C (1993) Hypothesis concerning the causes of the disappearance of boreal toads from the mountains of Colorado. *Conserv Biol* 7:355–362
- Carey C, Bruzgul JE, Livo LJ, Walling ML and others (2006) Experimental exposures of boreal toads (*Bufo boreas*) to a pathogenic chytrid fungus (*Batrachochytrium dendrobatidis*). *EcoHealth* 3:5–21
- Daszak P, Cunningham AA, Hyatt AD (2003) Infectious disease and amphibian population declines. *Divers Distrib* 9: 141–150
- De la Riva I (2005) Bolivian frogs of the genus *Telmatobius* (Anura: Leptodactylidae): synopsis, taxonomic comments, and description of a new species. *Monogr Herpetol* 7: 65–101
- De la Riva I, Köhler J, Lötters S, Reichle S (2000) Ten years of research on Bolivian amphibians: updated checklist, distribution, taxonomic problems, literature and iconography. *Rev Esp Herpetol* 14:19–164
- De la Riva I, Aparicio J, Ríos N (2005) New species of *Telmatobius* (Anura: Leptodactylidae) from humid paramo of Peru and Bolivia. *J Herpetol* 39:409–416
- Di Rosa I, Simoncelli F, Fagotti A, Pascolini R (2007) The proximate cause of frog declines? *Nature* 447:E4–E5
- Fellers GM, Green DE, Longcore JE (2001) Oral chytridiomycosis in the mountain yellow-legged frog (*Rana muscosa*). *Copeia* 2001:945–953
- Green DE, Kagarise Sherman C (2001) Diagnostic histological findings in Yosemite toads (*Bufo canorus*) from a die-off in the 1970s. *J Herpetol* 35:92–103
- IUCN, Conservation International, and NatureServe (2006) Global amphibian assessment. www.globalamphibians.org
- Kagarise Sherman C, Morton ML (1993) Population declines of Yosemite toads in the eastern Sierra Nevada of California. *J Herpetol* 27:186–198
- Knapp RA, Morgan JAT (2006) Tadpole mouthpart depigmentation as an accurate indicator of chytridiomycosis, an emerging disease of amphibians. *Copeia* 2006:188–197
- Lips KR, Brem F, Brenes R, Reeve JD and others (2006) Emerging infectious disease and the loss of biodiversity in a neotropical amphibian community. *Proc Natl Acad Sci USA* 103:3165–3170
- Merino-Viteri A, Coloma LA, Almendáriz A (2005) Los *Telmatobius* (Leptodactylidae) de los Andes del Ecuador y su declive poblacional. *Monogr Herpetol* 7:9–37
- Muths E, Corn PS, Pessier AP, Green DE (2003) Evidence for disease related amphibian decline in Colorado. *Biol Conserv* 110:357–365
- Pessier AP, Nichols DK, Longcore JE, Fuller MS (1999) Cutaneous chytridiomycosis in poison dart frogs (*Dendrobates* spp.) and White's tree frogs (*Litoria caerulea*). *J Vet Diagn Investig* 11:194–199
- Pounds JA, Bustamante MR, Coloma LA, Consuegra JA and others (2007) Global warming and amphibian losses; the proximate cause of frog declines? (Reply). *Nature* 447:E5–E6
- Rachowicz LJ (2002) Mouthpart pigmentation in *Rana muscosa* tadpoles: seasonal changes without chytridiomycosis. *Herpetol Rev* 33:263–265
- Rachowicz LJ, Hero JM, Alford RA, Taylor JW and others (2005) The novel and endemic pathogen hypothesis: competing explanations for the origin of emerging infectious diseases of wildlife. *Conserv Biol* 19:1441–1448
- Reichle S (2003) Anexo 2. Lista de anfibios presentes en Bolivia. In: Ibsch PL, Mérida G (eds) Biodiversidad: la riqueza de Bolivia: estado de conocimiento y conservación. Editorial FAN, Santa Cruz de la Sierra, p 583–585
- Reichle S (2006) Distribution, diversity and conservation status of Bolivian amphibians. PhD dissertation, Rheinische Friedrichs-Wilhelm Universität, Bonn
- Ron SR (2005) Predicting the distribution of the amphibian pathogen *Batrachochytrium dendrobatidis* in the New World. *Biotropica* 37:209–221
- Seimon TA, Hoernig G, Sowell P, Halloy S, Seimon A (2005) Identification of chytridiomycosis in *Telmatobius marmoratus* at 4450 m in the Cordillera Vilcanota of Southern Peru. *Monogr Herpetol* 7:273–281
- Seimon TA, Seimon A, Daszak P, Halloy SRP, and 6 others (2006) Upward range extension of Andean anurans and chytridiomycosis to extreme elevations in response to tropical deglaciation. *Glob Change Biol* 13:288–299
- Skerratt LF, Berger L, Speare R, Cashins S and others (2007) Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* 4: 125–134