

REVIEW

Treatment of gyrodactylid infections in fish

Bettina Schelkle¹, Andrew P. Shinn², Edmund Peeler³, Joanne Cable^{1,*}

¹School of Biosciences, Cardiff University, Cardiff CF10 3AX, UK

²Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK

³Centre for Environment, Fisheries and Aquaculture Science, Barrack Road, The Nothe, Weymouth, Dorset DT4 8UB, UK

ABSTRACT: Since Norway experienced the devastating *Gyrodactylus salaris* (Monogenea) epidemics in Atlantic salmon *Salmo salar*, there has been heightened interest in how to treat gyrodactylosis in fish. Here we summarize chemical treatments previously used against gyrodactylids and discuss the main problems associated with these control measures including efficacy, host toxicity, human health concerns and application of treatments. Unfortunately, for these reasons and because of the different methodologies and different parasite and host species used in previous studies, it is difficult to recommend effective chemotherapeutic treatments. However, we suggest a method for manual removal of gyrodactylids from the host suitable for use in small-scale research facilities.

KEY WORDS: *Gyrodactylus* · Control treatment · Efficacy · Toxicity

—Resale or republication not permitted without written consent of the publisher—

INTRODUCTION

A frequently asked, but rarely addressed, question is what is the most effective means of removing *Gyrodactylus* spp. from their hosts? With increasing interest in this large group of ectoparasitic worms, not only as important pathogens but also as model organisms (reviewed by Bakke et al. 2002, 2007, Cable & Harris 2002), it seems timely to document the methods that have been used thus far to remove gyrodactylids.

PROBLEMS WITH EXISTING TREATMENTS

Appendix 1 lists 88 compounds and treatment combinations that have previously been used to treat gyrodactylid infections in research facilities, aquaculture and the hobbyist market. However, selecting the best of these treatments is difficult, because few studies have compared compounds using the same methodologies and the majority of treatments have various associated problems. Formaldehyde, for instance, was found to be 100% effective at eliminating *Gyrodactylus salaris* experimentally (Buchmann & Kristensson 2003), but under large-scale aquaculture conditions it

does not eradicate gyrodactylids completely (Rintamäki-Kinnunen & Valtonen 1996, Rowland et al. 2006), and in some three-spined stickleback *Gasterosteus aculeatus* populations it is only 10% effective (J. Cable pers. obs.). Due to its broad anti-parasitic properties, formaldehyde is still commonly used in aquaculture, even though it is classed as a human carcinogen (IARC 2004). Mutagenic and carcinogenic effects are also known for malachite green (Srivastava et al. 2004), which like formaldehyde is widely used as an anti-parasitic treatment, but its effectiveness against *Gyrodactylus* spp. has not been evaluated. Malachite green is now banned in food fish production in Europe and North America as it is retained in fish flesh (European Council 1990; US Food and Drug Administration and the Canadian Food Inspection Agency, cited in Srivastava et al. 2004).

Alternative treatments include rotenone, an indiscriminate ATPase inhibitor which has been used to control *Gyrodactylus salaris* in Norway by killing all potential hosts (reviewed in Bakke et al. 2007), including other gill-breathing organisms. This is only partially effective and has potential negative effects on human health. Aqueous aluminium is another alternative that is being tested. However, although aluminium ap-

*Corresponding author. E-mail: cablej@cardiff.ac.uk

peared promising in laboratory studies (Soleng et al. 1999, 2005, Poléo et al. 2004), field trials in Norway using aluminium in combination with rotenone have been problematic because successful treatment of the whole river system requires maintaining a specific concentration without exceeding levels toxic to Atlantic salmon (P. Shave pers. comm.). Aluminium toxicity to fish increases under acidic water conditions (Birchall et al. 1989, Poléo 1995, Soleng et al. 1999), and elevated environmental aluminium levels can result in Alzheimer's disease (Doll 1993) and decreased agricultural and forestry productivity (Bi et al. 2001). Organophosphates, such as trichlorfon, are no longer in use against fish parasites as they cause irreversible effects in non-target species by phosphorylating acetylcholinesterase (see Kozlovskaya & Mayer 1984, Peña-Llopis et al. 2003, Costa 2006). Acute and chronic toxicity to other aquatic organisms has also been reported for benzimidazoles (e.g. Oh et al. 2006); however, due to their low efficacy (see Appendix 1), these are not preferred over broad antiparasitic treatments such as formalin.

In addition to efficacy, environmental and human health issues, the main problem associated with current gyrodactylid treatments is their toxicity to the host (Schmahl & Taraschewski 1987, Santamarina et al. 1991, Tojo et al. 1992, Scholz 1999, Ekanem et al. 2004, Srivastava et al. 2004). Even widely used compounds, such as formaldehyde, may significantly change the host's gill structure and epidermis (Speare et al. 1997, Sanchez et al. 1998, Buchmann et al. 2004). For instance, although zinc exposure initially stimulates host mucus production, mucus is subsequently depleted leaving the fish more susceptible to microbial infections (McGeer et al. 2000). If these fish are subsequently used for experimental infections without a sufficient recovery period, they may show an abnormal response to infection (see review by Bakke et al. 2007). Host respiratory problems are also a common side effect of gyrodactylid treatments due to direct interference with gill function and indirectly via reduction of water quality (e.g. decrease in pH and/or dissolved oxygen: Smith & Piper 1972, Ross et al. 1985, Rowland et al. 2006). Toxicity in many cases is dose-dependent, but may also be affected by one or a combination of the following factors: temperature, pH, salinity, mechanism of delivery, species and exposure time (Schmahl & Taraschewski 1987, Schmahl et al. 1989, Scholz 1999), especially where multiple treatments are used.

Gyrodactylid treatments are applied either orally (with food) or topically (added to the water). Both methods usually lead to application of overly high doses to compensate for lack of control over drug administration (Scholz 1999), which may lead to environmental contamination. Oral administration of some drugs can also reduce host food consumption (e.g.

piperazine, Fugotenil® and Neguvon®; Tojo & Santamarina 1998), which increases the dosage needed per unit of food for efficacy. Additionally, compounds that are partially effective as baths may not be effective if administered orally (e.g. trichlorfon; Santamarina et al. 1991, Tojo & Santamarina 1998). Moreover, resistance of parasites against anthelmintics has been reported, with Goven & Amend (1982) and Schmahl et al. (1989) finding resistance of gyrodactylids to dimethyl phosphonate and trichlorfon, respectively. Breeding parasite-resistant fish or developing a vaccine against gyrodactylosis would override many of the above problems and bypass the potential issue of drug resistance. Breeding experiments with wild Atlantic salmon *Salmo salar* that show resistance against *Gyrodactylus salaris* are currently under way in Norway (R. Salte et al. unpubl.), but we are unaware of any plans for vaccine development. In Norway destocking and drying of fish farms was used successfully to eradicate *G. salaris* (see Mo 1994); however, this method is not suitable for establishments which receive potentially infected fish on a regular basis and infected fish would still have to be treated.

A major problem with many of the studies presented in Appendix 1 is that they used sub-sampling methods, such as mucus scrapings (e.g. Tojo et al. 1992, Tojo & Santamarina 1998) or partial examination of fish populations in field trials (e.g. Goven & Amend 1982). This is a crude and unreliable estimate of parasite infection, particularly where the gyrodactylid is not identified to species level (e.g. Tojo & Santamarina 1998, Chansue 2007) as different *Gyrodactylus* spp. show marked site specificities. Furthermore, the majority of compounds trialled successfully have been tested only once. For those with at least 2 independent studies, results have been variable probably because of different methodologies. Hence, many seemingly successful compounds may not be effective when it comes to treating a different host-parasite system. Field studies are of particular importance as they provide an indication of how efficient treatments are on a large scale, potentially for use in the ornamental or food fish industries. However, there have only been a few such studies (see Appendix 1) with only Rach et al. (2000) claiming 100% efficiency for the compound they tested (hydrogen peroxide). Other authors defined a treatment in the field as successful if it just reduced the pathogen burden (e.g. Lewis 1967, Rintamäki-Kinnunen & Valtonen 1996).

MANUAL REMOVAL OF GYRODACTYLIDS

Due to toxicity and/or difficulties in administration (such as the need for prolonged exposure), no treat-

ment detailed in Table 1 is entirely satisfactory or 100% effective, although several treatments indicate 100% efficacy against gyrodactylosis (e.g. formaldehyde and aqueous aluminium). In laboratory trials with small, hardy fish, the most effective means of removing gyrodactylids (at least those that predominantly occur on the skin and fins) is a licensed Home Office procedure in the UK which requires the host to be lightly anaesthetised (using e.g. 0.02% MS-222, chlorbutanel) whilst the living parasites are removed manually. Following anaesthesia, the fish is transferred to a shallow dish, but kept fully immersed in dechlorinated water and screened using a stereo-microscope with fibre-optic illumination. The latter is essential to prevent the host and parasites overheating during examination on the microscope stage. Parasites can be most easily detected by their movement, but it is still essential to scan all surfaces of the host. Small fish, such as guppies and sticklebacks, can be manipulated using a plastic disposable pipette tip. Once a parasite is detected, the worm can be removed and crushed using watchmaker's forceps. It is essential to ensure that the parasite is not released directly back into the water as even a damaged worm may re-attach to its host and successfully give birth. Once all parasites have been removed, the same screening procedure should be repeated until all fish in a population have been screened clear of parasites on 3 consecutive occasions separated by approximately 1 wk, ideally by 2 independent, trained researchers. If a parasite is found on any fish during this process, then the screening process should begin again for the entire host population.

For small fish, such as poeciliids, manual removal of parasites can be effective without any chemical intervention. However, for larger fish (or where gyrodactylids are abundant and/or show a preference for the gills), chemical treatment prior to screening may be the only practical solution to parasite removal. In our laboratories we use levamisole, but this is only available by veterinarian prescription and only reliable if applied to single fish that are closely monitored for signs of toxicity. Screening is still essential to ensure that fish are free of gyrodactylids following chemical treatment, with additional screening on subsequent occasions to ensure the experimental fish are free from infection. For larger fish, such as adult Atlantic salmon *Salmo salar*, sub-sampling by examining mucus scrapings or selected fins may be the only practical way of screening, but this is not accurate. It is also stressful for the fish as it damages the fish's epithelium, so increasing the risk of secondary infections. An alternative or ideally supplementary procedure to manual removal, but again only suitable for small fish, is to maintain each fish in isolation in a closed system with regular water changes. As fish become immune and gyro-

dactylids are shed, there is no opportunity for the parasites to transfer to new hosts. However, even long periods of isolation are not always effective for certain host-parasite combinations (see King et al. 2008) and the accompanying high maintenance is extremely time consuming and costly. Whatever the control method used (chemical, manual or host isolation), thorough screening is essential to ensure parasite extinction. The longer the interval between the 3 consecutive screens the more reliable the procedure. A single (transparent, <1 mm long) gyrodactylid missed in an earlier screen will have had ample time to reproduce *in situ* (as little as 24 h at 25°C) generating a larger number of parasites which are less likely to be overlooked on subsequent screens.

CONCLUSION

Although the majority of compounds tested against *Gyrodactylus* spp. are reportedly effective (see Appendix 1), 100% efficacy has not been achieved without toxicity to hosts. Currently, the lack of comparative data makes the selection of a drug difficult. Leaving just one (hermaphrodite, viviparous) worm can be sufficient to initiate a new disease outbreak. For research projects, manual removal of skin gyrodactylids from small fish is an option; however, this is not feasible for gill parasites, larger hosts or for use in aquaculture. Further research into different treatments and their application on different species of parasite and host is necessary to combat the existing problems caused by *Gyrodactylus* spp. epizootics.

Acknowledgements. This work was supported by the Natural Environment Research Council, UK (NER/J/S/2002/00706).

LITERATURE CITED

- Anttila P, Kuusela J, Koski P (2007) Disinfection of *Gyrodactylus salaris* by Virkon S and heat. Poster presentation at the 13th International Conference of Fish and Shellfish Diseases by the European Association of Fish Pathologists 17–21 September 2007, Grado, Italy. EAAP Abstract Book, p 334. Available at: <http://eafp.org/eafp-information/2009/2/20/grado-2007-information-archived.html>
- Bakke TA, Harris PD, Cable J (2002) Host specificity dynamics: observations on gyrodactylid monogeneans. *Int J Parasitol* 32:281–308
- Bakke TA, Cable J, Harris PD (2007) The biology of gyrodactylid monogeneans: the 'Russian-doll killers'. *Adv Parasitol* 64:161–376
- Bi SP, Yang XD, Zhang FP, Wang XL, Zou GW (2001) Analytical methodologies for aluminium speciation in environmental and biological samples—a review. *Fresenius J Anal Chem* 370:984–996
- Birchall JD, Exley C, Chappell JS, Phillips MJ (1989) Acute toxicity of aluminium to fish eliminated in silicon-rich acid waters. *Nature* 338:146–148

- Buchmann K (1997) Salinity tolerance of *Gyrodactylus derjavini* from rainbow trout *Oncorhynchus mykiss*. Bull Eur Assoc Fish Pathol 17:123–125
- Buchmann K, Kristensson RT (2003) Efficacy of sodium percarbonate and formaldehyde bath treatments against *Gyrodactylus derjavini* infestations of rainbow trout. N Am J Aquac 65:25–27
- Buchmann K, Bresciani J, Jappe C (2004) Effects of formalin treatment on epithelial structure and mucous cell densities in rainbow trout, *Oncorhynchus mykiss* (Walbaum), skin. J Fish Dis 27:99–104
- Cable J, Harris PD (2002) Gyrodactylid developmental biology: historical review, current status and future trends. Int J Parasitol 32:255–280
- Caboni P, Sherer TB, Zhang N, Taylor G, Na HM, Greenamyre JT, Casida JE (2004) Rotenone, deguelin, their metabolites and the rat model of Parkinson's disease. Chem Res Toxicol 17:1540–1548
- Carter PKL (2003) The impact of low concentrations of cadmium on host–monogenean interactions. PhD dissertation, University of Stirling
- Chansue N (2007) Effects of dried Indian almond leaf (*Terminalia catappa* L.) extract on monogenean parasites in goldfish (*Carassius auratus*). Wien Tierärztl Mon Schr 94: 269–273
- Costa LG (2006) Current issues in organophosphate toxicology. Clin Chim Acta 366:1–13
- Crigel P, Losson B, Defour J (1995) Utilisation de la quinaldine, un anesthésique pour poissons, à des fins antiparasitaires. Ann Med Vet 139:343–348
- Doll R (1993) Review: Alzheimer's disease and environmental aluminium. Age Ageing 22:138–153
- Ekanem AP, Wang M, Simon JE, Obiekezie AI, Morah F (2004) *In vivo* and *in vitro* activities of the seed extract of *Piper guineense* Schum and Thonn against skin and gill monogenean parasites of goldfish (*Carassius auratus auratus*). Phytother Res 18:793–797
- El-Khatib NRH (2003) Biological eradication of some parasitic diseases in fishes using *Bacillus thuringiensis* Agerin® product. Vet Med J Giza 51:19–28
- European Council (1990) Regulation No 2377/90 of 26 June 1990 laying down a community procedure for the establishment of maximum residue limits for veterinary medicinal products in foodstuffs of animal origin. Off J Eur Union L 224:1–8
- FishDoc (2008) Malachite green and formalin—a good general-purpose anti-parasite treatment. www.fishdoc.co.uk/treatments/malachite.htm (accessed 15 Dec 2008)
- Gheorgiu C, Marcogliese DJ, Scott M (2005) Concentration-dependent effects of waterborne zinc on population dynamics of *Gyrodactylus turnbulli* (Monogenea) on isolated guppies (*Poecilia reticulata*). Parasitology 132:225–232
- Goven BA, Amend DF (1982) Mebendazole/trichlorfon combination: a new anthelmintic for removing monogenetic trematodes from fish. J Fish Biol 20:373–378
- Harris PD, Shinn AP, Cable J, Bakke TA, Bron J (2008) GyroDb: gyrodactylid monogeneans on the web. Trends Parasitol 24:109–111
- IARC (International Agency for Research on Cancer) (2004) Monographs on the evaluation of carcinogenic risks to humans, Vol 88, formaldehyde, 2-butoxyethanol and 1-tert-butoxy-2-propanol. IARC, Lyons
- King TA, Harris PD, Cable J (2008) Long-term *Gyrodactylus lomi* infection on isolated juvenile chub, *Leuciscus cephalus*. J Parasitol 94(6):1426–1427
- Kou GH, Liu CI, Liu CK (1988) Handbook of fish disease. Pig Research Institute Taiwan, Miaoli (in Chinese)
- Kozlovskaya VI, Mayer FL (1984) Brain acetylcholinesterase and backbone collagen in fish intoxicated with organophosphate pesticides. J Gt Lakes Res 10:261–266
- Lewis SD (1967) Prophylactic treatment of minnow hatchery ponds with paraformaldehyde to prevent epizootics of *Gyrodactylus*. Prog Fish-Cult 29:160–161
- Liao C, Guo JJ, Su MS (1996) The use of chemicals in aquaculture in Taiwan, province of China. In: Arthur JR, Lavilla-Pitogo CR, Subasinghe RP (eds) Proceedings of the meeting on the use of chemicals in aquaculture in Asia, 20–22 May 1996, Tigbauan, Iloilo. SEAFDEC, Philippines, p 193–206
- McGeer JC, Szedebinsky C, McDonald DG, Wood CM (2000) Effects of chronic sublethal exposure to waterborne Cu, Cd or Zn in rainbow trout. 1: iono-regulatory disturbance and metabolic costs. Aquat Toxicol 50:231–234
- Mo TA (1994) Status of *Gyrodactylus salaris* problems and research in Norway. In: Pike AW, Lewis JW (eds) Parasitic diseases of fish. Samara Publishing, Tresaith, p 43–56
- Oh SJ, Park J, Lee MJ, Park SY, Lee JH, Choi K (2006) Ecological hazard assessment of major veterinary benzimidazoles: acute and chronic toxicities to aquatic microbes and invertebrates. Environ Toxicol Chem 25:2221–2226
- Peña-Llopis S, Ferrando MD, Peña JB (2003) Fish tolerance to organophosphate-induced oxidative stress is dependent on the glutathione metabolism and enhanced by N-acetylcysteine. Aquat Toxicol 65:337–360
- Pesticide Action Network UK (2001) Rotenone. Pesticides News 54:20–21
- Pettersen RA, Vøllestad LA, Flodmark LEW, Poléo ABS (2006) Effects of aqueous aluminium on four fish ectoparasites. Sci Total Environ 369:129–138
- Poléo ABS (1995) Aluminium polymerization—a mechanism of acute toxicity of aqueous aluminium of fish. Aquat Toxicol 31:347–356
- Poléo ABS, Schjolden J, Hansen H, Bakke TA, Mo TA, Rosse-land BO, Lydersen E (2004) The effect of various metals on *Gyrodactylus salaris* (Platyhelminthes, Monogenea) infections in Atlantic salmon (*Salmo salar*). Parasitology 128:169–177
- Rach JJ, Gaikowski MP, Ramsay RT (2000) Efficacy of hydrogen peroxide to control parasitic infestations on hatchery-reared fish. J Aquat Anim Health 12:267–273
- Rintamäki-Kinnunen P, Valtonen ET (1996) Finnish salmon resistant to *Gyrodactylus salaris*: a long-term study at fish farms. Int J Parasitol 26:723–732
- Ross LG, Ward KMH, Ross B (1985) The effects of formalin, malachite green and suspended solids on the respiratory activity of rainbow trout, *Salmo gairdneri* Richardson. Aquac Res 16:129–138
- Rowland ST, Nixon M, Landos M, Mifsud C, Read P, Boyd P (2006) Effects of formalin on water quality and parasitic monogeneans on silver perch (*Bidyanus bidyanus* Mitchell) in earthen ponds. Aquac Res 37:869–876
- Russo R, Curtis EW, Yanong RPE (2007) Preliminary investigations of hydrogen peroxide treatment of selected ornamental fishes and efficacy against external bacteria and parasites in green swordtails. J Aquat Anim Health 19:121–127
- Sanchez JG, Speare DJ, Sims DE, Johnson GJ (1998) Morphometric assessment of epidermal and mucous-biofilm changes caused by exposure of trout to chloramine-T or formalin treatment. J Comp Pathol 118:81–87
- Santamarina MT, Tojo J, Ubeira FM, Quinteiro P, Sanmartín ML (1991) Anthelmintic treatment against *Gyrodactylus* sp. infecting rainbow trout *Oncorhynchus mykiss*. Dis Aquat Org 10:39–44

- Schmahl G (1993a) Treatment of fish parasites. 10. Effects of a new triazine derivative, HOE-092-V, on Monogenea—a light and transmission electron-microscopy study. *Parasitol Res* 79:559–566
- Schmahl G (1993b) Up to date chemotherapy against monogenea: a review. *Bull Fr Pêche Piscic* 328:74–81
- Schmahl G, Mehlhorn H (1988) Treatment of fish parasites. 4. Effects of sym. triazinone (toltrazuril) on Monogenea. *Parasitol Res* 75:132–143
- Schmahl G, Taraschewski H (1987) Treatment of fish parasites 2. Effects of praziquantel, niclosamide, levamisole, hydrochloride and metrifonate on Monogenea (*Gyrodactylus aculeati* and *Diplozoon paradoxum*). *Parasitol Res* 73:341–351
- Schmahl G, Taraschewski H, Mehlhorn H (1989) Chemotherapy of fish parasites. *Parasitol Res* 75:503–511
- Scholz T (1999) Parasites in feral and cultured fish. *Vet Parasitol* 84:317–355
- Smith CE, Piper RG (1972) Pathological effects in formalin treated rainbow trout (*Salmo gairdneri*). *J Fish Res Board Can* 29:328–329
- Soleng A, Bakke TA (1997) Salinity tolerance of *Gyrodactylus salaris* (Platyhelminthes, Monogenea): laboratory studies. *Can J Fish Aquat Sci* 54:1837–1864
- Soleng A, Bakke TA, Hansen LP (1998) Potential for dispersal of *Gyrodactylus salaris* (Platyhelminthes, Monogenea) by sea-running stages of the Atlantic salmon (*Salmo salar*): field and laboratory studies. *Can J Fish Aquat Sci* 55:507–514
- Soleng A, Poléo ABS, Alstad NEW, Bakke TA (1999) Aqueous aluminium eliminates *Gyrodactylus salaris* (Platyhelminthes, Monogenea) infections in Atlantic salmon. *Parasitology* 119:19–25
- Soleng A, Poléo ABS, Bakke TA (2005) Toxicity of aqueous aluminium to the ectoparasitic monogenean *Gyrodactylus salaris*. *Aquaculture* 250:616–620
- Speare DJ, Arsenault G, MacNair N, Powell MD (1997) Branchial lesions associated with intermittent formalin bath treatment of Atlantic salmon, *Salmo salar* L., and rainbow trout, *Oncorhynchus mykiss* (Walbaum). *J Fish Dis* 20:27–33
- Srivastava S, Sinha R, Roy D (2004) Toxicological effects of malachite green. *Aquat Toxicol* 66:319–329
- Steverding D, Morgan E, Tkaczynski P, Walder F, Tinsley R (2005) Effect of Australian tea tree oil on *Gyrodactylus* spp. infection of the three-spined stickleback *Gasterosteus aculeatus*. *Dis Aquat Org* 66:29–32
- Tojo JL, Santamarina MT (1998) Oral pharmacological treatments for parasitic diseases of rainbow trout *Oncorhynchus mykiss*. II. *Gyrodactylus* sp. *Dis Aquat Org* 33:187–193
- Tojo J, Santamarina MT, Ubeira FM, Estevez J, Sanmartin ML (1992) Anthelmintic activity of benzimidazoles against *Gyrodactylus* sp. infecting rainbow trout *Oncorhynchus mykiss*. *Dis Aquat Org* 12:185–189
- Tojo J, Santamarina MT, Ubeira FM, Leiro J, Sanmartin ML (1993a) Efficacy of antiprotozoal drugs against gyrodactylosis in rainbow trout (*Oncorhynchus mykiss*). *Bull Eur Assoc Fish Pathol* 13:79–82
- Tojo J, Santamarina MT, Ubeira FM, Estevez J, Leiro J, Sanmartin ML (1993b) Efficacy of anthelmintic drugs against gyrodactylosis in rainbow trout (*Oncorhynchus mykiss*). *Bull Eur Assoc Fish Pathol* 13:45–48

Appendix 1. Treatments against infections of *Gyrodactylus* species. With the exception of orally administered treatments, other compounds below were applied directly to the water

Compound	Dose	Application	<i>Gyrodactylus</i> spp.	Host	Efficacy	Source
(1) Botanicals						
Rotenone	Used extensively to treat <i>G. salaris</i> on salmonids in Norwegian rivers. Kills both fish and parasites. Linked to Parkinson's disease. ^a					Pesticide Action Network UK (2001), Harris et al. (2008), Caboni et al. (2004) Chansue (2007)
<i>Terminalia catappa</i> extract	1.7, 3.4 & 5.1 g l ⁻¹	4 wk. Lab study	<i>Gyrodactylus</i> sp.	<i>Carassius auratus</i>	Not effective (1.7 g l ⁻¹) to 100% effective after 14 d (5.1 g l ⁻¹) ^b	
Tea tree oil and Tween 80	3, 10 & 30 ppmv in 0.01% Tween	48 h. 13°C. Lab study	<i>Gyrodactylus</i> spp. (probably <i>G. gasterostei</i> and/or <i>G. arcuatus</i>)	<i>Gasterosteus aculeatus</i>	Reduced parasite mean from 12.6 worms fish ⁻¹ to 1.4 (30 ppmv) ^b	Steverding et al. (2005)
	0.01% Tween				Reduced parasite mean from 12.6 worm fish ⁻¹ and reduced prevalence by 45% ^b	
<i>Piper guineense</i> extract	0.5, 1.0, 1.5 & 2 mg l ⁻¹	96 h. Test solution renewed every 24 h. Lab study	<i>G. elegans</i>	<i>Carassius auratus</i> (10 fish treatment ⁻¹)	Dose dependent, but none 100% effective ^{a,b,c}	Ekanem et al. (2004)
(2) General chemotherapeutics						
Acaprin (Acaprina®)	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	96% effective ^{d,e}	Tojo et al. (1993a)
Agerin®	10, 500 & 1000 ppm	Indefinite time, 2 & 1 h, respectively. 18°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oreochromis niloticus</i>	Not effective ^c	El-Khatib (2003)
Albendazole (Oversol®)	25 & 200 mg l ⁻¹	12 & 3 h, respectively. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective (200 mg l ⁻¹) to 95.5% effective (25 mg l ⁻¹) ^{a,d,e}	Tojo et al. (1992)
Aminoisidine (Gabbrocol®)	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Aminoisidine (Gaboral®)	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
Amprolium (Prolsal®)	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Benznidazole	200 mg l ⁻¹	3 h. 11°C. Lab. study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Bithionol	4, 20 & 60 mg l ⁻¹	3 h. 15°C. Lab. study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	98.6% effective (4 mg l ⁻¹) to 100% effective (20 & 60 mg l ⁻¹) ^{a,d,e}	Santamarina et al. (1991)
	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Caridazole (Spartrix®)	10 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
Chloroquine	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Chloroquine diphosphate	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
Closantel (Flukiver®)	0.25 & 0.125 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	99.6% effective (0.125 mg l ⁻¹) to 100% effective (0.25 mg l ⁻¹) ^{a,d,e}	Santamarina et al. (1991)

Appendix 1 (continued)

Compound	Dose	Application	<i>Gyrodactylus</i> spp.	Host	Efficacy	Source
Diethylcarbamazine	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993b)
	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^d	Tojo & Santamarina (1998)
Dimetridazole	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
Diminazene aceturate (Berenil [®])	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	98.6% effective ^{d,e}	Tojo et al. (1993a)
Febantel (Rintal [®])	2 & 10 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	Not effective (2 mg l ⁻¹) to toxic to host (10 mg l ⁻¹ ; 4 trout dead after 1 h, expt. terminated) ^{a,d,e}	Santamarina et al. (1991)
Fenbendazole (Panacur [®])	0.77, 1.5, 6.2, 12.5 & 25 mg l ⁻¹	12 h (additionally, 3 h for 25 mg l ⁻¹) 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective (0.77 mg l ⁻¹ & 25 mg l ⁻¹ for 3 h) to 100% effective (>1.5 mg l ⁻¹) ^{a,d,e}	Tojo et al. (1992)
Flubendazole	25 & 200 mg l ⁻¹	12 & 3 h, respectively. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^{a,e}	Tojo et al. (1992)
Formaldehyde	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^e	Tojo & Santamarina (1998)
Formalin	2.5, 5, 10, 20, 40 & 60 mg l ⁻¹	18 h. Lab study	<i>G. aculeati</i>	<i>Gasterosteus aculeatus</i>	Not effective (2.5 mg l ⁻¹) to 100% effective (20, 40 & 60 mg l ⁻¹) ^g	Buchmann & Kristensson (2003)
	20, 25, 30 & 40 mg l ⁻¹	24.1–26.9°C or 13.2–15.7°C. Field trial: applied to ponds	<i>Gyrodactylus</i> sp.	<i>Bidyanus bidyanys</i>	Not effective (20, 25 mg l ⁻¹) to sig. decrease (30, 40 mg l ⁻¹) ^{b,c}	Rowland et al. (2006)
Formalin; malachite green & formalin; salt bath	0.2 g l ⁻¹ ; 4 g l ⁻¹ & 0.2 g l ⁻¹ ; 1.5–2%, respectively	Farm study: rotation of treatments	Putatively <i>G. salaris</i>	<i>Salmo salar</i> , <i>S. trutta trutta</i> & <i>S. t. lacustris</i>	Formalin partially effective (decreased parasite burden); malachite green + formalin, and/or salt not effective ^{b,g}	Rintamäki-Kinnunen & Valtonen (1996)
Formalin (For) & malachite green (MG)	Recommended use in combination: 3.68 g MG in 1 l For, used at 0.025 ml l ⁻¹ of water for 60 min, or 3.3 g MG in 1 l For used at 0.015 ml l ⁻¹ of water as a prolonged immersion. Synergistic effect of compounds in combination, but toxic to both parasite and host					Srivastava et al. (2004), Fishdoc (2008)
HOE 092 V (triazine derivative)	0, 2, 5, 10 & 15 µg l ⁻¹	1, 2, 3 & 4 h. 22°C. Lab study	<i>G. arcuatus</i>	<i>Gasterosteus aculeatus</i>	Not effective (0–5 µg l ⁻¹) to effective after 4 h (10 µg l ⁻¹) & 1 h (15 µg l ⁻¹) ^{a,b,c}	Schmahl (1993a)
Hydrogen peroxide	10 mg l ⁻¹ 170, 280 & 560 mg l ⁻¹	3 h 3 × 30 min over 5 d. Field trials	<i>G. groschaffi</i> <i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i>	Effective ^b 100% effective ^{b,c}	Cited in: Schmahl (1993b) Rach et al. (2000)
Ivermectin (Ivomec [®])	11 & 16 mg l ⁻¹ ; 3 & 6 mg l ⁻¹	1 h; 24 h. 23.5°C. Lab study	<i>Gyrodactylus</i> spp.	<i>Xiphophorus helleri</i> (51–75 fish treatment ⁻¹)	Not effective ^{b,c}	Russo et al. (2007)
	0.031 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	100% host mortality ^{a,c,e}	Santamarina et al. (1991)
	0.1 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	96% effective ^{d,e}	Tojo et al. (1993b)
Ivermectin + closulon (Ivomec-F [®])	0.025, 0.25, 0.05 & 0.5 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	59% effective (0.025 & 0.25 mg l ⁻¹) to 100% effective (0.05 & 0.5 mg l ⁻¹) ^{a,d,e}	Santamarina et al. (1991)
Ketoconazole	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)

Appendix 1 (continued)

Compound	Dose	Application	Host	Efficacy	Source
Levamisol hydrochloride	0, 10, 20, 50 & 100 µg l ⁻¹	25, 30, 60, 90 & 120 min.	<i>G. aculeati</i>	Decrease of parasite burden ^{b,c}	Schmahl & Taraschewski (1987)
	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	Not effective ^f	Tojo & Santamarina (1998)
Levamisol hydrochloride (Citarin-L®)	100 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	Not effective ^{a,e}	Santamarina et al. (1991)
Malachite green	Recommended use: 0.5–0.8 ppm for 1 d, or 0.015 ppm for prolonged immersion. Remains in fish tissue for up to 1 mo. Highly toxic to fish: carcinogenic, mutagenic, chromosomal fractures, teratogenicity and respiratory toxicity. Now banned in the Europe and N. America for food fish. Most effective used together with formalin. ^a				Kou et al. (1988), Liao et al. (1996), Srivastava et al. (2004)
Mebendazole	0.1 mg l ⁻¹	24 h. 23°C. Lab study	<i>G. elegans</i>	100% effective ^{b,c}	Goven & Amend (1982)
	25 & 100 mg l ⁻¹	12 & 3 h, respectively. 15°C. Lab study	<i>G. salaris</i>	Not effective (100 mg l ⁻¹) to 95% effective (25 mg l ⁻¹) ^{a,e}	Tojo et al. (1992)
Mebendazole (Meb) & trichlorfon (Tri)	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	Not effective ^f	Tojo & Santamarina (1998)
	1 mg l ⁻¹ 0.01 mg l ⁻¹ 0.1–0.4 mg l ⁻¹ Meb & 0.46–1.8 mg l ⁻¹ Tri 0.5 mg l ⁻¹ Meb & 2.3 mg l ⁻¹ Tri	24 h 24 h Field tests.	<i>G. bullatarudis</i> <i>G. elegans</i> <i>G. elegans</i>	Effective ^{g,h} Effective ^{g,h} 100% effective ^{b,c}	Cited in: Schmahl (1993b) Cited in: Schmahl (1993b) Goven & Amend (1982)
Meglumine (Glucantime®)	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	Not effective ^e	Tojo et al. (1993a)
Methylene blue	Recommendations: 8–10 ppm, bath 1 d; 2 ppm, bath 3–5 d. Highly toxic to fish.				Liao et al. (1996)
Metronidazole	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	Not effective ^e	Tojo et al. (1993a)
Metronidazole (Flagyl®)	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	Not effective ^f	Tojo & Santamarina (1998)
	20 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	Not effective ^{d,e}	Tojo et al. (1993b)
Netobimin (Hapasi®)	2 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	Not effective ^{a,e}	Santamarina et al. (1991)
Niclofolan (Bilevon®)	0.025, 0.05 & 0.2 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	91% effective (0.025 mg l ⁻¹) to 100% effective (0.05 & 0.2 mg l ⁻¹ , but all host fish dead) ^{a,c,e}	Santamarina et al. (1991)
	0, 0.05, 0.075 & 0.1 µg l ⁻¹	30, 60, 90 & 120 min. 20°C. Lab study	<i>G. aculeati</i>	Reduction of parasite burden ^b	Schmahl & Taraschewski (1987)
Niclosamide (Fugotemil®)	6 & 1.5 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	100% effective ^{d,e}	Tojo et al. (1993b)
	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	Not 100% effective ^f	Tojo & Santamarina (1998)
Niridazole (Ambilhar®)	100 & 200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	Not effective (100 mg l ⁻¹) to 100% effective (200 mg l ⁻¹) ^e	Tojo et al. (1993b)

Appendix 1 (continued)

Compound	Dose	Application	<i>Gyrodactylus</i> spp.	Host	Efficacy	Source
Nitroscanate (Lopato1.500®)	40, 6.25, 0.63, 6.25 g kg ⁻¹ feed	10 d, 2 d, 1 d, 1 d oral treatment, respectively. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	100% effective ^f	Tojo & Santamarina (1998)
Nitroscanate (Lopato1®)	0.04, 0.07 & 156.25 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	99% effective (0.04 mg l ⁻¹) to 100% effective (0.07, 156.25 mg l ⁻¹) ^{a,e}	Santamarina et al. (1991)
Nitroxynil (Distomicine®)	30 & 50 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective (30 mg l ⁻¹) to 100% effective (50 mg l ⁻¹) ^{d,e}	Tojo et al. (1993b)
Nitroxynil	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Oxfendazole	25 & 200 mg l ⁻¹	3 h. 15°C. Lab. study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^{a,e}	Tojo et al. (1992)
Oxibendazole	25 & 200 mg l ⁻¹	12 & 3 h, respectively. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective (200 mg l ⁻¹ , 3 h) to 86% effective (25 mg l ⁻¹ , 12 h) ^{a,d,e}	Tojo et al. (1992)
Parafomaldehyde	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Parbendazole	Ca. 10 g l ⁻¹	Field trial: ponds sized 0.1–0.5 acre (0.04–0.2 ha), ~3 ft (1 m) deep	<i>G. elegans</i>	<i>Notemigonus crysoleucas</i>	Treatment was successful ^{h,g}	Lewis (1967)
Piperazine	25 & 200 mg l ⁻¹	12 & 3 h, respectively. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective (200 mg l ⁻¹ , 3 h) to 91% effective (25 mg l ⁻¹ , 12 h) ^{a,e}	Tojo et al. (1992)
Piperazine citrate	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Piperazine dihydrochloride	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Praziquantel	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	Not effective ^{a,e}	Santamarina et al. (1991)
Praziquantel (Droncit®)	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	Not effective ^{a,e}	Santamarina et al. (1991)
Praziquantel	0, 1, 5, 10, 20 & 50 µg l ⁻¹	15, 30, 60, 90, 120 min, 4 h & 16 h, 20°C. Lab study	<i>G. aculeati</i>	<i>Gasterosteus aculeatus</i> (10 fish treatment ⁻¹)	Decrease in parasite burden ^{b,d}	Schmahl & Taraschewski (1987)
Pyrantel pamoate (Trilombrin®)	10 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	98% effective ^{a,e}	Santamarina et al. (1991)
Quinacrine HCl (Atabrine®)	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Quinaldine	50 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993b)
Rafoxanide (Ramide®)	25, 100 & 200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	98% effective (25 mg l ⁻¹) to 100% effective (100, 200 mg l ⁻¹) ^{d,e}	Tojo et al. (1993a)
	0, 26, 104 & 260 mg l ⁻¹	Ca. 20, 40 & 60 s. 20–22°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Carassius auratus</i> , <i>Cyprinus auratus</i>	Effective (26 mg l ⁻¹ , ca. 60 s, time needed to anaesthetise fish) ^{b,c}	Crigel et al. (1995)
	50 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993b)

Appendix 1 (continued)

Compound	Dose	Application	<i>Gyrodactylus</i> spp.	Host	Efficacy	Source
Ronidazole	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^d	Tojo & Santamarina (1998)
Secnidazole	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	84% effective ^e	Tojo et al. (1993a)
Sodium percarbonate	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Sodium sulphha-quinoxaline Suramin (Nagano [®])	10, 20, 40, 80 & 160 mg l ⁻¹	18 h. Lab study	<i>G. derjavini</i> (= <i>G. derjavinoidea</i>)	<i>Oncorhynchus mykiss</i>	Not effective (10, 20 mg l ⁻¹) to 100% effective (80, 160 mg l ⁻¹) ^g	Buchmann & Kristensson (2003)
Sodium	200 mg l ⁻¹	3 h. 11°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993a)
Thiophanate (Triasox [®])	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993b)
Thiophanate (Nermafax 20 [®])	100 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	95% effective ^{d,e}	Tojo et al. (1993b)
Thiophanate	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Toltrazuril	10 & 100 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^{a,d,e}	Tojo et al. (1992)
Thiophanate	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective ^e	Tojo et al. (1993b)
Thiophanate	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Toltrazuril	5, 10, 20, 30 & 50 µg l ⁻¹	0.3, 1, 2, 3, 4 & 6 h. 20°C. Lab study	<i>G. arcuatus</i>	<i>Gasterosteus aculeatus</i> (10–15 fish treatment ⁻¹)	Not effective (5, 10 µg l ⁻¹) to effective after 2 h (50 µg l ⁻¹) ^{a,b,c}	Schmahl & Mehlhorn (1988)
Trichlorfon	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective	Tojo et al. (1993a)
Trichlorfon (Metrifonate)	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Trichlorfon (Neguvon [®])	Up to 2 mg l ⁻¹	24 h. 23°C. Lab study	<i>G. elegans</i>	<i>Carassius auratus</i> (10 fish treatment ⁻¹)	Not effective ^{b,c}	Goven & Amend (1982)
Trichlorfon	0.25 mg l ⁻¹	Continuous	<i>Gyrodactylus</i> sp.			Cited in: Schmahl (1993b)
Trichlorfon (Metrifonate)	0, 10 & 50 µg l ⁻¹	25, 30, 60 & 90 min. 20°C. Lab study	<i>G. aculeati</i>	<i>Gasterosteus aculeatus</i> (10 fish treatment ⁻¹)	Not effective ^g	Schmahl & Taraschewski (1987)
Trichlorfon (Neguvon [®])	200 mg l ⁻¹	3 h. 15°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (4–10 fish treatment ⁻¹)	98% effective ^{a,e}	Santamarina et al. (1991)
Triclabendazol (Fasinex [®])	40 g kg ⁻¹ feed	10 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Not effective ^f	Tojo & Santamarina (1998)
Triclabendazol	6.2, 12.5 & 25 mg l ⁻¹	12 h (additionally, 25 mg l ⁻¹ for 3 h)	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> (5 fish treatment ⁻¹)	Not effective (6.2 mg l ⁻¹ for 12 h, 25 mg l ⁻¹ for 3 h) to 10% effective (25 mg l ⁻¹ for 12 h) ^a	Tojo et al. (1992)
Triclabendazol	40 g kg ⁻¹ feed	10 d & 5 d oral treatment. 15°C. Lab study	<i>Gyrodactylus</i> sp.	<i>Oncorhynchus mykiss</i> (20 fish treatment ⁻¹)	Time dependent; parasite reduction (5 d) to 100% effective (10 d) ^f	Tojo & Santamarina (1998)
Virkon S + heat	Low concentration	25, 30, 35 & 40°C. Lab study	<i>G. salaris</i>	<i>Oncorhynchus mykiss</i> fin clips	Temperature dependent; parasite survival ranges from 119.7 min (25°C) to 9 s (40°C) ^g	Anttila et al. (2007)

Appendix 1 (continued)

Compound	Dose	Application	<i>Gyrodactylus</i> spp.	Host	Efficacy	Source
(3) Metal salts						
Aluminium	25, 50, 100 & 200 µg l ⁻¹	Continuous exposure (11–15 d) until fish died/recovered. 8°C. Lab study	<i>G. salaris</i>	<i>Salmo salar</i> (>50 parasites host ⁻¹)	Not effective (25 µg l ⁻¹) to 100% effective after 3 d (200 µg l ⁻¹) ^{b,c}	Poléo et al. (2004)
Aluminium (& acidification)	92–250 µg l ⁻¹ at pH 5.0, 5.2, 5.6 & 5.9	Continuous exposure (4–30 d) 12°C. Lab study	<i>G. salaris</i>	<i>Salmo salar</i> (15 fish treatment ⁻¹)	No effect of pH, but dose-dependent effect. At pH 5.0 effect of Al was enhanced: 100% effective after 9 d ^{b,c}	Soleng et al. (1999)
	Al-rich & Al-poor water (pH 5.8) & control (pH 6.3)	Continuous exposure. 11–13°C. Lab study	<i>G. derjavini</i> (<i>G. derjavinoïdes</i>)	<i>Salmo trutta</i> (10 fish treatment ⁻¹)	Al-rich water: 100% effective after 3 d. Al-poor water: decrease in parasites ^g	Pettersen et al. (2006)
			<i>G. macronychus</i>	<i>Phonixus phonixus</i> (15 fish treatment ⁻¹)	Al-rich water: 100% effective after 8 d. Al-poor water: not effective. ^g	
Cadmium	5 µg l ⁻¹	20 & 30 d. 23°C. Lab study	<i>G. bullatarudis</i>	<i>Poecilia reticulata</i>	Not effective (after 5 & 15 d in 20 d trials) to decrease in parasite burden after 30 d ^d	Carter (2003)
Copper	5 µg l ⁻¹ 10, 20, 40 & 80 µg l ⁻¹	60 d. 23°C. Lab study Continuous exposure until fish died/recovered. 8°C. Lab study	<i>G. turbulli</i> <i>G. salaris</i>	<i>Poecilia reticulata</i> <i>Salmo salar</i> (>50 parasites host ⁻¹)	Not effective ^d	Carter (2003) Poléo et al. (2004)
Iron	25, 50, 100 & 200 µg l ⁻¹	Continuous exposure until fish died/recovered. 8°C. Lab study	<i>G. salaris</i>	<i>Salmo salar</i> (>50 parasites host ⁻¹)	Not effective ^c	Poléo et al. (2004)
Manganese	100, 200, 400 & 800 µg l ⁻¹	Continuous exposure (11–15 d) until fish died/recovered. 8°C. Lab study	<i>G. salaris</i>	<i>Salmo salar</i> (>50 parasites host ⁻¹)	Not effective	Poléo et al. (2004)
Zinc	50, 100, 200 & 400 µg l ⁻¹ 8.3, 23.7, 38.1, 68.5, 129.5 & 260.3 µg l ⁻¹	Continuous exposure (11–15 d) until fish died/recovered. 8°C. Lab study 1 wk pre-exposure to required conc.; after infection continuous exposure until isolated fish died/recovered (up to 29 d) 25°C. Lab study	<i>G. salaris</i> <i>G. turbulli</i>	<i>Salmo salar</i> <i>Poecilia reticulata</i>	Not effective (50 µg l ⁻¹) to 100% effective after 3 d (400 µg l ⁻¹) ^c Decrease in peak parasite burden, but higher infection establishment (240 µg l ⁻¹). Combined effects of parasite infection & Zn exposure more detrimental to host than parasite ^c	Poléo et al. (2004) Gheorgiu et al. (2005)
4. Salt						
Salinity	5, 7.5, 10, 15, 20 & 33‰ salinity 33‰ salinity 5–20‰ NaCl	Continuous exposure. 1.4, 6 & 12°C. Lab study 5, 15, 30 & 60 min. 12°C. Lab study 4 d. 11°C. Lab study	<i>G. salaris</i> <i>G. salaris</i> <i>G. derjavini</i> (= <i>G. derjavinoïdes</i>)	<i>Salmo salar</i> (fins of 6 fish for 33‰ salinity; for all others, 12 fish treatment ⁻¹) <i>Salmo salar</i> (15 fish treatment ⁻¹) <i>Oncorhynchus mykiss</i> (21–23 fish treatment ⁻¹)	Not effective (5.0‰) to 100% effective after a few minutes (33‰) ^{b,d} Time dependent effect: no effect (5 min) to 100% effective (60 min) ^{b,c} Not effective (5–7‰) to 100% effective (10–20‰) ^{b,c}	Soleng & Bakke (1997) Soleng et al. (1998) Buchmann (1997)

^aIn vitro tests included in study; ^bInfection intensity was determined by sub-sampling methods other than screening the entire fish; ^cCauses mortality in some or all hosts; ^dSigns of host toxicity; ^eInfection intensity was determined 24 h post-treatment by ^fpelvic fin clips, or ^gmucus scrapings; ^hMortality not indicated by author; ⁱNo primary literature available