NOTE

Histological evaluation of sodium percarbonate exposure on the gills of rainbow trout

James M. Forwood^{1,*}, James O. Harris¹, Matt Landos², Marty R. Deveney^{3,4}

¹School of Biological Sciences, Flinders University, GPO Box 2100, Adelaide, SA 5001, Australia

²Future Fisheries Veterinary Services Pty Ltd, East Ballina, NSW 2478, Australia

³South Australian Research and Development Institute (SARDI), and ⁴Marine Innovation South Australia (MISA), West Beach, SA 5024, Australia

ABSTRACT: *Ichthyophthirius multifiliis* is a recurring problem in Australian rainbow trout *Oncorhynchus mykiss* farms and requires strategically timed, repeat treatments for effective management. Sodium percarbonate (SPC) is permitted for use in Australia, with host safety margins based on the toxicity of acute exposures to hydrogen peroxide (HP), the active product released when SPC is added to water. The effects of exposure to HP released by SPC, of repeated doses and of doses exceeding 100 mg l⁻¹ on rainbow trout are unknown. We exposed juvenile rainbow trout (mean weight: 30.5 ± 9 g) to repeated doses of 50, 150 and 250 mg l⁻¹ SPC for 1 h on Days 1, 2, 7 and 8 of a treatment regime. The effect of SPC was assessed by histological evaluation of structural changes in gill tissue. Survival was 100 % in all groups, but some fish exposed to 250 mg l⁻¹ SPC displayed impaired swimming performance, and on Day 9 after the final treatment, oedema was present in 9.8 % of lamella, which was significantly higher than the mean occurrence of 1.7, 4.2 and 1.3 % in fish treated with 0, 50 and 150 mg l⁻¹ SPC, respectively. These changes resolved within 24 h of the cessation of treatment. We conclude that SPC is safe to use on rainbow trout in doses of ≤150 mg l⁻¹ at 17°C, however caution is advised at doses approaching 250 mg l⁻¹. Water temperature, fish age, fish size and maturity, intensity of parasite infection and stocking density could alter the sensitivity of rainbow trout to SPC treatments.

KEY WORDS: *Oncorhynchus mykiss* · Sodium percarbonate · Gills · Histology

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Sodium percarbonate (SPC) is a granular, water-soluble compound that dissociates in water into sodium carbonate and hydrogen peroxide (HP), the latter of which is a powerful oxidant (Bostek 1989) that has been used extensively in aquaculture as a therapeutic compound for control of ectoparasitic infections of fish (Noga 2000). HP decomposes into water and carbon dioxide and does not form cumulative or toxic metabolites (Pedersen et al. 2006); it therefore has low environmental impact, and treated fish are food-safe. However, treating fish using liquid HP can be problematic, since obtaining the desired

dose is difficult due to the effects of water biogeochemistry (Bishop et al. 1968). There are also workplace health and safety concerns with handling HP (Eul et al. 2001), and security restrictions on transport and storage of the product (Department of Homeland Security 2007). Conversely, SPC is a granular solid with lower workplace health and safety risk and no current security sensitivity; its gradual dissociation in water to HP also provides a wider fish safety margin (Buchmann et al. 2003).

SPC is effective against free-living stages of *Ichthyophthirius multifiliis* (Ichthyophthiriidae) *in vitro* (Buchmann et al. 2003, Heinecke & Buchmann 2009, Forwood et al. 2014a). The *I. multifiliis* trophont, which

*Corresponding author: james.forwood@flinders.edu.au

resides in the host's epidermis, is unaffected by SPC treatment, therefore effective management of *I. multifiliis* requires repeat treatments specifically timed to reduce or eliminate reinfection and interrupt the parasite's life cycle (Picón-Camacho et al. 2012). SPC can be used in Australia pursuant to an Australian Pesticides and Veterinary Medicines Authority (APVMA) minor use permit (MUP) (PER12944) to treat ectoparasites of finfish at doses of up to 100 mg l⁻¹, based on Buchmann et al. (2003) and an analysis of teleost HP toxicity data (authors' unpubl. data) repeated over 3 consecutive days. However, results from *in vitro* trials indicated that doses above 100 mg l⁻¹ are required to effectively control all free-living stages of *I. multifiliis* (see Forwood et al. 2014a). The impacts of HP on Atlantic salmon *Salmo salar* (Salmonidae) (see Johnson et al. 1993, Kiemer & Black 1997) and rainbow trout *Oncorhynchus mykiss* (Salmonidae) (see Speare et al. 1999, Tort et al. 2002) have been investigated, and results indicated that structural damage was restricted to the gills (Roth et al. 1993, Kiemer & Black 1997, Tort et al. 2002). Alkalinity and pH could affect the efficacy and host toxicity of HP treatment (Tort et al. 2002), although the tolerance of rainbow trout to HP slowly released from SPC with significantly increased alkalinity from carbonate is unknown.

In this study, we investigated the histological effect on the gills of rainbow trout in response to baths of SPC applied at a frequency designed to interrupt the life cycle of *I. multifiliis*.

MATERIALS AND METHODS

Experimental design

A total of 192 *Oncorhynchus mykiss* with a mean weight of 30.5 ± 9 g were sourced from a commercial trout farm and transferred to Flinders University, South Australia. Fish were randomly selected, and 16 were transferred to each of 12 indoor 50 l recirculating aerated experimental tanks where they were acclimated for 6 d. Fish were fed 2% body weight daily of 3 mm Ridley Aqua-feed (Ridley AgriProducts®); no food was provided on treatment days. The recirculation system contained filtered dechlorinated municipal water; water hardness, pH (YSI 9300 photometer, YSI), temperature and dissolved oxygen (Handy Polaris H01P, OxyGuard®) were measured daily. Water hardness was 145 mg l⁻¹, alkalinity 20 mg l⁻¹ as CaCO₃, pH 6.1, temperature was maintained at 17 ± 1°C and dissolved oxygen was 7.8 ± 0.81 mg l⁻¹.

Dosing method and duration

At the end of the acclimation period, each tank was randomly assigned to 1 of 4 treatments (3 tanks treatment⁻¹): no dose (CON; control), 50, 150 and 250 mg l⁻¹ SPC (nominal concentration). Fish were exposed to SPC at the designated dose for 1 h, 4 times on Days 1, 2, 8 and 9, followed by a 10 d recovery period. On treatment days, the designated dose of granular SPC (Sigma-Aldrich®, lot no. MKBB5394V) was mixed in a 10 l bucket of water taken from the treatment tank, left to dissolve for 30 min, stirred, and then added to a treatment tank and mixed further. Fish were transferred from the 12 experimental recirculation tanks into 12 static treatment tanks for 1 h and then returned to the experimental tanks. The concentration of HP released was measured photometrically (YSI 9300 photometer) immediately prior to the fish being added to the treatment tank, and immediately after the fish were removed. Water in the experimental tanks was exchanged on the second and fourth treatment days.

Fish sampling

Rainbow trout were observed throughout the treatment period for signs of toxicity such as impaired swimming performance, colour change or loss of equilibrium. Any mortality was recorded. Fish were sampled immediately after treatment on Days 2 and 9, and additional sampling was carried out on Days 3, 7, 10, 14 and 18. Randomisation was ensured by capturing all the fish using a hand net, removing them one at a time and randomly assigning 2 fish numbers to be sampled; 6 fish for each treatment. Sampled fish were replaced with fin-clipped fish of a similar size during treatment periods to maintain the same organic load. The sampled fish were immediately euthanised with an overdose (40 ml 1000 l⁻¹ bath) of Aqui-S® (Aqui-S NZ). Histology samples were taken from the right first gill arch and immediately fixed in 10% neutral buffered formalin (NBF) until histological processing.

Histology processing and evaluation

Fixed tissues were dehydrated in a graded series of ethanol, cleared in xylene and embedded in paraffin wax. After processing, tissues were sectioned at 5 µm and stained with haematoxylin and eosin. Three well-oriented representative filaments from

the first gill arch of each fish with no *Ichthyophthirius multifiliis* infection were assessed. Slides were coded and examined blind. Structural changes in gill tissues (see Fig. A1 in the Appendix) were assessed and quantified by examination using a compound microscope (Olympus CH-2, Olympus Australia). To quantify the status of the gills, morphometric indices based on a method developed for branchial tissue by Speare & Ferguson (1989) and modified by Sanchez et al. (1997) were used. Each filament was evaluated for (1) total number of lamellae on the filament suitable for assessment; (2) lamellar oedema: the percentage of lamellae with visible separation of the 2 layers of epithelial cells, involving at least 50 % of the given lamella; (3) epithelial hyperplasia: the percentage of lamellae with one or more points of thickened areas of epithelial tissue (>3 cell layers thick), excluding those lamellae where this occurred exclusively at the tip (see index 7); (4) lamellar fusion: the percentage of lamellae that had greater than one-third fused to the adjacent lamella; (5) lamellar inflammation: the percentage of lamellae that had one or more clusters (>5 cells) of inflammatory cells located in the regions between the pillar cells and the outer layer of the epithelium; (6) interlamellar inflammation: the percentage of zones between 2 adjacent lamellae that had clusters (>5 cells) of inflammatory cells; and (7) clubbed lamellae: the percentage of lamellae with an accumulation of epithelial cells near the tip of the lamella that increases the number of cell layers at the lamellar tip, appearing as a 'club-shape'.

The presence of *I. multifiliis* in the histology sections was recorded and the prevalence and intensity of infection (Bush et al. 1997) was generated for each hemibranch.

Statistical analysis

The values for each index from a given fish were expressed as the pooled mean from the 3 filaments evaluated. Normality of the data was tested using the Kolmogorov-Smirnov test and variances were tested using Levene's test. Indices that failed to meet these assumptions were log-transformed prior to analysis. Differences in indices between treatment and control groups were analysed using 3-way nested ANOVAs with sample day and dose as orthogonal factors, and treatment tank nested within dose. Where significant differences were detected in these ANOVAs, an unpaired *t*-test was used as the post-test for specific comparisons. Statistical analyses were performed using SPSS Statistics v.20.0 (IBM), and significance

for all tests was judged at $p < 0.05$. Results are presented as mean $\pm$ SD (range).

RESULTS

The mean measured concentration of HP was 18 ± 2.5 (10 to 22) mg l^{-1} released from 50 mg l^{-1} SPC, 49 ± 6.4 (36 to 60) mg l^{-1} released from 150 mg l^{-1} SPC and 72 ± 7.7 (52 to 82) mg l^{-1} released from 250 mg l^{-1} SPC.

Survival was 100 % in all groups. Prevalence of *Ichthyophthirius multifiliis* was 59 % (95 % CI = 50 to 67 %) and mean intensity was 3.7 ± 1.6 (1 to 9). Gross observations of rainbow trout exposed to 50 and 150 mg l^{-1} SPC showed no adverse reactions to treatment, but reduced movement and slowed escape response was evident in fish exposed to 250 mg l^{-1} SPC. Two fish in one replicate exposed to 250 mg l^{-1} lost equilibrium during the second treatment and were returned to the recovery tank 5 min before the end of the treatment period; these fish recovered with no apparent negative effects and were returned to the randomization sampling.

Occurrence of lamellar oedema showed a time-dependent response to dose (3-way ANOVA: $F_{18,62} = 7.603$, $p < 0.001$), and there was variation between replicate tanks within doses (3-way ANOVA: $F_{47,62} = 7.640$, $p < 0.001$). In fish treated with 250 mg l^{-1} SPC, a mean of 9.8 ± 3.6 % (2.4 to 21.5 %) of lamella had oedemas on Day 9 after the final treatment, which was significantly higher than the mean of 1.7 ± 0.7 % (0 to 0.9 %), 4.2 ± 2.8 % (0.6 to 15.3 %), and 1.3 ± 0.4 % (0.8 to 2.4 %) in fish treated with 0, 50 and 150 mg l^{-1} SPC, respectively (Fig. 1). Epithelial hyperplasia and lamellar fusion changed significantly between sample days (3-way ANOVAs: $F_{6,62} = 3.933$, $p < 0.002$ and $F_{6,62} = 2.697$, $p < 0.022$, respectively) but was unaffected by dose; lamellar inflammation varied between sample days and replicate tanks in the same treatment doses (3-way ANOVA: $F_{47,62} = 1.775$, $p < 0.017$). There was no significant difference in the occurrence of clubbed lamellae or interlamellar inflammation between doses or sample days.

DISCUSSION

We observed no mortality in fish exposed to up to 250 mg l^{-1} SPC; doses of up to 150 mg l^{-1} SPC appear to be safe for use on rainbow trout at 17°C. The SPC used in this study contained 30 to 40 % HP, releasing up to 82 mg l^{-1} HP from a 250 mg l^{-1} SPC treatment.

Fig. 1. Percentage change in lamellar oedema in rainbow trout *Oncorhynchus mykiss* exposed to sodium percarbonate (SPC) on Days 1, 2, 8 and 9, at 0 (CON; control), 50, 150 and 250 mg l⁻¹ for 1 h. Fish were sampled immediately after treatment on Days 2, 3, 7, 9, 10, 14 and 18. Superscript letters indicate significant differences between sample days; numeric superscripts indicate significant difference between doses on the fourth sampling day (Day 9) ($p < 0.05$). Error bars: SE

This HP concentration is below the reported no observable effect concentration (NOEC) and lethal concentration (LC₅₀) values for juvenile rainbow trout. Gaikowski et al. (1999) reported that the NOEC for HP in rainbow trout in 60 min exposures administered on Days 1, 3 and 5 at 12°C was 188 mg l⁻¹ in fry and 162 mg l⁻¹ in fingerlings, which is within the range of NOEC values for juvenile rainbow trout reported by Rach et al. (1997) at 12°C. Tort et al. (2002) reported that the LC₅₀ of HP to 30 to 60 d old rainbow trout during a single 60 min exposure was approximately 180 mg l⁻¹ at 14°C.

HP oxidises more rapidly with increasing water temperature (Rach et al. 1997) and with decreasing pH (Bishop et al. 1968), and is therefore more toxic at higher temperatures and lower pH. The dissociation of SPC also releases sodium carbonate, which increases alkalinity and increases and buffers pH (Buchmann et al. 2003), stabilising the oxidative rate of HP. Increasing the dose of SPC to provide a dose of HP comparable with the NOEC for HP would provide a better understanding of the effects of increased alkalinity on the toxicity of HP in rainbow trout. Alkalinity does not have a significant effect on the efficacy of SPC against the free-living stages of *Ichthyophthirius multifiliis* (see Forwood et al. 2014a), and this change may also have a limited effect on the host.

Rainbow trout exposed to 250 mg l⁻¹ SPC for 1 h showed impaired swimming (including slowed reaction time and diminished flight response), and 2 fish lost equilibrium and were removed from the treatment tank. Fish exposed to 250 mg l⁻¹ SPC also had significantly more gill oedema after the fourth treatment than other treatment groups and controls. Rainbow trout exposed to ≥ 100 mg l⁻¹ HP for 1 h displayed epithelial lifting and gill necrosis (Tort et al. 2002) which can impair basic gill function including oxygen transport in high exposures (500 mg l⁻¹) (Powell & Perry 1997). Derksen et al. (1999) attributed epithelial cell swelling in the gills of rainbow trout exposed to HP to the oxidant effect of HP overcoming innate antioxidant defences. Oxidising agents denature the protein components of the outer cell membrane, impairing their permeability and leading to hydropic degradation (Derksen et al. 1999). Thus, the increase in oedema we observed at 250 mg l⁻¹ SPC is likely due to oxidation of surface membranes. We also observed a significant difference in epithelial hyperplasia and lamellar fusion between sample days, but it was unrelated to dose. Healthy fish can display idiopathic epithelial hyperplasia (Speare & Ferguson 1989), and it is likely that the effect we detected was unrelated to the treatment procedure. Our results showed that SPC treatments at 250 mg l⁻¹ at 17°C may cause gill oedema and loss of equilibrium in treated fish, but that treatments of up to 150 mg l⁻¹ are likely safe. Further research investigating the physiological effects of SPC at high doses is required to determine the effect on ionic transport and dysfunction of blood gases (Powell & Perry 1997). The stocking densities in this study were low, whereas in commercial aquaculture operations stocking densities are likely to be higher, which can increase stress levels (Conte 2004), leaving fish more sensitive to the negative effects of chemical treatments. Higher doses should therefore be used with caution.

The fish used in this study were infected with *I. multifiliis*. Rainbow trout can mount an immune response to *I. multifiliis* (see Buchmann et al. 2001), but the inflammatory response associated with this parasite is highly variable (Ventura & Paperna 1985). Changes associated with individual *I. multifiliis* trophonts were excluded from the analysis, and prevalence and intensity were low. Trophonts were randomly located throughout the lamellae. The significant occurrence of generalised lamellar inflammation may be due to *I. multifiliis* infection, but it is unlikely to have been linked to HP exposure.

SPC is effective against all free-living stages of *I. multifiliis* at 64 mg l⁻¹ at 17°C, but 250 mg l⁻¹ is re-

quired to be effective against the tomont at 12°C (Forwood et al. 2014a). The upper recommended dose of SPC on the MUP in Australia could be safely increased to 150 mg l⁻¹ for juvenile rainbow trout in water temperatures ≤17°C, improving treatment efficacy for *I. multifiliis*. Fish age, intensity of parasite infection and stocking density should, however, be taken into consideration when making decisions about doses of SPC or HP to manage parasites. Dose (and therefore efficacy) is also influenced by a range of system dynamics (Rach & Ramsay 2000, Forwood et al. 2014b) and HP concentrations should therefore be monitored throughout the treatment to maintain the target dose (Forwood et al. 2014b).

HP toxicity varies with fish size (Rach et al. 1997) and among species (Gaikowski et al. 1999). Tolerance of doses higher than those currently recommended needs to be assessed for other species.

Acknowledgements. We are grateful to the staff at Snobs Creek Hatchery in Snobs Creek, Victoria for providing and transporting rainbow trout, to the Victorian Department of Environment and Primary Industries (DEPI) for funding and support of the project, and to the Victorian Trout Growers Association (VTGA). This work was supported by funds provided by the project tactical research fund: "Optimisation of treatment of *Ichthyophthirius multifiliis* in farmed trout", and was supported by funding from the Fisheries Research and Development Corporation on behalf of the Australian Government (Project 2011/255). All procedures approved by the Flinders University Animal Welfare Committee (Permit E376).

LITERATURE CITED

- Bishop DE, Stern G, Fleischman M, Marshall S (1968) Hydrogen peroxide catalytic oxidation of refractory organics in municipal water. *Ind Eng Chem Process Des Dev* 7: 110–117
- Bostek CC (1989) Oxygen toxicity: an introduction. *AANA J* 57:231–237
- Buchmann K, Sigh J, Nielsen CV, Dalgaard M (2001) Host responses against the fish parasitizing ciliate *Ichthyophthirius multifiliis*. *Vet Parasitol* 100:105–116
- Buchmann K, Jensen PB, Kruse KD (2003) Effects of sodium percarbonate and garlic extract on *Ichthyophthirius multifiliis* theronts and tomocysts: *in vitro* experiments. *N Am J Aquaculture* 65:21–24
- Bush AO, Lafferty KD, Lotz JM, Shostak AW (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. *J Parasitol* 83:575–583
- Conte FS (2004) Stress and the welfare of cultured fish. *Appl Anim Behav Sci* 86:205–223
- Department of Homeland Security (2007) Appendix to chemical facility anti-terrorism standards. *Fed Reg* 72: 65420–65435
- Derksen JA, Ostland VE, Ferguson HW (1999) Effects of hydrogen peroxide on clearance of formalin-killed *Flavobacterium branchiophilum* from the gills of rainbow trout, *Oncorhynchus mykiss* (Walbaum). *J Fish Dis* 22: 59–67
- Eul W, Moeller A, Steiner N (2001) Hydrogen peroxide. In: Kirk-Othmer encyclopedia of chemical technology. Wiley Online Library, <http://onlinelibrary.wiley.com/10.1002/0471238961.0825041808051919.a01.pub2/abstract> (accessed 25 Jan 2013)
- Forwood JM, Harris JO, Landos M, Deveney MR (2014a) Minimum effective concentrations of formalin and sodium percarbonate on the free-living stages of an Australian isolate of *Ichthyophthirius multifiliis*. *Parasitol Res* 113:3251–3258
- Forwood J, Harris JO, Landos M, Deveney M (2014b) Evaluation of treatment methods using sodium percarbonate and formalin on Australian rainbow trout farms. *Aquacult Eng* 63:9–15
- Gaikowski MP, Rach JJ, Ramsay RT (1999) Acute toxicity of hydrogen peroxide treatments to selected lifestages of cold-, cool-, and warmwater fish. *Aquaculture* 178: 191–207
- Heinecke RD, Buchmann K (2009) Control of *Ichthyophthirius multifiliis* using a combination of water filtration and sodium percarbonate: dose-response studies. *Aquaculture* 288:32–35
- Johnson SC, Constible JM, Richard J (1993) Laboratory investigations on the efficacy of hydrogen peroxide against the salmon louse *Lepeophtheirus salmonis* and its toxicological and histopathological effects on Atlantic salmon *Salmo salar* and chinook salmon *Oncorhynchus tshawytscha*. *Dis Aquat Org* 17:197–204
- Jørgensen TR, Buchmann K (2007) Stress response in rainbow trout during infection with *Ichthyophthirius multifiliis* and formalin bath treatment. *Acta Ichthyol Piscat* 37: 25–28
- Kiemer MCB, Black KD (1997) The effects of hydrogen peroxide on the gill tissues of Atlantic salmon, *Salmo salar* L. *Aquaculture* 153:181–189
- Noga EJ (2000) Fish disease diagnosis and treatment. Iowa State University Press, Ames, IA
- Pedersen LF, Pedersen PB, Sortkjær O (2006) Dose-dependent decomposition rate constants of hydrogen peroxide in small-scale bio filters. *Aquacult Eng* 34:8–15
- Picón-Camacho SM, Marcos-Lopez M, Bron JE, Shinn AP (2012) An assessment of the use of drug and non-drug interventions in the treatment of *Ichthyophthirius multifiliis* Fouquet, 1876, a protozoan parasite of freshwater fish. *Parasitology* 139:149–190
- Powell MD, Perry SF (1997) Respiratory and acid-base pathophysiology of hydrogen peroxide in rainbow trout (*Oncorhynchus mykiss* Walbaum). *Aquat Toxicol* 37: 99–112
- Rach JJ, Ramsay RT (2000) Analytical verification of waterborne chemical treatment regimens in hatchery raceways. *N Am J Aquaculture* 62:60–66
- Rach JJ, Schreier TM, Howe GE, Redman SD (1997) Effects of species, life stage, and water temperature on the toxicity of hydrogen peroxide to fish. *Prog Fish-Cult* 59: 41–46
- Roth M, Richards RH, Sommerville C (1993) Current practices in the chemotherapeutic control of sea lice infestations in aquaculture: a review. *J Fish Dis* 16: 1–26
- Sanchez JG, Speare DJ, Johnson GJ (1997) Morphometric and histochemical assessment of the branchial tissue response of rainbow trout, *Oncorhynchus mykiss* (Wal-

- baum), associated with chloramine T treatment. J Fish Dis 20:375–381
- Speare DJ, Ferguson HW (1989) Fixation artifacts in rainbow trout (*Salmo gairdneri*) gills: a morphometric evaluation. Can J Fish Aquat Sci 46:780–785
 - Speare DJ, Carvajal V, Horney BS (1999) Growth suppression and branchitis in trout exposed to hydrogen peroxide. J Comp Pathol 120:391–402
 - Tort MJ, Jennings-Bashore C, Wilson D, Wooster GA, Bowser PR (2002) Assessing the effects of increasing hydrogen peroxide dosage on rainbow trout gills utilizing a digitized scoring methodology. J Aquat Anim Health 14:95–103
 - Ventura MT, Paperna I (1985) Histopathology of *Ichthyophthirius multifiliis* infections in fishes. J Fish Biol 27: 185–203E

Appendix.

Fig. A1. Light micrographs of histological sections of rainbow trout *Oncorhynchus mykiss* gills stained with haematoxylin and eosin. (A) Control fish gill with normal gills; (B) gill of fish exposed to 250 mg l⁻¹ sodium percarbonate (SPC) displaying lamellar oedema (arrow); (C) gill of fish exposed to 250 mg l⁻¹ SPC displaying epithelial hyperplasia (arrow); (D) *Ichthyophthirius multifiliis* trophont (arrow); (E) gill of fish exposed to 150 mg l⁻¹ SPC displaying epithelial hyperplasia (arrow); (F) gill of fish exposed to 250 mg l⁻¹ SPC displaying epithelial hyperplasia (arrow). Scale bars = 100 µm