

Seasonal variation in nitrification and denitrification in estuarine sediment colonized by benthic microalgae and bioturbating infauna

Søren Rysgaard^{1,*}, Peter Bondo Christensen¹, Lars Peter Nielsen²

¹National Environmental Research Institute, Vejlsvøvej 25, DK-8600 Silkeborg, Denmark

²Department of Microbial Ecology, Institute of Biological Science, University of Aarhus, Ny Munkegade, Bygning 540, DK-8000 Aarhus C, Denmark

ABSTRACT: Measurements of seasonal variation in oxygen fluxes, nutrient fluxes, and denitrification were obtained in an estuarine sediment inhabited by benthic microalgae and bioturbating infauna. Oxygen dynamics in the upper sediment strata were found to be controlled by the microalgae and there was a net flux of O₂ out of the sediment during spring and autumn. High assimilation by the microalgae reduced the efflux of NH₄⁺ and PO₄³⁻ from the sediment to the water column during daytime. Denitrification based on NO₃⁻ from the water column (D_w) only occurred in winter and spring, when NO₃⁻ was present in the water column, and activity was proportional to the water column NO₃⁻ concentration. The rate of D_w was reduced during daytime when the upper oxic zone of the sediment increased due to O₂ production by benthic microalgae. Coupled nitrification-denitrification (D_n) in the sediment was stimulated by the O₂ production during winter and spring, at which times NO₃⁻ and NH₄⁺ were present in the water column in high concentrations. In contrast, during summer, when the concentration of NO₃⁻ and NH₄⁺ in the water column was low, benthic microalgae inhibited D_n by competing with nitrifying bacteria for NH₄⁺. D_w accounted for 80% of the total denitrification during winter, while on an annual basis, D_w and D_n each accounted for 50% of the total denitrification activity. Benthic infauna, such as *Corophium* spp., *Hydrobia* spp., and *Nereis* spp., occurred in densities of up to several thousand ind. m⁻² from May to October. Oxygen consumption, D_w and D_n were linearly correlated with the density of the amphipod *Corophium* spp., all the processes studied being stimulated by the pumping of O₂- and NO₃⁻-rich water through the burrows in the upper 2 to 6 cm of the sediment. During summer, the D_n activity was, therefore, the net result of the inhibitory effect by benthic microalgae and the stimulatory effect of the benthic infauna. However, as the concentration of inorganic nitrogen in the overlying water and the sediment nitrification potential are both low in shallow coastal waters during summer, when benthic infauna density is high, we conclude that the stimulatory effect of bioturbating infauna on both D_w and D_n is of minor importance to the annual denitrification budget.

KEY WORDS: Denitrification · Nitrification · Microalgae · Bioturbation · ISN regulation

INTRODUCTION

Denitrification provides a sink in the global nitrogen budget and thereby plays an important part in controlling the degree of eutrophication in waters subjected to substantial anthropogenic input of nutrients. Denitrification in estuarine sediments thus decreases the

transport of nitrogen from land to the open sea (Seitzinger 1988). The process may be supported either by NO₃⁻ diffusing from the overlying water into the sediment or by NO₃⁻ being produced within the sediment by nitrification (Vanderborght & Billen 1975, Nishio et al. 1983, Jenkins & Kemp 1984). Diffusion of NO₃⁻ from overlying water is mainly controlled by a concentration gradient determined by the water NO₃⁻ concentration and the length of the diffusion path through the oxic zone (Christensen et al. 1990). Nitrifi-

*E-mail: fvsvr@dmu.dk

cation activity in sediments is mainly controlled by the availability of NH_4^+ and O_2 , as well as by population dynamics of nitrifying bacteria (Hansen et al. 1981, Henriksen et al. 1981).

Major seasonal and diurnal variation in nitrification and denitrification in shallow water sediments is explicable by changes in oxygen penetration depth caused by benthic microalgal growth and mineralization (Christensen et al. 1990, Risgaard-Petersen et al. 1994). Nitrification is preferentially stimulated in the daytime due to photosynthetic production of oxygen and deeper oxygen penetration, while the diffusion of nitrate from the water column is stimulated during night due to high oxygen demand for mineralization and low oxygen penetration. Benthic microalgae also assimilate nitrogen and may successfully compete with nitrifiers and denitrifiers for NH_4^+ and NO_3^- (Sundbäck & Granéli 1988, Nielsen et al. 1990, Nielsen & Sloth 1994). In addition, nitrifying bacteria in sediment with active phototrophs may become inhibited by high pH, high O_2 concentrations, CO_2 limitation, and toxic organic products (Henriksen & Kemp 1988).

Benthic infaunal activity also affects the physical and chemical processes within sediments, e.g. through burrow building, bioturbation and irrigation (Rhoads 1974, Kristensen 1984, Aller 1988). Recently, it has been shown that the amphipod *Corophium volutator* stimulates oxygen uptake, denitrification of water phase NO_3^- and coupled nitrification-denitrification by mass transport of water into its burrows (Pelegri et al. 1994).

The purpose of the present study was to measure and explain the diurnal and seasonal variation in O_2 consumption, nutrient fluxes, nitrification and denitrification in a shallow estuarine sediment colonized by benthic microalgae and bioturbating infauna.

MATERIALS AND METHODS

Study site. The study was carried out in Kertinge Nor, a small shallow estuary located on the east coast of the island of Funen, Denmark (Fig. 1). Kertinge Nor is connected to the sea through a narrow entrance to the east. The system receives only minor amounts of freshwater, mostly from small streams and precipitation, and the water residence time at the sampling locality is ca 4 to 6 wk (Christensen 1994). Salinity fluctuates on an annual basis but is generally around 17‰. During the study year, temperature varied from 2 to 4°C during winter up to 20 to 24°C during summer. The sampling site was located in shallow water (0.5 m depth) where the sediment was sandy. In early spring and late autumn,




Fig. 1. Study area and sampling site in the innermost part of Kertinge Nor estuary, Denmark

a dark-brown layer of benthic microalgae was observed on the sediment surface. Amphipods (*Corophium* spp.) dominated the benthic fauna from May to August at densities of up to 20 000 ind. m^{-2} . Later in summer, a mixture of polychaetes (*Nereis* spp.), oligochaetes, mud snails (*Hydrobia* spp.) and amphipods (*Corophium* spp.) were present in the sediment at a total density of up to 50 000 ind. m^{-2} .

Sampling. The sediment was sampled on 8 different dates during 1992. On each sampling date, 16 intact sediment cores were sampled by hand in 30 cm long and 52 mm wide Plexiglas tubes and brought to the laboratory within 4 h. In the laboratory the sediment cores were adjusted to give a sediment depth of ca 11 cm and a water column of ca 20 cm. The water column was stirred by a 2.5 cm teflon-coated magnet positioned 5 cm above the sediment surface the magnets receiving momentum from an external rotating magnet (60 rpm). The adjusted cores were left uncapped at the *in situ* temperature in a water bath containing 10 l of water from the locality. Five cores were incubated in the dark and five at the *in situ* light conditions. During winter, spring and autumn the sediment surfaces were illuminated with ca 70 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, and during summer ca 120 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$.

Flux measurements. Net fluxes of oxygen, inorganic nitrogen and inorganic phosphorus were measured by closing the cores with a rubber stopper and incubating for 1 to 12 h depending on the season. The incubation time was adjusted to ensure that the oxygen consumption in the cores never reduced the initial O_2 concentration by more than 20%. Water samples were collected just before closing the cores and after the incubation. Parallel incubated cores containing only water were used to correct for water column activity. Water samples were analyzed for O_2 by Winkler titration within a few hours of sampling and GF/C filtered samples for NO_3^- , NH_4^+ and PO_4^{3-} were immediately

frozen for later analysis. *In situ* fluxes of oxygen and inorganic nitrogen and phosphorus were calculated for each sediment core from the change in concentration during incubation and expressed as the rate per square meter. On several sampling dates, time series were made to ensure that the changes in oxygen and nutrient concentrations were linear over the incubation times used.

Denitrification measurements. Following measurement of the oxygen and nutrient fluxes, denitrification activity was determined on the same sediment cores by means of the isotope pairing technique (Nielsen 1992). $^{15}\text{NO}_3^-$ (20 to 60 μM) was added to the water column. However, in winter when high *in situ* NO_3^- concentrations were present, 250 μM $^{15}\text{NO}_3^-$ was added in order to obtain a uniform mixing of the isotopes. The added $^{15}\text{NO}_3^-$ was allowed to equilibrate with sediment porewater NO_3^- before the cores were closed with rubber stoppers. The cores were dark or light incubated for the same period as used for the flux measurements.

After incubation, samples of the water column and sediment porewater were collected for analysis of the ^{15}N labelling of N_2 , NH_4^+ and NO_3^- . The water column was sampled immediately upon removal of the stopper. 250 μl ZnCl_2 solution (50% w/w) was then added to the sediment surface to stop all bacterial activity and the sediment porewater and water column were carefully mixed with a Plexiglas rod. A sample of the resultant sediment slurry was taken by syringe. All samples for ^{15}N isotope analysis were preserved in gastight containers (Exetainers, Labco, High Wycombe, UK) with 2% (vol) of the ZnCl_2 solution. Finally, the sediment cores were sieved through a 1 mm sieve to recover the benthic infauna. Sediment porosity was measured at each sampling date in separate cores.

Test incubation. A test incubation was performed in order to find the optimal $^{15}\text{NO}_3^-$ concentration range for the denitrification measurements (Nielsen 1992). Four different concentrations of $^{15}\text{NO}_3^-$ in the overlying water (20, 40, 50 and 80 μM) were selected, and for each concentration, 4 intact sediment cores (22 cm^2 , 11 cm sediment and 20 cm water) were incubated as described above.

Analysis and calculations. The concentration of $\text{NO}_3^- + \text{NO}_2^-$ was determined on a flow injection analyzer (Tecator, Höganäs, Sweden) using the method described by Grasshoff et al. (1983). NH_4^+ concentration was determined manually using the method of Bower & Hansen (1980), and PO_4^{3-} was determined by a standard colorimetric method described by Grasshoff et al. (1983). The $^{15}\text{N}_2$ ($^{15}\text{N}^{15}\text{N}$ and $^{14}\text{N}^{15}\text{N}$) in the water and slurry samples was extracted into a helium headspace introduced in the Exetainers. After 5 min of vigorously shaking most of the N_2 is found in the head-

space, less than 2% of the N_2 gas being dissolved in the water at equilibrium. The gas in the headspace was then injected into a gas chromatograph coupled to a triple-collector isotopic ratio mass spectrometer (Robo-Prep-G⁺ in line with TracerMass, Europa Scientific, Crewe, UK) and the abundance and concentrations of $^{14}\text{N}^{15}\text{N}$ and $^{15}\text{N}^{15}\text{N}$ analyzed.

The production rate of the isotopes [$p(^{14}\text{N}^{15}\text{N}$ and $^{15}\text{N}^{15}\text{N})$] was calculated as follows:

$$p(^{14}\text{N}^{15}\text{N} \text{ and } ^{15}\text{N}^{15}\text{N}) = \frac{[V_1(C_{\text{water}} - C_{\text{ini}})] + [(C_{\text{slurry}} - C_{\text{ini}})V_2]}{At} \quad (1)$$

where C_{water} and C_{slurry} are the concentrations of the isotope in the water column and the sediment slurry, respectively, C_{ini} is the initial concentration of the isotope, V_1 is the volume of the sampled water, V_2 is the volume of porewater plus the remaining water column after the initial sampling, A is the area, and t is the incubation time.

Denitrification rates were estimated from the production of ^{15}N isotopes (Nielsen 1992):

$$D_{15} = p(^{14}\text{N}^{15}\text{N}) + 2p(^{15}\text{N}^{15}\text{N}) \quad (2)$$

$$D_{14} = \frac{p(^{14}\text{N}^{15}\text{N})}{2p(^{15}\text{N}^{15}\text{N})} D_{15} \quad (3)$$

where D_{15} and D_{14} are the rates of denitrification based on $^{15}\text{NO}_3^-$ and $^{14}\text{NO}_3^-$, respectively, and $p(^{14}\text{N}^{15}\text{N})$ and $p(^{15}\text{N}^{15}\text{N})$ are the rates of production of the 2 labelled N_2 species ($^{14}\text{N}^{15}\text{N}$ and $^{15}\text{N}^{15}\text{N}$, respectively). While D_{15} expresses denitrification activity of added $^{15}\text{NO}_3^-$, D_{14} expresses the total *in situ* denitrification activity.

The proportion of D_{14} that is based on NO_3^- from the water phase (D_w) was calculated from D_{15} and the $^{14}\text{N}:^{15}\text{N}$ ratio of water column NO_3^- :

$$D_w = D_{15} [^{14}\text{NO}_3^-]_w / [^{15}\text{NO}_3^-]_w \quad (4)$$

where $[^{14}\text{NO}_3^-]_w$ is the concentration of unlabelled NO_3^- and $[^{15}\text{NO}_3^-]_w$ the concentration of labelled NO_3^- in the water column. Finally, *in situ* denitrification of NO_3^- produced by nitrification (D_n) was calculated as:

$$D_n = D_{14} - D_w \quad (5)$$

To estimate D_w it was, as indicated above, necessary to measure the ^{15}N labelling of the water column NO_3^- . A pure culture of denitrifying *Pseudomonas nauticus* was used to convert $^{14}\text{NO}_3^-$ and $^{15}\text{NO}_3^-$ into N_2 gas composed of $^{28}\text{N}_2$, $^{29}\text{N}_2$ and $^{30}\text{N}_2$, which was subsequently analyzed by mass spectrometry (Risgaard-Petersen et al. 1993). Labelling of NO_3^- in the water column of the sediment cores was then calculated from the $^{29}\text{N}_2:^{30}\text{N}_2$ ratio in the analyzed gas.

RESULTS

Oxygen and nutrient fluxes

Marked seasonal variation was observed in the water column concentration and sediment-water flux of oxygen, inorganic nitrogen and phosphorus (Fig. 2). The benthic microalgae at the sediment surface produced O_2 throughout the year, as indicated by the net efflux or reduced net uptake of O_2 in the light-incubated sediment cores (Fig. 2A, B). Oxygen uptake in the dark-incubated cores, which represents the total sediment oxygen consumption, was significantly higher during the summer months of May until October (Fig. 2B). During summer, the water phase O_2 concentration exceeded 100% atmospheric saturation due to O_2 production by phytoplankton.

The NO_3^- concentration in the water column of Kertinge Nor displayed distinct seasonal fluctuation, the concentrations being high in winter and spring but negligible ($<0.2 \mu M$) throughout the summer. NO_3^- uptake by the sediment correlated to the water column NO_3^- concentration in both light- and dark-incubated sediment (Fig. 2C, D); as a result uptake was high in spring and winter but negligible during the summer period. The water column NH_4^+ concentration followed that of NO_3^- , although at a lower level. NH_4^+ efflux from the sediment was highest during the summer, with the rate being maximal in August (Fig. 2E, F). Throughout the growth season, NH_4^+ efflux was significantly lower in the light- than in the dark-incubated cores. The PO_4^{3-} flux changed during the year from sediment uptake in winter to efflux during summer. Moreover, PO_4^{3-} release from the sediment was higher in the dark-incubated sediment (Fig. 2H) than in the light-incubated sediment (Fig. 2G).




Fig. 2. Seasonal variation (1992) in O_2 , NO_3^- , NH_4^+ , and PO_4^{3-} fluxes in light (open bars) and dark (solid bars) incubated sediment shown together with the respective *in situ* concentration. Error bars on the flux measurements indicate SE ($n = 5$)

Optimization of the $^{15}\text{NO}_3^-$ concentration for the denitrification assay

Denitrification of $^{15}\text{NO}_3^-$ (D_{w15}) was proportional to the $^{15}\text{NO}_3^-$ concentration over the range 20 to 80 μM (Fig. 3A). Further, coupled nitrification-denitrification activity (D_n) was independent of the water column $^{15}\text{NO}_3^-$ concentration (Fig. 3A), thus verifying that incomplete isotope mixing due to heterogeneity was not a problem (Nielsen 1992, Pelegri et al. 1994, Rysgaard et al. 1994). On this basis we concluded that addition of 20 to 80 μM $^{15}\text{NO}_3^-$ would give real values of D_n . In February, however, when the *in situ* NO_3^- concentration was higher than 1000 μM , $^{15}\text{NO}_3^-$ was added to a final concentration of 250 μM .

The concentrations of both $^{14}\text{N}^{15}\text{N}$ and $^{15}\text{N}^{15}\text{N}$ increased linearly with incubation time (Fig. 3B). The intercept with the x-axis ($y = 0$) represents the time necessary for the NO_3^- profile to stabilize within the surface sediment following addition of the $^{15}\text{NO}_3^-$.

In situ denitrification activity

Seasonal variation in *in situ* denitrification is shown both as denitrification based on water phase NO_3^- (D_w ; Fig. 4A) and denitrification based on NO_3^- from nitrification (D_n ; Fig. 4B). In general, the rate of total denitrification ($D_w + D_n$) was highest in late winter and




Fig. 3. (A) Denitrification of $^{15}\text{NO}_3^-$ (D_{w15}) and coupled nitrification-denitrification (D_n) as a function of water column $^{15}\text{NO}_3^-$ concentration. Error bars indicate SE ($n = 5$). (B) Production of $^{29}\text{N}_2$ and $^{30}\text{N}_2$ as a function of time after the addition of $^{15}\text{NO}_3^-$. The experiment was undertaken in March at a temperature of 5°C




Fig. 4. Seasonal variation (1992) in (A) denitrification of NO_3^- from the overlying water (D_w) and (B) denitrification coupled to nitrification in the sediment (D_n). Open and solid bars represent the rates in light- and dark-incubated sediment, respectively. *In situ* NO_3^- concentration of the water column is given in (A). Error bars indicate SE ($n = 5$)

spring in both light- and dark-incubated sediment (Fig. 4). Denitrification based on water phase NO_3^- correlated with the NO_3^- concentration in the overlying water; it decreased from ca 60 $\mu\text{mol N m}^{-2} \text{h}^{-1}$ in February to <1 $\mu\text{mol N m}^{-2} \text{h}^{-1}$ in May and remained low throughout the summer period until the NO_3^- concentration increased in the autumn (Fig. 4A). Denitrification of water phase NO_3^- was always slightly higher in dark-incubated cores than in light-incubated cores.

Denitrification of NO_3^- produced by nitrification within the sediment was almost constant from February to June, but was lower and more variable during the rest of the year (Fig. 4B). Coupled nitrification-denitrification activity was significantly higher in light- than in dark-incubated cores during winter and spring, with the pattern being opposite in the summer period, when the concentration of inorganic nitrogen in the overlying water was very low. Coupled nitrification-denitrification accounted for ca 50% of total denitrification on an annual basis.

The presence of benthic infauna had a marked effect on oxygen consumption and denitrification rates, sediment oxygen consumption, D_w and D_n all being stimulated at increasing amphipod density (*Corophium* spp.) under both light and dark conditions (Fig. 5).

DISCUSSION

Use of the isotope pairing technique for measuring denitrification in estuarine sediments

Correct determination of actual *in situ* denitrification using the isotope pairing technique requires that 3 important assumptions are fulfilled (Nielsen 1992).




Fig. 5. Oxygen consumption denitrification based on NO_3^- from the water column (D_w) and coupled nitrification-denitrification (D_n) in light and dark as a function of the density of the amphipod *Corophium* spp. in May 1992. Linear regressions are shown. Each data point represents 1 sediment core

(1) Addition of $^{15}\text{NO}_3^-$ must not alter the rate of denitrification based on *in situ* NO_3^- . Microsensor and modelling studies (Christensen et al. 1989, 1990, Nielsen et al. 1990) have demonstrated a 1st order kinetic relationship between denitrification based on water phase NO_3^- and the NO_3^- concentration in the overlying water, denitrification of water phase NO_3^- primarily being determined by the NO_3^- concentration gradient within the upper oxic surface layer of the sediment. Since D_{w15} of the Kertinge Nor sediment was linearly correlated to the water phase $^{15}\text{NO}_3^-$ concentration (Fig. 3A), the first assumption was therefore fulfilled.

(2) The added $^{15}\text{NO}_3^-$ must mix uniformly with the NO_3^- already present in water column and in the sediment. Heterogeneous topography, bioturbation, inhomogenous nitrification activity, etc., may cause local variations in the transport of $^{14}\text{NO}_3^-$ and $^{15}\text{NO}_3^-$ to the anoxic denitrification zone, and hence underestimate *in situ* denitrification activity (D_{14}) since $^{14}\text{N}^{15}\text{N}$ production would then be less than that predicted on the assumptions of homogeneity (Broast et al. 1988). As demonstrated by several authors (Nielsen 1992, Pellegrini et al. 1994, Rysgaard et al. 1994) this possible underestimation can be analyzed by incubating the sediment cores at different $^{15}\text{NO}_3^-$ concentrations. At increasing $^{15}\text{NO}_3^-$ concentration, an increased denitrification of $^{14}\text{NO}_3^-$ will be detected directly as $^{14}\text{N}^{15}\text{N}$ on the mass spectrometer, thereby minimizing the possible underestimation of D_{14} . As demonstrated in the optimization experiment, coupled nitrification-denitrification was independent of the water phase NO_3^- concentration at concentrations greater than 20 μM , thus indicating uniform mixing of the added $^{15}\text{NO}_3^-$

(Fig. 3A). We always used a higher $^{15}\text{NO}_3^-$ concentration, thereby ensuring correct determination of both coupled denitrification (D_n) and total denitrification (D_{14}).

(3) A stable NO_3^- concentration gradient must be established in the surface layer of the sediment within a short time of $^{15}\text{NO}_3^-$ addition relative to the duration of incubation. If this is not the case, denitrification activity will be underestimated since the added $^{15}\text{NO}_3^-$ will not be immediately available to the denitrifying bacteria in the anoxic zone of the sediment. The time needed to establish a stable NO_3^- gradient depends on the O_2 penetration depth. During summer, when oxygen typically penetrates 1 mm down into shallow sediments, the 90% equilibration time is ca 5 min (Nielsen 1992). During winter, when the O_2 penetration is deeper, the establishment of a new NO_3^- gradient takes longer. Nevertheless, as the optimization experiment shows that production of ^{15}N -dinitrogen was linear after 30 min in March (Fig. 3B), the establishment of a stable NO_3^- profile took less than 30 min, this being a short period compared to the total incubation time of up to 12 h during winter. The third assumption was therefore also fulfilled at the Kertinge Nor sediment.

Effect of benthic microalgae and infauna on oxygen and nutrient dynamics

When benthic microalgae colonize the sediment surface of shallow waters, they may greatly influence oxygen and nutrient dynamics at the sediment-water interface. Oxygen production by benthic microalgae

may increase the oxygen penetration into the sediment by several mm (Revsbech & Jørgensen 1983) and thereby influence sediment metabolism as well as the turnover and flux of nutrients on both a diurnal and seasonal basis (Sundbäck 1986, Sundbäck & Granéli 1988, Risgaard-Petersen et al. 1994). Even though NH_4^+ was present at a high concentration in the sediment porewater, the benthic microalgae reduced the NH_4^+ flux to the water column significantly as a result of nitrogen assimilation as demonstrated by Rysgaard et al. (1993). Efflux of NH_4^+ was not measurable when photosynthesis was taking place, except in August (Fig. 2E). The microalgae therefore acted as an efficient filter, adsorbing the flux of ammonium from the deeper, anoxic sediment layers. Henriksen et al. (1980) and Sundbäck (1986) have also reported that a thin layer of benthic microalgae is able to control the flux of inorganic nitrogen from sediment to the overlying water. When NO_3^- was present in the water column in spring and early winter, the flux of NO_3^- was generally directed into the sediment, this being attributable to benthic assimilation in the surface layers and denitrification in the deeper sediment layers. Assimilation of inorganic nutrients by the benthic microalgae also influenced the PO_4^{3-} flux, a lower efflux being observed in light- than in dark-incubated sediment. The PO_4^{3-} flux into the sediment during winter was most likely due to binding of phosphate to oxidized iron, which is more abundant within the sediment during the cold season, when oxygen demand is lower and oxygen penetration into the sediment therefore deeper (Rasmussen & Jørgensen 1992).

Benthic oxygen production may also greatly influence sediment nitrification and denitrification. In the sediment from Kertinge Nor, we found that D_w was slightly less in the light than in the dark (Fig. 4A). However, it has recently been demonstrated that photosynthesis by benthic microalgae reduced denitrification based on water phase NO_3^- (D_w) by ca 50% during the day as a result of deeper O_2 penetration into the sediment when photosynthesis was taking place (Risgaard-Petersen et al. 1994). This deeper oxygen penetration enhances the diffusion path from the water column to the denitrifying zone, thereby reducing the NO_3^- supply for denitrification (Christensen et al. 1989, Nielsen et al. 1990). The difference in inhibition of denitrification in the 2 studies can be ascribed to the relatively higher activity (as judged from the much higher O_2 production) of benthic microalgae in the laboratory experiment by Risgaard-Petersen et al. (1994) as compared to our measurements of *in situ* activity in Kertinge Nor.

During winter and early spring when the availability of inorganic nitrogen was high, benthic photosynthesis stimulated coupled nitrification-denitrification (Fig. 4B),

this probably being due to oxygen stimulation of nitrification. This is in agreement with the study of Risgaard-Petersen et al. (1994), who found that benthic microalgae stimulated D_n during the day as a result of their O_2 excretion. However, both findings are in conflict with the hypothesis of Henriksen & Kemp (1988) that photosynthesis by benthic diatoms reduces nitrification due to a combination of a high competition for NH_4^+ , high pH and high O_2 concentration in the upper sediment layers, all of which inhibit the nitrification process, and thereby also the coupling between nitrification and denitrification. The competition between nitrifiers and benthic microalgae for inorganic nitrogen is particularly intense during periods of illumination; moreover, the concentration of available inorganic nitrogen in the overlying water and within the sediment is of major importance when evaluating the effect of benthic photosynthesis on nitrogen processes. Thus, during the summer period from May until November, when both NO_3^- and NH_4^+ was almost absent in the water column of Kertinge Nor, the photosynthetic activity of benthic microalgae actually reduced the activity of coupled nitrification-denitrification (Fig. 4B). Further, D_n reflected the NH_4^+ efflux from the sediment (Fig. 2F), thus indicating strong competition for NH_4^+ between nitrifiers and benthic microalgae in the surface layers of the sediment during the summer period, as has been suggested by Henriksen & Kemp (1988). Since benthic microalgae can assimilate NO_3^- and NH_4^+ at high rates for up to 60 h after sediment has been darkened, they therefore represent an efficient competitor for nitrifying bacteria at low nitrogen concentrations (Rysgaard et al. 1993). However, when NO_3^- or NH_4^+ are present at high concentrations in the overlying water column they can act as a nitrogen source for the benthic assimilatory demand, and thereby reduce the competition with nitrifiers for porewater NH_4^+ . This was the situation in Kertinge Nor during winter and spring as well as in the laboratory experiments of Risgaard-Petersen et al. (1994), where benthic primary production stimulated the coupled nitrification-denitrification. Thus, as indicated by the present study, there may be both diurnal and seasonal variation in the effect of benthic microalgae on the coupling between nitrification and denitrification in estuarine sediments, i.e. microalgal photosynthesis may stimulate denitrification during the cold season, when nitrogen availability is high, and inhibit denitrification during summer, when nitrogen availability is low.

Oxygen and nitrogen dynamics within the sediment may also be affected by bioturbating infauna (Henriksen et al. 1980, Aller 1982, Kristensen et al. 1991). In the sandy sediment of Kertinge Nor, sediment oxygen respiration and denitrification was significantly enhanced at increasing densities of the amphipod *Coro-*

phium spp. (Fig. 5). Stimulation of sediment oxygen consumption, D_n and D_w in the presence of *Corophium* spp. can be explained by mass transport of O_2 - and NO_3^- -rich water within the U-shaped amphipod burrows, which may penetrate as much as 2 to 6 cm into the sediment (Pelegri et al. 1994). Infauna density was particularly high from May until November (Fig. 6). During this period, the water column NO_3^- concentration was very low (Fig. 2) and the effect of bioturbating fauna on D_w was consequently negligible; the high infauna density primarily stimulated D_n activity (Fig. 5). The stimulation of coupled nitrification-denitrification activity was higher in light, probably because the infauna were more active during the day and therefore able to pump more O_2 -rich water into the sediment. Coupled nitrification-denitrification activity in May ranged from $8 \mu\text{mol N m}^{-2} \text{h}^{-1}$ in the dark to $13 \mu\text{mol N m}^{-2} \text{h}^{-1}$ in the light (Fig. 4B). From the data presented in Fig. 5, the activity in sediments devoid of *Corophium* spp. can be extrapolated to be less than $3 \mu\text{mol N m}^{-2} \text{h}^{-1}$. It is therefore obvious that the presence of *Corophium* spp. had a marked influence on coupled nitrification-denitrification activity in Kertinge Nor during the summer period. However, total nitrification activity is generally reduced in shallow coastal waters during summer since the population of nitrifiers in the sediment is reduced due to lower O_2 availability and higher NH_4^+ competition (Hansen et al. 1981). As a consequence, coupled nitrification-denitrification activity was low during summer in Kertinge Nor compared to that obtained during winter (Fig. 4B). The stimulatory effect of the bioturbating infauna (mainly present during summer) on D_w and D_n was therefore limited by the predominantly low water column NO_3^- concentration and the reduced nitrifying population in the sediment, respectively.

Stimulation of coupled nitrification-denitrification with increasing infauna density was 2-fold greater




Fig. 6. Seasonal variation (1992) in benthic infauna density at the study site. Error bars indicate SE ($n = 8$ to 16)




Fig. 7. Oxygen uptake and denitrification coupled to nitrification (D_n) as a function of amphipod density. The relative rates are rates measured in bioturbated sediment cores divided by rates measured in non-bioturbated sediment cores. The horizontal line at 1 thus represents activity in amphipod-free sediment cores. Data are from both light- and dark-incubated cores

than that of O_2 consumption rate (Fig. 7), a finding consistent with the observations of Pelegri et al. (1994). Nitrate produced by nitrification in a sediment devoid of bioturbating animals will diffuse both upwards to the water column and downwards to the anoxic denitrification zone; assuming homogeneous distribution of nitrifiers throughout the oxic surface layers of the sediment, approximately half of the NO_3^- will diffuse in each direction (Nielsen et al. 1990, Pelegri et al. 1994). Nitrate diffusing out of the oxic layer in an amphipod burrow can be denitrified further down the burrow, however, and coupling between nitrification and denitrification will therefore be much closer in bioturbated sediment. Thus, as a general rule, bioturbation should stimulate nitrification and sediment oxygen uptake to an equivalent extent, while coupled nitrification-denitrification should be stimulated twice as much as oxygen consumption when coupling of nitrification and denitrification is almost 100% (Pelegri et al. 1994).

Control and relative importance of D_w and D_n

Denitrification based on water phase NO_3^- (D_w) was highly correlated to the NO_3^- concentration in the water, activity being high during winter and spring and almost negligible throughout summer (Fig. 4A). A model relating D_w to O_2 uptake and the water column concentrations of O_2 and NO_3^- (Christensen et al. 1990) was tested on the present data set. The model is based on the fact that O_2 penetration within the sediment is a function of O_2 concentration and O_2 uptake. The O_2 penetration depth represents the diffusional path for NO_3^- to the underlying, anaerobic denitrification zone

and the concentration of NO_3^- in the water column divided by the diffusional path of NO_3^- thus represents the concentration gradient of NO_3^- . The flux of NO_3^- to the denitrification zone can therefore be calculated by multiplying this concentration gradient with the diffusion coefficient of NO_3^- in the sediment:

$$D_w = F_{\text{O}_2} \times \alpha \times \left[\left(1 + \frac{D_{\text{NO}_3^-}}{D_{\text{O}_2}} \times \frac{C_{\text{NO}_3^-}}{C_{\text{O}_2}} \times \frac{1}{\alpha} \right)^{\frac{1}{2}} - 1 \right] \quad (6)$$

where F_{O_2} is the sediment oxygen consumption, C_{O_2} and $C_{\text{NO}_3^-}$ are the respective concentrations in the water column, D_{O_2} and $D_{\text{NO}_3^-}$ are the respective coefficients of diffusion, and α is the ratio between the volume specific denitrification and oxygen respiration rates. The model is based on the fact that D_w depends on the water column NO_3^- concentration, the volume specific denitrification rate in the anoxic zone, and the length of the diffusion path through the oxic zone. The thickness of the oxic zone, in turn, is a function of O_2 uptake and O_2 concentration in the water column, assuming the same volume specific O_2 consumption rate throughout the oxic zone. The diffusion coefficients $D_{\text{NO}_3^-}$ and D_{O_2} need not to be measured since the ratio between them is invariably 0.8 in an aquatic medium. The volume specific denitrification and oxygen respiration rates are not determined either, but the ratio between them (α) is set to 0.8 on the basis of bacterial kinetics and microsensor studies (Christensen et al. 1989). As demonstrated in Fig. 8, the measured rates of D_w corresponded very well with the rates predicted by the model. Denitrification based on NO_3^- may therefore be estimated by this model using easily obtainable parameters such as O_2 consumption and water column O_2 and NO_3^- concentrations.

D_n activity is related to the nitrification rate and the efficiency with which the nitrification process is coupled to denitrification. In the sediment from Kertinge Nor,




Fig. 8. Measured (circles) and predicted (line) D_w (denitrification rate based on NO_3^- from the water column) during 1992. Error bars indicate SE ($n = 5$)

the highest rates of coupled nitrification-denitrification were observed during the cold months, when the temperature was below 4°C (Fig. 4B). Total microbial respiration in the sediment is reduced by the cold temperatures, thereby reducing total sediment O_2 consumption. Oxygen penetration within the sediment is therefore enhanced during winter (Rasmussen & Jørgensen 1992), which may activate nitrifying bacteria located deeper in the sediment, and thereby enhance total nitrification activity. A seasonal horizontal shift that ensures maximal nitrifying activity in the deeper sediment layers during winter will result in tight coupling between nitrification and denitrification in this part of the year, as demonstrated by Rysgaard et al. (1994) and Jensen et al. (1994).

Denitrification based on water phase NO_3^- and denitrification coupled to nitrification within the sediment both played an important role in nitrogen removal from Kertinge Nor estuary. At the shallow, sandy sampling site, D_w accounted for ca 50% of total annual denitrification. It has previously been stated that coupled nitrification-denitrification is the most important denitrification process in aquatic sediments (Seitzinger 1988). However, the present study indicates that D_w may be important in shallow coastal systems that receive significant nutrient input. Moreover, as the study also demonstrates, the relative importance of the 2 processes may change dramatically throughout the year: In February, when NO_3^- was present in high concentrations, D_w accounted for more than 80% of the total denitrification, while in the summer months, almost all denitrification was due to coupled nitrification-denitrification (Fig. 4).

We conclude that benthic microalgae may have a strong regulating effect on the efflux of nutrients from the sediment surface to the overlying water in shallow estuarine waters. Due to their assimilatory demand, they may efficiently reduce the flux of inorganic nutrients during the day and may also cause significant uptake of, for example, NO_3^- from the water column. From this and previous studies it is evident that the presence of benthic microalgae also has a marked influence on the diurnal and seasonal variation in sediment denitrification. The rather complex means whereby they regulate these processes can be summarized as follows. (1) Oxygen production by benthic microalgae increases the oxic zone within the sediment, thereby lengthening the diffusional path for NO_3^- from the water column to the anoxic denitrification zone and hence reducing denitrification activity based on water phase NO_3^- (D_w). (2) When inorganic nitrogen is present in excess, benthic oxygen production stimulates nitrification and the increased oxic surface layers caused by benthic primary production may stimulate coupled nitrification-denitrification (D_n).

(3) During summer, when the water column concentration of NO_3^- and NH_4^+ is low, there is strong competition for inorganic nitrogen between benthic microalgae, nitrifiers and denitrifiers. In this situation benthic primary production will reduce nitrification activity and thereby D_n , and nitrogen assimilation can decrease the activity of D_w as well.

A further conclusion to be drawn from this study is that bioturbation by benthic infauna can significantly stimulate sediment oxygen consumption, D_w and D_n within the sediment as long as excess inorganic nitrogen is present in the water column. However, since benthic infauna density is generally highest during summer, when the NO_3^- concentration is low, their effect on D_w will be of minor importance on an annual basis. Moreover, as the nitrification potential of sediment in shallow coastal waters is reduced during summer, the stimulatory effect on D_n of bioturbation by benthic infauna will also be of minor importance on an annual basis.

Acknowledgements. We thank Tage Dalsgaard for helpful discussions and Pia Nygård Christensen, Kitte Gerlich, Marlene Jessen and Lars Møller for valuable technical assistance. This research was supported by the Danish Agency of Environmental Protection through the Marine Research 90 programme.

LITERATURE CITED

- Aller RC (1982) The effects of macrobenthos on chemical properties of marine sediment and overlying water in animal-sediment relations. McCall PL, Tevesz MJS (eds). Plenum Press, New York, p 53–102
- Aller RC (1988) Benthic fauna and biogeochemical processes in marine sediments: the role of burrow structures. In: Blackburn TH, Sørensen J (eds) Nitrogen cycling in coastal marine environments. John Wiley and Sons Ltd, Chichester, p 301–338
- Bower E, Hansen HA (1980) A salicylate-hypochlorite method for determining ammonia in seawater. *Can J Fish Aquat Sci* 37:794–798
- Broast CW, Mulvaney RL, Baveye P (1988) Evaluation of nitrogen-15 tracer techniques for direct measurements of denitrification in soil: 1. Theory. *Soil Sci Soc Am J* 52: 1317–1322
- Christensen PB (ed) (1994) Nutrient transport and dynamics in Kertinge Nor/Kerteminde Fjord, Denmark. The Marine Research 90 Programme No. 43. The Danish Agency of Environmental Protection (in Danish with English summary)
- Christensen PB, Nielsen LP, Revsbech NP, Sørensen J (1989) Microsonation of denitrification activity in stream sediments as studied with a combined oxygen and nitrous oxide microsensor. *Appl Environ Microbiol* 55(5):1234–1241
- Christensen PB, Nielsen LP, Sørensen J, Revsbech NP (1990) Denitrification in nitrate-rich streams: diurnal and seasonal variation related to benthic oxygen metabolism. *Limnol Oceanogr* 35(3):640–651
- Grasshoff K, Erhardt M, Kremling K (1983) Methods of seawater analysis, 2nd edn. Verlag Chemie, Weinheim
- Hansen JI, Henriksen K, Blackburn TH (1981) Seasonal distribution of nitrifying bacteria and rates of nitrification in coastal marine sediments. *Microb Ecol* 7:297–304
- Henriksen K, Hansen J, Blackburn TH (1980) The influence of benthic infauna on exchange rates of inorganic nitrogen between sediment and water column. *Ophelia* (Suppl) 1: 249–256
- Henriksen K, Hansen JI, Blackburn TH (1981) Rates of nitrification, distribution of nitrifying bacteria, and nitrate fluxes in different types of sediment from Danish waters. *Mar Biol* 61:299–304
- Henriksen K, Kemp W (1988) Nitrification in estuarine and coastal marine sediments. In: Blackburn TH, Sørensen J (eds) Nitrogen cycling in coastal marine environments. John Wiley and Sons, p 201–249
- Jenkins MC, Kemp WM (1984) The coupling of nitrification and denitrification in two estuarine sediments. *Limnol Oceanogr* 29:609–619
- Jensen K, Sloth NP, Risgaard-Petersen N, Rysgaard S, Revsbech NP (1994) Estimation of nitrification and denitrification from microprofiles of sediment. *Appl Environ Microbiol* 60:2094–2100
- Kristensen E (1984) Effect of natural concentrations of nutrient exchange between a polychaete burrow in estuarine sediment and the overlying water. *J exp mar Biol Ecol* 75: 171–190
- Kristensen E, Jensen MH, Aller C (1991) Direct measurements of dissolved inorganic nitrogen exchange and denitrification in individual polychaete (*Nereis virens*) burrows. *J mar Res* 49:355–377
- Nielsen LP (1992) Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbiol Ecol* 86: 357–362
- Nielsen LP, Christensen PB, Revsbech NP, Sørensen J (1990) Denitrification and photosynthesis in stream sediment studied with microsensor and whole-core technique. *Limnol Oceanogr* 35(5):1135–1144
- Nielsen LP, Sloth NP (1994) Denitrification, nitrification and nitrogen assimilation in photosynthetic microbial mats. In: Stahl LJ, Caumette P (eds) Microbial mats, structure, development and environmental significance. NATO ASI Series G, Ecological Sciences, Vol 35. Springer Verlag, Berlin, p 319–324
- Nishio T, Koike I, Hattori A (1983) Estimation of denitrification and nitrification in coastal and estuarine sediments. *Appl Environ Microbiol* 45:444–450
- Pelegri SP, Nielsen LP, Blackburn TH (1994) Denitrification in estuarine sediment stimulated by the irrigation activity of the amphipod *Corophium volutator*. *Mar Ecol Prog Ser* 105:285–290
- Rasmussen H, Jørgensen BB (1992) Microelectrode study of seasonal oxygen uptake in a coastal sediment; role of molecular diffusion. *Mar Ecol Prog Ser* 81:289–303
- Revsbech NP, Jørgensen BB (1983) Photosynthesis of benthic microflora measured with high spatial resolution by the oxygen microprofile method: capabilities and limitations of the method. *Limnol Oceanogr* 28:749–756
- Rhoads DC (1974) Organism-sediment relations on the muddy sea floor. *Oceanogr mar Biol A Rev* 12:263–300
- Risgaard-Petersen N, Rysgaard S, Revsbech NP (1993) A sensitive assay for determination of $^{14}\text{N}/^{15}\text{N}$ isotope distribution in NO_3^- . *J Microbiol Meth* 17:155–164
- Risgaard-Petersen N, Rysgaard S, Nielsen LP, Revsbech NP (1994) Diurnal variation of denitrification in sediments colonized by benthic microphytes. *Limnol Oceanogr* 39: 573–579
- Rysgaard S, Risgaard-Petersen N, Nielsen LP, Revsbech NP (1993) Nitrification and denitrification in lake and estu-

- arine sediments measured by ^{15}N dilution technique and isotope pairing. *Appl Environ Biol* 59(7):2093–2098
- Rysgaard S, Risgaard-Petersen N, Sloth NP, Jensen K, Nielsen LP (1994) Oxygen regulation of nitrification and denitrification in freshwater sediments. *Limnol Oceanogr* 39:1643–1652
- Seitzinger SP (1988) Denitrification in freshwater and coastal marine ecosystems: ecological and geochemical significance. *Limnol Oceanogr* 33:702–724
- Sundbäck K (1986) What are the benthic microalgae doing on the bottom of Lahom Bay? *Ophelia (Suppl)* 4:273–286
- Sundbäck K, Granéli W (1988) Influence of microphytobenthos on the nutrient flux between sediment and water: a laboratory study. *Mar Ecol Prog Ser* 43:63–69
- Vanderborght JP, Billen G (1975) Vertical distribution of nitrate concentration in interstitial water of marine sediments with nitrification and denitrification. *Limnol Oceanogr* 20:953–961

This article was submitted to the editor

Manuscript first received: October 26, 1994

Revised version accepted: April 5, 1995