

# Effect of photoinhibition during bottle incubations on the measurement of seasonal primary production in a shallow coastal water\*

Joel C. Goldman and Mark R. Dennett

Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA

**ABSTRACT:** Primary productivity in Vineyard Sound, Massachusetts (USA), a shallow well-mixed coastal water, was determined at monthly intervals during May–October, 1981. Over the course of each study day fresh samples first were obtained at 3 h intervals and then incubated until sunset with time-course measurements of  $\text{H}^{14}\text{CO}_3$  uptake made at frequent intervals. Two seasonal patterns of  $^{14}\text{C}$  uptake were apparent. First, rapid initial uptake of  $^{14}\text{C}$  over each incubation during the spring and fall was followed by reduced uptake during periods of strong sunlight. The duration over which rapid uptake was sustained was a function of the prevailing light intensity at the start of the incubation. Under these conditions estimates of near surface primary productivity based on summing total production over the first 3 h of each back to back incubation were higher by 50 to 130 % than estimates based on single end-point measurements from full-day incubations. During summer, however, uptake of  $^{14}\text{C}$  was linear throughout each incubation when both total daily irradiation and temperature were maximal so that the 2 techniques for estimating productivity led to comparable results. The reasons for these seasonal differences in the pattern of  $^{14}\text{C}$  uptake are unclear, but may be related to changes in temperature or phytoplankton species, or both. Our results for a shallow well-mixed water are in general accord with the conclusion of Harris (1980) that when vertical mixing is sufficient to prevent prolonged exposure of phytoplankton to photoinhibiting light intensities at the surface, photoinhibition could be far more pronounced during a bottle incubation than is occurring in the water column.

## INTRODUCTION

Utility of the  $^{14}\text{C}$  bottle incubation technique for measuring marine primary productivity is based, to a large extent, on the premise that biological activity in a confined and small (hundreds of ml) sample of water under a light regime representative of the depth from where the sample was obtained and over incubation periods extending up to tens of hours will be comparable to what would have occurred had the sample been left undisturbed in the water column. In recent years, however, evidence has been accumulating that, for a variety of reasons, the longer a sample is confined the more pronounced may be the deviation between true water column primary productivity and the measured bottle rate (Harris, 1980; Goldman et al., 1981).

One of the major concerns with long incubations is

that photoinhibition, which commonly is observed in surface samples exposed to correspondingly high irradiation, may be far more pronounced in the incubation bottle than actually is taking place in the water column, thereby leading to underestimates of true primary productivity (Harris, 1980). The major problem is that photoinhibition is a time-dependent process (Kok, 1956; Takahashi et al., 1971) so that, depending on the prevailing patterns of vertical water movements, cells in the water column may have a significantly different temporal history of exposure to inhibiting irradiation than when confined in a bottle (Harris and Lott, 1973; Harris and Piccinin, 1977; Marra, 1978a, b). It is not surprising, therefore, that higher rates of photosynthesis have been observed on numerous occasions when cells were exposed to fluctuating rather than constant irradiation (Jewson and Wood, 1975; Jones, 1978; Marra, 1978a, b; Savidge, 1980; Gallegos et al., 1980; Gallegos and Platt, 1982), when short-term rates were compared with longer-term rates

\*Contribution No. 5437 from the Woods Hole Oceanographic Institution

under greater than saturating levels of irradiation (Ichimura and Saijo, 1958; Barnett and Hirota, 1967; Harris and Lott, 1973; Harris and Piccinin, 1977), or when the summed production from back to back incubations of short duration involving fresh samples taken over the course of a day were compared with production from a single incubation representing the same total time interval as the back to back incubations (Rodhe, 1958; Vollenweider and Nauwerck, 1961; Hammer et al., 1973; Gieskes et al., 1979).

In the present study we performed a series of back to back  $^{14}\text{C}$  incubations, each involving time-course measurements of primary productivity of fresh samples from a shallow location in Vineyard Sound, Massachusetts, midway between Martha's Vineyard and Woods Hole. We carried out these studies at monthly intervals over a 6 mo period to assess both the daily and seasonal interactions of sunlight and temperature upon the measurement of primary production.

## METHODS

### Primary production measurements

On each of the 6 study days between May 14, 1981 and October 29, 1981 we collected a series of samples at approximately 3 h intervals from pre-dawn until late afternoon (Table 1). Each sample was collected from 1 m depth with a 6 l Niskin bottle, filtered through 202  $\mu\text{m}$  Nitex mesh, and returned to the Woods Hole Oceanographic Institution (WHOI) where the incubations were performed outdoors. We incubated each sample for the balance of the day and made time-course measurements of  $\text{H}^{14}\text{CO}_3^-$  incorporation into organic matter. Our incubation system, consisting of ten 1 l vessels, was identical to the one used previously in other Vineyard Sound studies (Goldman et al., 1981;

Glibert et al., 1982). Each vessel was held at ambient temperature by circulating water from a temperature-controlled bath and was mixed continuously by magnetic bar stirring. Irradiation incident to each vessel was regulated at 60 % of full sunlight ( $I_0$ ) with neutral density screens.

Two additional samples were incubated for a full day starting on June 11, one always exposed to full sunlight and the other exposed to a variable light regime on two occasions (incident irradiation alternated in 30 min intervals between 100 %  $I_0$  and 60 %  $I_0$  on June 11 and between 60 %  $I_0$  and 30 %  $I_0$  on July 16) and 30 %  $I_0$  on the remaining dates. The last incubation of the day on September 29 was performed under full sunlight as well as at 60 %  $I_0$ .

Daily variations in solar irradiation (visible plus IR region) during the course of each study were measured at the study site with an International Light Radiometer (No. 700), whereas total daily irradiation (UV, visible, and IR) was recorded on an Eppley pyroheliometer located on the roof of the Clark Laboratory at WHOI, a short distance from where the experiments were performed.

We performed time-course measurements of  $^{14}\text{C}$  uptake by introducing  $\text{H}^{14}\text{CO}_3^-$  at the start of an incubation to yield a specific activity of 80 to 100  $\mu\text{Ci} \cdot \text{mMole DIC}^{-1}$ , followed by subsampling (10 ml) at about 15 min intervals for the first 30 min, then at 90 min intervals, and finally at 60 min intervals for the balance of the incubation. The samples were processed by pipeting the 10 ml into a scintillation vial containing enough 18 N  $\text{H}_2\text{SO}_4$  to lower the pH to 1.3. The vial contents were then sparged with pure nitrogen gas (200 ml  $\text{min}^{-1}$ ) for 15 min to remove  $^{14}\text{CO}_2$ , and 5 ml were added to another scintillation vial containing 9 ml of Aquasol. Radioactivity was measured by liquid scintillation counting on a Beckman LS-100C instru-

Table 1. Summary of environmental conditions and rates of primary productivity in Vineyard Sound, Massachusetts, based on full-day single end-point measurements at  $I_0$ , 60 %  $I_0$ , and 30 %  $I_0$  light levels

Date 1981	Water temperature (°C)	Incubation duration (h)	Total daily surface irradiation ( $\text{W h m}^{-2}$ )	Weather conditions (*)	Primary productivity ( $\mu\text{g C l}^{-1} \text{h}^{-1}$ )		
					$I_0$	60 % $I_0$	30 % $I_0$
May 14	10.5	14.5	6426	S→PC→S	N.M.**	1.4	N.M.
June 11	17.0	16.0	7716	S→IC	3.2	9.4	(9.1) <sup>+</sup>
Jul 16	22.0	15.0	8018	S	1.3 (1.7) <sup>++</sup>	3.8	N.M.
Aug 5	23.0	11.0	3091	OC→R	13.9	13.2	11.5
Sep 29	17.0	12.0	5031	OC→S	1.7	4.3	6.4
Oct 29	12.5	10.5	1057	OC	5.7	7.3	6.7

\* S – full sun, PC – partly cloudy, IC – intermittent clouds, OC – overcast, R – rain  
 \*\* Not measured  
 + Alternating 30 %  $I_0$  and 60 %  $I_0$  during 0.5 h intervals  
 ++ Alternating 60 %  $I_0$  and 100 %  $I_0$  during 0.5 h intervals

ment. The measured radioactivity represents the sum of labelled particulate and non-volatile dissolved organic carbon (Schindler and Holmgren, 1971).

On 3 occasions (June 11, August 5, September 29) additional filtered samples representing only labelled particulate carbon from the full day (60%  $I_0$ ) incubations were processed for  $^{14}\text{C}$  activity. 20 ml subsamples, taken intermittently during the course of the incubations, were filtered immediately through Whatman GF/F glass-fiber filters. The filters first were rinsed with 50 ml of filtered seawater and then placed in scintillation vials containing 10 ml of Aquasol. Radioactivity was measured, as described above.

### Analytical methods

Particulate carbon and particulate nitrogen were measured with a Perkin Elmer 240 elemental analyzer on samples retained on precombusted Whatman GF/F glass-fiber filters. Dissolved inorganic carbon (DIC) was measured on a Dohrmann PR-1 carbon analyzer (Goldman, 1979).

Portions of those samples collected before sunrise on each study day were filtered through 202  $\mu\text{m}$  Nitex mesh and preserved in Lugol's solution; phytoplankton cell counts and identifications were performed at a later date on 50 ml subsamples according to the Utermöhl method (Lund et al., 1958).




Fig. 1. Sequential incubations involving time-course measurements of  $^{14}\text{CO}_2$  fixation by natural marine populations in Vineyard Sound, Massachusetts, May 14, 1981. (A) Solar irradiance. (B)  $^{14}\text{CO}_2$  fixation of samples incubated at 60%  $I_0$ : incubations started at 0530 ●, 0800 ○, 1100 ▲, 1400 △, 1700 ■. Curves in all figures were drawn by inspection. Solid lines: presumed linear uptake; dashed lines: possible rapid response to changing irradiation




Fig. 2. Sequential incubations involving time-course measurements of  $^{14}\text{CO}_2$  fixation by natural marine populations in Vineyard Sound, Massachusetts, June 11, 1981. (A) Solar irradiance. (B)  $^{14}\text{CO}_2$  fixation of samples incubated at 60%  $I_0$ : incubations started at 0500 ●; filtered samples \*, 0800 ○, 1100 ▲, 1400 △, 1700 ■. (C) Full day incubations at 100%  $I_0$  (○); alternating light between 30%  $I_0$  and 60%  $I_0$  in 0.5 h intervals △. Dashed line: full day incubation curve at 60%  $I_0$  from Panel (B)

## RESULTS

### Climatic conditions and water characteristics

Climatic conditions varied tremendously between study days over the 6 mo period (Table 1). Only 1 of the 6 d, July 11, was cloudless (Panel A in Fig. 1 to 6). On 2 occasions otherwise sunny days were interrupted by mid-day clouds: heavy overcast on May 14 (Fig. 1 A) and intermittent clouds on June 11 (Fig. 2 A). September 29, in contrast, was characterized by morning overcast, followed by mid-day clearing (Fig. 5 A). On 2 dates, August 5 and October 29, full day overcast prevailed (Fig. 4 A and 6 A). The August 5 experiment, in fact, had to be terminated at 1600 h due to a thunderstorm. With the exception of the 2 overcast dates, total daily irradiation on the study days varied between about 5000 and 8000  $\text{W h m}^{-2}$  (Table 1). Water temper-

Table 2. Summary of particulate carbon and particulate nitrogen data of individual samples obtained from Vineyard Sound, Massachusetts during 6 mo study period

Date 1981	Sampling time	Particulate carbon ( $\mu\text{g C l}^{-1}$ )	Particulate nitrogen ( $\mu\text{g N l}^{-1}$ )	C:N ratio (wt:wt)
May 14	0530	264	31	8.5
	0800	627	77	8.1
	1100	457	59	7.7
	1400	962	103	9.3
	1700	847	99	8.6
Jun 11	0500	750	80	9.4
	0800	730	81	9.0
	1100	685	74	9.3
	1400	690	73	9.5
	1700	985	79	12.5
Jul 16	0500	748	89	8.4
	0800	454	66	6.9
	1100	632	79	8.0
	1400	691	69	10.0
	1700	633	70	9.0
Aug 5	0500	623	86	7.2
	0800	573	78	7.3
	1100	1110	136	8.2
	1400	670	85	7.9
Sep 29	0615	637	77	8.3
	0915	740	92	8.0
	1215	718	91	7.9
	1515	1083	134	8.1
Oct 29	0600	340	41	8.3
	0900	261	33	7.9
	1200	190	24	7.9
	1430	152	20	7.6

atures varied in a more predictable fashion, increasing from 10.5 °C on May 14 to a maximum of 23 °C on August 5, followed by a decrease to 12.5 °C on October 29 (Table 1).

We could not find any evidence of diel or seasonal changes in the concentrations of particulate carbon and particulate nitrogen during the first five study days (Table 2). For the most part, the particulate carbon and particulate nitrogen concentrations varied irregularly between about 600–750  $\mu\text{g l}^{-1}$  and 60–90  $\mu\text{g l}^{-1}$ , respectively, over the course of each study day, although higher and lower concentrations of both constituents were measured from time to time. Concentrations of both particular constituents, not only were minimal on October 29, but decreased over the course of the day. The particulate C:N ratio was reasonably constant at about 8 (wt:wt) during most of the study period, although on June 11 it was > 9.

Assorted diatoms and cryptomonads were the most abundant phytoplankton species during the six month period (Table 3). The cryptomonads *Chroomonas*

*amphioxeia* and *C. minuta* comprised between 64 and 85 % of the total number of phytoplankton cells between May 14 and July 16. Assorted diatoms made up the balance of the cell population during this period. The diatom fraction (dominated by *Leptocylin-drus danicus* on August 5, and *Rhizosolenia stolter-fothii* on September 29) increased from late summer through early fall, reaching 92 % of the total cell population by September 29; the cryptomonad contribution, in contrast, decreased dramatically to only 7 % of the total population during this same period. By October 29 when biomass concentrations were lowest (Table 2) there was an even balance between diatoms and cryptomonads (43 % each), with an unidentified dinoflagellate comprising the remainder of the population.

### Time course of primary production

The 2 patterns of  $^{14}\text{C}$  uptake we observed seemed to be correlated with season. On the first 2 study dates in May and June (Fig. 1 B and 2 B) and again in September (Fig. 5 B) rapid initial uptake, which occurred during all but the last daily back to back incubation of each day, was followed by reduced uptake over the balance of the incubation period and, in some cases, cessation of uptake late in the day. The duration over which the initial rapid phase was sustained seemed to decrease with increasing time of day. This type of response was most apparent on May 14 (Fig. 1 B) and on September 29 (Fig. 5 B). In fact, total  $^{14}\text{C}$  uptake over the first 3 h of the day on these 2 dates accounted for 51 % and 53 %, respectively, of the full day production (Table 4). In contrast, uptake during the last incubation of each day, not only was linear, but was about as rapid as during the first several hours of the first daily incubation.

Although we drew straight (solid) lines through the data to indicate the trends of uptake in Fig. 1 B, 2 B, and 5 B, it appeared, particularly during the May 14 and September 29 experiments, that mid-day suppression of  $^{14}\text{C}$  uptake occurred when there were no or intermittent clouds. This suppression seemed to be alleviated either when heavy clouds passed over, as on May 14, or during the latter part of the afternoon on all 3 dates when sunlight intensity decreased significantly. This trend, as shown by the dashed lines in Fig. 1 B, 2 B, and 5 B, was apparent in all but the last incubation of each day. It was impossible to assign error bars to the individual datum points in each experiment because we only sampled once at each time interval and the incubations were not replicated. Thus there is some uncertainty as to whether the apparent rapid responses to changing sunlight were real. The consistency of the trends from one incubation to

Table 3. Summary of major phytoplankton species present in Vineyard Sound, Massachusetts at time of pre-dawn sampling during 6 mo study period between May 14, 1981 and October 29, 1981

Species	Cell Count (cells ml <sup>-1</sup> )					
	May 14	Jun 11	Jul 16	Aug 5	Sep 29	Oct 29
Diatoms						
<i>Navicula</i> sp.					13	
<i>Cylindrotheca closterium</i>	3	42	4			11
<i>Leptocylindrus danicus</i>	82			106	26	33
<i>Rhizosolenia imbricata</i> var. <i>schrubsolei</i>			4	40		4
<i>Rhizosolenia delicatula</i>			12	24		5
<i>Rhizosolenia calcar-avis</i>				13		
<i>Rhizosolenia setigera</i>		11				
<i>Rhizosolenia stolterfothii</i>	13				314	23
<i>Skeletonema costatum</i>		21				
<i>Skeletonema menzili</i>		37				
<i>Thalassiosira pseudonana</i>			14			
Assorted other diatoms	18		9	24	8	16
Dinoflagellates						
20 µm unident. naked dinoflag.						28
Assorted other dinoflagellates		5	11	3	5	
Cryptomonads						
<i>Pyramimonas</i> sp.	8	26	11		3	
<i>Chroomonas amphioxeia</i>	145	16				9
<i>Chroomonas minuta</i>	26	227	275	153	21	48
Unknown sp.	21	137	21			21
Assorted other cryptomonads	8		1		3	14
Others						
Unidentified 3 µm sp.		42		79		
<i>Dunaliella</i> sp.				5		
Total (Percentage)						
Diatoms	36	20	12	47	92	43
Dinoflagellates	0	<1	3	<1	1	14
Cryptomonads	64	73	85	34	7	43
Other and Unidentified	0	7	0	19	0	0

Table 4. Summary of total and incremental daily primary production estimates at 60% I<sub>0</sub> determined by full day, single end-point measurements (TP), by summed back to back 3 h rates (Σ P), and by first h rate from each back to back incubation extrapolated to 3 h and summed over full day (Σ P')

Date 1981	Total daily production (µg C l <sup>-1</sup> )			Incremental daily production % Σ P (% TP)					Σ P : TP ratio	Σ P' : TP ratio
	TP	Σ P	Σ P'	0-3 h	3-6 h	6-9 h	9-12 h	12-15 h		
May 14	22.2	33.6	39.4	33 (51)	18	14	15	20	1.5	1.8
Jun 11	148.2	141.5	169.8	29 (19)	17	25	15	23	0.9	1.2
Jul 16	55.8	47.9	82.1	23 (20)	14	14	18	31	0.9	1.5
Aug 5	139.0	136.2	128.4	19 (19)	27	22	32*		1.0	0.9
Sep 29	51.4	117.2	145.5	35 (53)	22	10	33		2.0	2.8
Oct 29	75.0	113.8	127.1	22 (33)	32	32	14*		1.5	1.7

\* Two h incubation

another and the similarity in the results between the May and September studies, however, appears to be more than coincidental, thereby providing us with some assurance that the rapid changes in <sup>14</sup>C uptake were real.

The second uptake pattern was one in which linear uptake was more persistent during each incubation. This response was observed on July 16 (Fig. 3 B) when there were virtually no clouds and high total irradiation (8018 W h m<sup>-2</sup>) and on both August 5 (Fig. 4 B) and

October 29 (Fig. 6 B), days marked by heavy overcast and very low total irradiation (3091 and 1057  $W h m^{-2}$ , respectively).

**Response to different light regimes**

A common feature of each experiment was that regardless of weather conditions, uptake of  $^{14}C$  over the first few hours of the full day incubations was maximal and unaffected by variations in the imposed light regime (Fig. 2 C-6 C). On sunny days, however, we found that  $^{14}C$  uptake quickly ceased after a few hours in those incubation vessels exposed to full sunlight for the entire day (Fig. 2 C, 3 C and 5 C) and that the mid-day suppression in productivity (which was common to the incubations at 60 %  $I_0$ ) was less pronounced when the light was reduced to 30 %  $I_0$  (Fig. 5 C). Rapid suppression of  $^{14}C$  uptake likewise occurred on a sunny day when the last incubation of the day was performed without screening, although recovery in uptake under these conditions seemed to occur just before sunset (Fig. 5 C). On cloudy days there was virtually no impact of light attenuation (down to 30 %  $I_0$ ) on the time course of  $^{14}C$  uptake over the entire day (Fig. 4 C and 6 C).

Alternating the light between 30 %  $I_0$  and 60 %  $I_0$  had about the same effect on  $^{14}C$  uptake as did con-




Fig. 3. Same as Fig. 2, except performed on July 16, 1981 and  $\Delta$  in Panel (C) represents alternating light between 60 %  $I_0$  and 100 %  $I_0$  in 0.5 h intervals




Fig. 4. Same as Fig. 2, except performed on August 5, 1981 and  $\Delta$  in Panel (C) represents incubation under 30 %  $I_0$




Fig. 5. Same as Fig. 4, except performed on September 29, 1981 and  $\bullet$  in Panel (C) represents incubation starting at 1500 under 100 %  $I_0$ . Dashed line: 1500 incubation curve at 60 %  $I_0$  from Panel (B)




Fig. 6. Same as Fig. 4, except performed on October 29, 1981

stant light reduced to 60 %  $I_0$  (Fig. 2 C). Similarly, there was no appreciable improvement in  $^{14}\text{C}$  uptake in the vessel exposed to alternating light between 60 %  $I_0$  and 100 %  $I_0$  compared to the one maintained constantly at full sunlight (Fig. 3 C).

### Primary productivity estimates

We estimated primary productivity in 3 different ways. First, total daily production based on single end-point measurements from the full day incubations was normalized to an hourly rate (TP). By this technique we observed that rates of productivity on a given day generally were saturated by at least 30 %  $I_0$ . These rates were sustained or only slightly altered at 60 %  $I_0$ , but did decrease significantly under full sunlight on June 11, July 16 and September 29 when total irradiation exceeded  $5000 \text{ W h m}^{-2}$  (Table 1). On the 2 overcast days, August 5 and October 29, exposure to full sunlight had little adverse effect on productivity.

There was no discernable seasonal trend in productivity rates (Table 1). Maximum rates of  $\sim 13 \mu\text{gC l}^{-1} \text{ h}^{-1}$  occurred on August 5 when water temperature was  $23^\circ\text{C}$  and total irradiation was  $3091 \text{ W h m}^{-2}$ , whereas lowest rates of  $1.4 \mu\text{gC l}^{-1} \text{ h}^{-1}$  were found on May 14 when water temperatures were  $10.5^\circ\text{C}$  and total irradiation was  $6426 \text{ W h m}^{-2}$ . The general trend was for

productivity to be highest when days were overcast and lowest when sunny weather prevailed.

The other estimates of productivity were based on summing total production at 60 %  $I_0$  over the first 3 h of each back to back incubation during the course of a day ( $\Sigma P$ ) and extrapolating the first hour rate from each back to back incubation to 3 h total production values and then summing these values as in the previous method ( $\Sigma P'$ ). These estimates of total daily production were then compared with estimates based on single-end-point measurements from the full day incubations at 60 %  $I_0$  (TP). Accordingly, we found the ratio  $\Sigma P : \text{TP}$  to be  $> 1$  in the spring and fall, ranging between 1.5 on May 14 and October 29 to 2.0 on September 29. During the summer period  $\Sigma P : \text{TP}$  was between 0.9 and 1.0 (Table 4). With the exception of the August 5 experiment, the ratio  $\Sigma P' : \text{TP}$  was greater than  $\Sigma P : \text{TP}$  on each study date (Table 4).

On the 3 occasions when we compared the time course of total  $^{14}\text{C}$  uptake (particulates plus excretion products) with uptake into the particulate fraction only (Fig. 2 B, 4 B, and 5 B), we found no difference between the 2 methods, indicating that there was little, if any, excretion of dissolved organics.

### Mixing in Vineyard Sound

Our sampling station in Vineyard Sound was located in a shallow region ( $\sim 10 \text{ m}$  depth) where tidal flushing is strong and vertical mixing is virtually complete throughout the year, as judged from numerous bathythermograms compiled at WHOI (Bumpus et al., 1971). Our inability to observe thermal stratification at this location over a 15 mo period in 1979–1980 (unpubl.) in conjunction with another study (Glibert et al., 1982) confirms this conclusion. In addition,  $> 97\%$  of incident irradiation always was attenuated at 10 m depth (unpubl.). Thus, although we have no information on vertical mixing velocities, it appears reasonable to assume that the resident phytoplankton populations that we collected were never photoadapted to any particular light level.

## DISCUSSION

### Time course of photoinhibition

As summarized in Harris (1978, 1980), photoinhibition is a well-established, if poorly understood phenomenon. Often little or no distinction is made between the various processes that can cause a decrease in photosynthesis at supersaturating light intensity (e.g. UV and visible photochemical decay, photorespiration, photooxidation); thus in most

studies, including this one, the term 'photoinhibition' is used to lump together all processes leading to a decrease in the rate of  $\text{CO}_2$  fixation at high irradiation.

The manifestation of photoinhibition during bottle incubations of natural water samples depends on at least 3 key factors: (1) light history of the confined population (stratified versus well-mixed phytoplankton), (2) incubation irradiation, and (3) duration of exposure to greater than saturating irradiation. The last factor, although recognized as being of crucial importance in the interpretation of bottle productivity measurements (Harris, 1978, 1980) and in the modeling of P versus I relationships (Platt et al., 1980), is poorly characterized.

From the earlier time-course studies of Myers and Burr (1940) and Kok (1956), it was evident that algae, when exposed to light intensities far greater than saturating, were capable of sustaining maximum photosynthetic rates for short (minutes) duration before a progressive decay in photosynthesis occurred. Others have observed this phenomenon in studies on cultured and natural populations of phytoplankton (Takahashi et al., 1971; Harris and Piccinin, 1977; Belay, 1981). Harris (1980) suggested that photoinhibitory effects not only would become progressively manifested at light intensities  $> 200 \mu\text{E m}^{-2} \text{ s}^{-1}$  ( $\sim 50 \text{ W m}^{-2}$ ), but that

of Harris (1980). In fact, on close scrutiny of the May 14 and September 29 results when photoinhibition was most apparent, it appeared that, even though the light level leading to photoinhibition was ca.  $130 \text{ W m}^{-2}$  (as judged by the light level corresponding to the point of departure from initially rapid  $\text{CO}_2$  fixation during the first incubation of the day [Fig. 1 B and 5 B]), it was possible for initially rapid photosynthetic rates to be sustained for 15 to 30 min periods during mid-day incubations when light levels had exceeded  $300 \text{ W m}^{-2}$  (Table 5).

However, during summer (June–August), not only was photoinhibition hardly evident during the 60%  $I_0$  incubation experiments, but in the unscreened experiments it seemed to occur at increasingly higher sunlight intensities as the summer progressed. For example, on July 16 (a cloudless day) photoinhibition developed only when irradiation reached  $400 \text{ W m}^{-2}$  (Fig. 3 C); and on August 5 (a variably cloudy day) photoinhibition never was experienced even though sunlight intensities reached  $600 \text{ W m}^{-2}$  at times (Fig. 4 C). The reasons for this seasonal change in the patterns of photoinhibition are unclear, but the results are strikingly similar to those of Marra (1978b) and Hobson and Hartley (1983), both of whom found no evidence for suppression in surface productivity of coastal waters

Table 5. Effect of time of day when incubation was initiated on durations of exposure and the irradiation leading to strong photoinhibition on May 14, 1981 and on September 29, 1981

Date 1981	Time of day incubation initiated	Starting irradiation ( $\text{W m}^{-2}$ )	Irradiation for onset of strong photoinhibition* ( $\text{W m}^{-2}$ )	Duration of exposure before onset of strong photoinhibition (min)
May 14	0530	0	130	180
	0800	85	200	60
	1100	430	540	30
	1400	360	360	15
Sep 29	0630	0	130–260 (130)**	240 (135)***
	0915	150	320	75
	1215	330	325	15

\* Irradiation at which major reduction in slope of  $^{14}\text{C}$  uptake curve occurred (incubation vessels exposed to 60%  $I_0$ , Fig. 1 B and 5 B)  
 \*\* Irradiation leading to complete inhibition (incubation vessels exposed to 100%  $I_0$ , Fig. 5 C)  
 \*\*\* Time for complete inhibition to occur

exposure to sunlight irradiation  $> 800\text{--}1000 \mu\text{E m}^{-2} \text{ s}^{-1}$  ( $\sim 200\text{--}250 \text{ W m}^{-2}$ ) for periods exceeding 30 min would lead to severe photoinhibition and that maximum photosynthesis under such conditions was possible for only 10 to 20 min.

Our observation that the onset of photoinhibition generally was preceded by a period of rapid initial  $\text{CO}_2$  fixation, and that the duration of this period was inversely related to the light level at the start of an incubation (Table 5), is in accord with the conclusions

during late spring–early summer when surface irradiation was maximal, but pronounced surface suppression before or after this period. Hobson and Hartley (1983) attributed this seasonal suppression of photosynthesis to a greater susceptibility to UV-inhibition by diatoms, species which were dominant during the period of surface suppression. We cannot rule out this possibility in our experiments, but we suspect that little, if any, UV light passed the double glass walls and layer of cooling water of our incubation vessels.

Considering the intense mixing of Vineyard Sound throughout the year, it is also difficult to invoke a photoadaptive mechanism for the summer resistance to photoinhibition we observed. Major changes in water temperature (10.5 °C on May 14 to 23.0 °C on August 5) and phytoplankton species (Table 3) did occur during this period and one or both of these variables may be correlated with the seasonal change we saw in photoinhibition light intensities.

A strong interrelationship exists between temperature and light in regulating photosynthesis in phytoplankton (Cloern, 1977; Yoder, 1979). Yet, little is known of any possible relationship between temperature and the time-course of photoinhibition, although, as summarized by Harris (1980), recovery from photoinhibition appears to be temperature-dependent. In an earlier study (Goldman et al., 1981) we found that under constant light intensity the incubation period for which linear CO<sub>2</sub> fixation was maintained by the marine diatom *Thalassiosira weissflogii* decreased systematically with decreasing temperature. Whether this temperature-dependent reduction in photosynthetic capacity over time was a true photoinhibitory response is not known, but the results are suggestive of a role for temperature in regulating the exposure period over which a particular P vs I relationship holds.

Concomitant with the increased resistance to photoinhibition between May 14 and July 16 that occurred in the incubated samples there was a decrease in the diatom fraction and an increase in the cryptomonads (Table 3). Cryptomonads, because of their small size relative to the diatoms, probably contributed considerably less to total primary production than their numbers would indicate.

Comparisons of the photosynthetic characteristics of laboratory-grown cryptomonads and diatoms provide no real clue as to why the Vineyard Sound summer phytoplankton were uniquely resistant to photoinhibition. Light saturation in many diatoms (Richardson et al., 1983) and in the few studied cryptomonads (Faust and Gantt, 1973; Cloern, 1977; Morgan and Kalff, 1979) seems to occur at intensities in the range 75 to 150  $\mu\text{E m}^{-2} \text{s}^{-1}$  ( $\sim 40$  to  $80 \text{ W m}^{-2}$ ). Similarly, photoinhibitory light intensities in many diatoms and in one freshwater cryptomonad (Morgan and Kalff, 1979) have been reported to be typically in the range of 150 to  $> 330 \mu\text{E m}^{-2} \text{s}^{-1}$  ( $\sim 20$  to  $40 \text{ W m}^{-2}$ ), although the higher values are from one study in which meticulous care was taken to insure long-term adaptation to the imposed light intensities (Brand and Guillard, 1981). Yet, we never observed photoinhibition at any time during the 6 mo study until light intensities  $> 130 \text{ W m}^{-2}$  were reached. Moreover, because the maximum slopes of the times-course curves generally were attained at the onset of the first morning incubations,

sometimes in vessels attenuated to 30 % I<sub>0</sub> (Fig. 3 C to 6 C), and again during the last incubations of the day at 60 % I<sub>0</sub>, light saturation levels always appeared to be very low.

Although little is known of the light response characteristics of natural cryptomonad populations, there is some convincing evidence that natural diatom populations become photoinhibited at much lower sunlight levels during bottle incubations than do a variety of other classes of phytoplankton (Harris and Piccinin, 1977; Platt et al., 1980). Platt et al., for example, found that even though natural populations of marine diatoms from well-mixed waters photoinhibited at light levels far above the expected values ( $\sim 140 \text{ W m}^{-2}$ ), other classes of species, including flagellates from the same waters consistently did not display photoinhibition at sunlight intensities  $> 600 \text{ W m}^{-2}$ . These results are in accord with our findings and lead us to believe that the mechanisms controlling photoinhibition in natural populations are far different than what might be ascertained from a laboratory P vs I experiment. The notion that cryptomonads may, as a rule, be more resistant to photoinhibition than diatoms does not explain the August 5 results when the diatom fraction had risen back to 47 % of the total cell population and still there was no evidence of photoinhibition.

It is difficult to speculate further on the causes of the seasonal variations in photoinhibition we observed. Our results point toward rather complicated interactions between light, temperature, species, and exposure period controlling observed photoinhibition. Light intensity alone probably was not the main contributing factor in our experiments because total daily irradiation during May was comparable to mid-summer intensities (Table 1).

The link we observed between apparent recovery from photoinhibition and the combination of exposure duration and photoinhibiting light intensity (Fig. 5 C) is consistent with the early findings of Myers and Burr (1940). They showed that: (1) the greater the photoinhibiting light intensity and duration of exposure, the longer the period required before cultures of microalgae were able to recover from photoinhibition when placed under subsaturating light, and (2) photoinhibition was permanent above critical light levels and exposure periods. The differences between the rapid recovery from photoinhibition we saw on both May 14 when a cloud bank rapidly passed over at mid-day (Fig. 1 B) and September 29 when surface irradiation decreased late in the day (Fig. 5 C) and the permanent photoinhibition that occurred on July 16 under an alternating light regime (Fig. 3 C) may also have been species-related. Cryptomonads, although possibly able to withstand high supersaturating light intensities for short periods, may become permanently damaged

when photoinhibition finally is manifested. Initial photoinhibition could be the result of photochemical decay, which is repairable (Sorokin, 1960), followed by photooxidative processes leading to complete chlorophyll degradation (Myers and Burr, 1940). Diatoms, in contrast, may exhibit photochemical decay at relatively lower light levels, perhaps as a protective strategy against photooxidation so that rapid recovery is possible once the light level is reduced late in the afternoon or at other times either through downward transport of cells in the water column or by cloud cover. Differences in the accessory pigment structure between diatoms and cryptomonads (Goodwin, 1974) may have something to do with these different patterns of photoinhibition.

#### Bottle photoinhibition and primary production estimates

The extent to which water-column productivity in a shallow environment such as Vineyard Sound may be underestimated by use of bottle incubations is not easily predictable. The unexpected nature of the seasonal response of natural populations to full sunlight in our study and in those of Marra (1978b) and Hobson and Hartley (1983) raises perplexing questions about the very nature of photoinhibition and its impact on measurements of seasonal changes in water column productivity. There seems little doubt, however, that had we based our estimates of productivity on single endpoint sampling from full day incubations, serious

underestimates of surface productivity would have resulted in all but the summer experiments. What effect these underestimates would have had on measurements of integrated water column productivity in Vineyard Sound is difficult to ascertain because our studies were limited to several light levels  $\geq 30\%$   $I_0$ . By comparison, there has been a clear trend for  $\Sigma P:TP$  integrated over the water column to be ca. 1.3 to 1.5 in the few studies in which back to back incubations have been carried out, and, as in our study (Table 4), to be as high as 2 to 3 for the surface and near surface (Table 6). Jewson and Wood (1975) do point out, however, that when the depth of the mixed layer is greater than that of the euphotic zone, real reductions in total water column productivity result which could offset the apparent underestimate due to surface photoinhibition, as determined from a bottle incubation.

One conclusion, which is well documented in our study and confirmed in other studies – e.g. Rodhe (1958), Vollenweider and Nauwerck (1961) – is that the 'mid-day suppression' in primary production we observed was, to a large extent, an incubation artifact. To illustrate this point, consider that had only a single full-day incubation been employed during May and September, we would have drawn the mistaken conclusion that  $> 50\%$  of the total daily production had occurred during the first 3 h of the day (Table 4). Use of back to back 3 h incubations, by raising the total daily production, reduced those early morning contributions to  $\leq 35\%$ ; but, as seen by our estimates of  $\Sigma P:TP$  (Table 4), even 3 h incubations may have been too long in our case to minimize the effects of photoinhibition.

Table 6. Comparison of primary productivity estimates based on summed short-term ( $\Sigma P$ ) and full day, single end-point measurement (TP) from reported studies involving  $^{14}C$  bottle incubations

Water body	Date of study	Total incubation duration (h)	Incubation interval (h)	% $I_0$	$\Sigma P:TP$	Integrated $\Sigma P:TP^*$	Source
Lake Erken, Sweden (Freshwater)	17 May 1956	19.0	3–4	30–80	1.3–2.9	1.3	Rodhe (1958) Vollenweider and Nauwerck (1961)
	30 May 1956	20.0	4	30–55	1.3–1.4	1.3	
	1 Jul 1956	18.0	3–4	45–70	1.5–2.1	1.3	
Lake Werowrap, Australia (Freshwater)	2 Mar 1970	13.5	4–5	–	–	1.2	Hammer et al. (1973)
	23 Aug 1970	10.5	3.5	–	–	1.5	
	17 Sep 1970	12.0	3.5–5	–	–	1.6	
	26 Sep 1970	12.5	3.5–5	–	–	1.3	
North Equatorial Current (Marine)	30 Nov 1978	14.0	2	30	2.8	–	Gieskes et al. (1979)
Kaneohe Bay Hawaii (Marine)	13 Nov 1976	12.0	1	**	1.0	–	Redalje and Laws (1981)

\* Integrated values of  $\Sigma P$  and TP over entire water column  
 \*\*  $I_0$  not given; incubated at 1 m depth

The choice between 'long' and 'short' incubations has been a continuing dilemma for phytoplankton ecologists. On the one hand, advocates for incubations lasting for 4 to 24 h have argued that such durations are a necessary prerequisite for establishing  $^{14}\text{C}$  isotopic equilibrium so that net primary production is measured (Eppley and Sharp, 1975; Hobson et al., 1976; Marra et al., 1981). Li and Harrison (1982), in fact, have shown that even 24 h incubations do not guarantee the attainment of isotopic equilibrium. On the other hand, various problems with confinement in bottles have led others to promote the use of short (up to several h) incubations, even though this technique is limited to measurement of gross productivity (Harris and Piccinin, 1977; Venrick et al., 1977; Goldman et al., 1981).

From our own results, and from those summarized in Table 6, we conclude that the choice of incubation period cannot be established without considerable knowledge of the physics and biology of the water mass in question. Clearly, many factors are involved in this choice. Well-mixed coastal and upwelled phytoplankton may, as a general rule, be the most likely to exhibit unrepresentative photoinhibition during long confinement in bottles. Seasonal variations in this type of response for a given water body complicate tremendously the choice of incubation duration. Our own preference is to perform back to back incubations that span the entire day, with each incubation, not only involving time-course measurements, but continuing until sunset. Such an approach provides the greatest flexibility for data interpretation when trying to sort out artifacts of confinement in bottles from real water column effects. However, when integrated water column measurements of primary productivity are required, the increasing number of time-course incubations that are required as the day progresses probably negates the utility of this approach. The protocols adapted by Rodhe (1958) and Vollenweider and Nauwerck (1961), which are similar to those suggested above, but without time-course measurements, seem to be a reasonable trade-off.

*Acknowledgements.* This study was supported by Grant No. OCE-8024445 from the National Science Foundation. We thank Dr. Edward J. Carpenter for cell counts and species identifications and Carol B. Riley for her able technical assistance.

#### LITERATURE CITED

- Barnett, A. M., Hirota, J. (1967). Changes in the apparent rate of  $^{14}\text{C}$  uptake with length of incubation period in natural phytoplankton populations. *Limnol. Oceanogr.* 12: 349-353
- Belay, A. (1981). An experimental investigation of inhibition of phytoplankton photosynthesis at lake surfaces. *New Phytol.* 89: 61-74
- Brand, L. E., Guillard, R. R. L. (1981). The effects of continuous light and light intensity on the reproduction rates of twenty-two species of marine phytoplankton. *J. exp. mar. Biol. Ecol.* 50: 119-132
- Bumpus, D. F., Wright, W. R., Vaccaro, R. F. (1971). Sewage disposal in Falmouth, Massachusetts. II. Predicted effect on the proposed outfall. *J. Boston Soc. Civil Engrs.* 58: 255-277
- Cloern, J. E. (1977). Effects of light intensity and temperature on *Cryptomonas ovata* (Cryptophyceae) growth and nutrient uptake rates. *J. Phycol.* 13: 389-395
- Eppley, R. W., Sharp, J. H. (1975). Photosynthetic measurements in the Central North Pacific: the loss of carbon in 24-hour incubations. *Limnol. Oceanogr.* 20: 981-987
- Faust, M. A., Gantt, E. (1973). Effect of light intensity and glycerol on the growth, pigment composition, and ultrastructure of *Chroomonas* sp. *J. Phycol.* 9: 489-495
- Gallegos, C. L., Hornberger, G. M., Kelly, M. G. (1980). Photosynthesis-light relationships of a mixed culture of phytoplankton in fluctuating light. *Limnol. Oceanogr.* 25: 1082-1092
- Gallegos, C. L., Platt, T. (1982). Phytoplankton production and water motion in surface mixed layers. *Deep Sea Res.* 29: 65-76
- Gieskes, W. W. C., Kraay, G. W., Baars, M. A. (1979). Current  $^{14}\text{C}$  methods for measuring primary production: gross underestimates in oceanic waters. *Neth. J. Sea Res.* 13: 58-78
- Glibert, P. M., Goldman, J. C., Carpenter, E. J. (1982). Seasonal variations in the utilization of ammonium and nitrate by phytoplankton in Vineyard Sound, Massachusetts, USA. *Mar. Biol.* 70: 237-249
- Goldman, J. C. (1979). Bioengineering aspects of inorganic carbon supply to mass algal cultures. In: *Proceedings Third Annual Biomass Energy Systems Conference*. Rep. SERI/TP-33-285, Solar Energy Research Institute, Golden, Colorado, p. 25-32
- Goldman, J. C., Taylor, C. D., Glibert, P. M. (1981). Nonlinear time-course uptake of carbon and ammonium by marine phytoplankton. *Mar. Ecol. Prog. Ser.* 6: 137-148
- Goodwin, T. W. (1974). Carotenoids and biliproteins. In: Stewart, W. D. P. (ed.) *Algal physiology and biochemistry*. University of California Press, Berkeley, p. 176-203
- Hammer, U. T., Walker, K. F., Williams, W. D. (1973). Derivation of daily phytoplankton production estimates from short-term experiments in some shallow, eutrophic Australian saline lakes. *Aust. J. mar. Freshwat. Res.* 24: 259-266
- Harris, G. P. (1978). Photosynthesis, productivity and growth. The physiological ecology of phytoplankton. *Ergebn. Limnol.* 10: 1-171
- Harris, G. P. (1980). The measurement of photosynthesis in natural populations of phytoplankton. In: Morris, I. (ed.) *The physiological ecology of phytoplankton*. University of California Press, Berkeley, p. 129-187
- Harris, G. P., Lott, J. N. A. (1973). Light intensity and photosynthetic rates in phytoplankton. *J. Fish. Res. Bd Can.* 30: 1771-1778
- Harris, G. P., Piccinin, B. B. (1977). Photosynthesis by natural phytoplankton populations. *Arch. Hydrobiol.* 80: 405-457
- Hobson, L. A., Hartley, F. A. (1983). Ultraviolet irradiance and primary production in a Vancouver Island fjord, British Columbia, Canada. *J. Plankton Res.* 5: 325-331
- Hobson, L. A., Morris, W. J., Pirquet, K. T. (1976). Theoretical and experimental analysis of the  $^{14}\text{C}$  technique and its use in studies of primary production. *J. Fish. Res. Bd Can.* 33: 1715-1721

- Ichimura, S., Saijo, Y. (1958). On the application of  $^{14}\text{C}$ -method to measuring organic matter production in the lake. Bot. Mag. Tokyo 71: 174-180
- Jewson, D. H., Wood, R. B. (1975). Some effect on integral photosynthesis of artificial circulation of phytoplankton through light gradients. Verh. int. Verein. theor. Limnol. 19: 1037-1044
- Jones, R. I. (1978). Adaptations to fluctuating irradiance by natural phytoplankton communities. Limnol. Oceanogr. 23: 920-926
- Kok, B. (1956). On the inhibition of photosynthesis by intense light. Biochim. biophys. Acta 21: 234-244
- Li, W. K. W., Harrison, W. G. (1982). Carbon flow into the end-products of photosynthesis in short and long incubations of a natural phytoplankton population. Mar. Biol. 72: 175-182
- Lund, J. W. G., Kilpling, C., LeCren, E. D. (1958). The inverted microscope method of estimating algal numbers and the statistical basis of estimation by counting. Hydrobiologia 11: 143-170
- Marra, J. (1978a). Effect of short-term variations in light intensity on photosynthesis of a marine phytoplankton. A laboratory simulation study. Mar. Biol. 46: 191-202
- Marra, J. (1978b). Phytoplankton photosynthetic response to vertical movement in a mixed layer. Mar. Biol. 46: 203-208
- Marra, J., Landrian G., Ducklow, H. W. (1981). Tracer kinetics and plankton rate processes in oligotrophic oceans. Mar. Biol. Lett. 2: 215-223
- Morgan, K. C., Kalff, J. (1979). Effect of light and temperature interactions on growth of *Cryptomonas erosa* (Cryptophyceae). J. Phycol. 15: 127-134
- Myers, J., Burr, G. O. (1940). Studies on photosynthesis. Some effects of light of high intensity on *Chlorella*. J. gen. Physiol. 24: 45-67
- Platt, T., Gallegos, C. L., Harrison, W. G. (1980). Photoinhibition of photosynthesis in natural assemblages of marine phytoplankton. J. mar. Res. 38: 687-701
- Redalje, D. G., Laws, E. A. (1981). A new method for estimating phytoplankton growth rates and carbon biomass. Mar. Biol. 62: 73-79
- Richardson, K., Beardall, J., Raven, J. A. (1983). Adaptation of unicellular algae to irradiance: an analysis of strategies. New Phytol. 93: 157-191
- Rodhe, W. (1958). The primary production of lakes: some results and restrictions of the  $^{14}\text{C}$  method. Rapp. Cons. Explor. Mer 144: 122-128
- Savidge, G. (1980). Photosynthesis of marine phytoplankton in fluctuating light regimes. Mar. Biol. Lett. 1: 295-300
- Schindler, D. W., Holmgren, S. K. (1971). Primary production and phytoplankton in the experimental lakes area, North-western Ontario and other low carbonate waters, and a liquid scintillation method for determining  $^{14}\text{C}$  activity in photosynthesis. J. Fish. Res. Bd Can. 28: 189-201
- Sorokin, C. (1960). Injury and recovery of photosynthesis in cells of successive developmental stages. The effect of light intensity. Physiologia Pl. 13: 687-700
- Takahashi, M., Shimura, S., Vamaguchi, Y., Fujita, Y. (1971). Photo-inhibition of phytoplankton photosynthesis as a function of exposure time. J. Oceanogr. Soc. Japan 2: 43-50
- Venrick, E. L., Beers, J. R., Heinbokel, J. F. (1977). Possible consequences of containing microplankton for physiological rate measurements. J. exp. mar. Biol. Ecol. 26: 55-76
- Vollenweider, R. A., Nauwerck, A. (1961). Some observations on the  $\text{C}14$  method for measuring primary production. Verh. internat. Verein. Limnol. 14: 134-139
- Yoder, J. A. (1979). Effect of temperature on light-limited growth and chemical composition of *Skeletonema costatum* (Bacillariophyceae). J. Phycol. 15: 362-370

This paper was submitted to the editor; it was accepted for printing on September 9, 1983