

Carotenoid pigments in microzooplankton

G. S. Kleppel¹, Evelyn J. Lessard²

¹Nova University Oceanographic Center, 8000 North Ocean Drive, Dania, Florida 33004, USA

²School of Oceanography, University of Washington, Seattle, Washington 98195, USA

ABSTRACT: The carotenoid pigments astaxanthin and canthaxanthin were measured in cultured and field-collected heterotrophic dinoflagellates (2 species), ciliates (3 species) and copepod eggs (1 species). The heterotrophic dinoflagellates contained either astaxanthin or canthaxanthin as a single dominant pigment. Conversely, the choreotrichous ciliates contained both carotenoids. A small amount of canthaxanthin was detected only at high concentrations in the oligotrich *Strombidium* sp. Copepod eggs contained astaxanthin and xanthophyll esters. The esters were not detected in the protozoans. Cellular pigment content varied both within and between taxonomic groups, but was low compared with phytoplankton. Microzooplankton which have low C:carotenoid ratios (e.g. some dinoflagellates), and large organisms, even when not heavily pigmented (e.g. *Favella ehrenbergii*) are readily detected by carotenoid analysis. Small, weakly pigmented ciliates, however, are more difficult to detect when they are present in low concentrations. The pigment content of a heterotrophic dinoflagellate (*Gymnodinium* sp.) cultured at $29 \mu\text{E m}^{-2} \text{s}^{-1}$ (low-light group) was approximately 42 % of the average pigment content of the same species cultured in the same food environment but at $290 \mu\text{E m}^{-2} \text{s}^{-1}$ (high-light group). Inadvertent starvation of the cultures in the low-light group for ≤ 3 d resulted in the loss of all measureable pigmentation.

INTRODUCTION

Carotenoid pigments occur in representatives of all phyla of marine organisms (Goodwin 1976) and are widely distributed among the zooplankton and micro-nekton (Cheeseman et al. 1967, Herring 1972). The carotenoids astaxanthin and canthaxanthin are common in metazoans (Castillo et al. 1982), and have been reported in trace concentrations in a few types of phytoplankton (Liaenn-Jensen 1979). They are thought to be photoprotective (Hairston 1979) and may have metabolic roles as well (Cheeseman et al. 1967, Byron 1982).

Certain carotenoids are restricted to specific taxonomic groups and, as such, are widely used as markers for coarsely describing plankton assemblages (Strain et al. 1944, Mantoura & Llewellyn 1983, Smith et al. 1987, Kleppel 1988). In addition, carotenoid content is roughly proportional to the biomass of bacteria (Liaaen-Jensen 1979) and phytoplankton (Burkill et al. 1987). A general proportionality was also reported between microzooplankton biomass and carotenoid pigment content, though the distribution and composition of pigments within taxa was not specified (Kleppel et al. 1988).

Most research on zooplankton pigmentation has focused on metazoans, notably the crustaceans (Herring 1968, Hairston 1979, Castillo et al. 1982). Systematic studies of microzooplankton pigmentation are scarce, especially for the protozoans. To our knowledge, the carotenoids of heterotrophic ciliates have not been described, although chlorophyll and phycoerythrin from sequestered chloroplasts of ingested algae have been documented (McManus & Fuhrman 1986, Stoecker et al. 1987). Carotenoid pigmentation in heterotrophic dinoflagellates, as distinct from their photosynthetic counterparts, has been discussed in several papers (Neveaux & Soyer 1976, Withers & Haxo 1978, Balch & Haxo 1984, Carreto 1985) though the amount of information on these pigments and their significance is small compared to the data on metazoan pigmentation.

The lack of information on protozoan pigmentation is not surprising given that the abundance, diversity and trophic importance of these organisms in the sea has only recently been recognized (e.g. see recent reviews by Sherr et al. 1986, Stoecker & Capuzzo 1990, Lessard 1991, and references therein). Ultimately, pigments

may be useful for describing microzooplankton distributions and understanding the physiology of these organisms.

In this paper we describe the pigments of cultured and field-collected microzooplankton (defined here as protozoans and metazoans smaller than 200 μm in diameter). The taxa that we have examined are typical representatives of the coastal microfauna. We also present the results of an experiment which examined the effects of light intensity on protozoan pigmentation.

MATERIALS AND METHODS

Two sets of measurements were performed. The first consisted of a general screening of pigments in 4 representative protozoan taxa. These were an athecate, heterotrophic dinoflagellate, *Gymnodinium* sp.; an aloricate, oligotrichous ciliate, *Strombidium* sp.; an aloricate choreotrichous ciliate, *Strombidinopsis* sp.; and a loricate choreotrich, *Favella ehrenbergii*, grown in culture. For comparative purposes, data on the pigments in copepod eggs and field-caught specimens of the thecate, heterotrophic dinoflagellate *Protoperdinium* sp. were described. The second set of measurements was performed to evaluate the differential effects of light intensity on pigment content in *Gymnodinium* sp.

Cultures were kindly provided by S. Strom and maintained on algal food mixtures in seawater with $f/2$ trace metals added (with 10^{-7}M EDTA; Gifford 1985). The food algae, maintained in $f/2$ minus Si, consisted of *Isochrysis galbana* (I), *Chroomonas* sp. (C), photosynthetic *Gymnodinium* sp. (G), and *Synechococcus* sp. (S). Mixtures of I, C and G were fed to *Favella ehrenbergii* and *Strombidinopsis* sp. I, C, G and S were fed to *Strombidium* sp. and I and S were fed to *Gymnodinium* sp. At the beginning of each experiment, inocula from stock cultures were transferred to fresh medium, and mixtures of exponentially growing food organisms were added (at ca $100\ \mu\text{g C l}^{-1}$ each). The cultures were incubated at $13\ ^\circ\text{C}$ under cool white, fluorescent lights, on a 14 h/10 h light/dark cycle. For general pigment screening (the first set of measurements), light intensity was $29\ \mu\text{E m}^{-2}\ \text{s}^{-1}$. To study the response of *Gymnodinium* to different light intensities (the second set), cultures were grown at 29 and $290\ \mu\text{E m}^{-2}\ \text{s}^{-1}$. During pigment screening, cultures were harvested 7 d after feeding. During the differential light-intensity experiment, triplicate 80 ml subsamples from each of 3 replicate cultures (9 analyses at each light intensity) exposed to each light regime were harvested 10 d after feeding. These are referred to as HA, HB, HC (high-light) and LA, LB, LC (low-light).

For pigment screening, samples of varying volume (usually 3 different volumes), representing different biomasses (Table 1), were prepared from each culture. These were concentrated under gentle vacuum ($<5\ \text{mm Hg}$) onto Whatman GF/F filters. The filters were stored in liquid nitrogen prior to shipment from the culturing facility (Seattle, Washington, USA) to the analytical laboratory (Dania, Florida, USA). The samples were shipped on dry ice and returned to liquid nitrogen within 24 h. Analyses were performed within 2 mo. Samples for cell counting and biovolume estimates were preserved with acid Lugol's iodine solution. Carbon content was determined by using the conversion $0.19\ \text{pg C}\ \mu\text{m}^{-3}$ (Putt & Stoecker 1989). In general, we chose not to attempt mechanical separation of the protozoans from their food as this would increase the risk of damaging the delicate cells. Instead, we chromatographed cultures of food without protozoa and identified the protozoan pigments in experimental (protozoa and food) cultures by process of elimination.

Protoperdinium sp. (740 cells) were individually pipetted from a sample collected by towing a plankton net (26 μm Nitex mesh, 0.50 m mouth diameter) just below the surface at Port Everglades Inlet, Florida (26.3°N , 80.06°W) in November 1987. The sample was analyzed within 5 h of collection.

Copepod eggs were obtained by pipetting 24 adult female *Acartia tonsa* into each of three 2 l bottles containing $0.22\ \mu\text{m}$ filtered seawater. The copepods were placed in a darkened incubator at ambient temperature for 12 h, after which the adults were removed from the containers and the eggs were collected on 35 μm mesh screens. The material on each screen was washed onto a Whatman GF/C filter and frozen in liquid nitrogen pending analysis.

For both sets of measurements, pigments were measured by reverse phase, high performance liquid chromatography (HPLC). Analyses were performed in a cool, dimly lit room with chromatography grade reagents (EM Omnisolve). The approach was described by Kleppel et al. (1988). Each sample on a frozen filter was macerated by hand, in a glass tissue grinder in 90 % aqueous acetone and refrigerated ($5\ ^\circ\text{C}$) for 45 min to permit extraction of pigments. The sample was cleared of particulates by centrifugation (5000 rpm for 5 min). The extract was decanted, an ion pairing agent was added (Solution P; Mantoura & Llewellyn 1983) at 0.3 ml Solution P per ml of extract, and the extract was recentrifuged for 5 min. The extract was then injected onto a Beckman C-8 octyl ultrasphere (5 μm), monomeric reverse phase column (4.5 mm \times 15 cm stainless steel), protected by a stainless steel guard column, packed with ultrasphere octyl (10 μm) material. Pigments were eluted under a 2-solvent (A and B) mobile phase delivered along a continuous gradient from 80 % B : 20 % A

Table 1. Biomass characteristics and pigments of microzooplankton considered in this study. ND: not detected or uncertain identification, not included in analysis; -: no data or not appropriate

Species	Sub-sample	Cell C (pg)	Total C (ng)	Pigment (pg cell ⁻¹)		C : carotenoid ratio
				Canthaxanthin	Astaxanthin	
<i>Gymnodinium</i> sp. (naked, heterotrophic dinoflagellate)	1	0.19	29	ND	ND	-
	2	0.19	288	2.23	ND	129.1
	3	0.19	2888	1.71	ND	111.1
	Mean			1.97		120.1
	SD			0.26		9.0
<i>Strombidium</i> sp. (aloricate, oligotrichous ciliate)	1	2.35	2585	ND	ND	-
	2	2.35	6463	ND	ND	-
	3	2.35	12925	ND	0.05	4.7 × 10 ⁴
	Mean			-	0.05	-
	SD			-	-	-
<i>Strombidinopsis</i> sp. (aloricate, choreotrichous ciliate)	1	8.25	5775	ND	ND	-
	2	8.25	11550	9.60	6.64	859.4
	3	8.25	28182	16.00	5.78	515.6
	Mean			12.80	6.21	687.5
	SD			3.20	0.43	171.9
<i>Favella ehrenbergii</i> (loricate, choreotrichous ciliate)	1	26.00	9880	26.84	ND	968.7
	2	26.00	24700	10.53	2.95	2469.1
	3	26.00	49400	12.11	2.58	2147.0
	Mean			16.49	2.77	1861.6
	SD			7.34	0.19	644.9
<i>Proto-peridinium</i> sp. (thecate, heterotrophic dinoflagellate)	1	-	ND	ND	0.37	-
<i>Acartia tonsa</i> eggs (calanoid copepod)	1	31.00	9300	342.40	ND	90.6
	2	31.00	9300	320.00	ND	96.9
	3	31.00	9300	297.60	ND	104.2
	Mean			320.00		97.2
	SD			0.18		5.6

(v : v) to 100 % B over 10 min. Solvent B was delivered for an additional 10 min. Flow rate rose from 1.0 to 1.9 ml min⁻¹ over the first 10 min, and remained at 1.9 ml min⁻¹ thereafter. Solvent A consisted of an 8 : 1 : 1 solution of water : methanol : Solution P; Solvent B was an 8 : 2 solution of methanol : acetone.

Chromatography was performed with a Waters gradient elution system consisting of paired pumps (model 501), linked to a model 680 gradient controller. Samples were injected into a U6K sampling module and eluting pigments were detected with a model 481, variable wavelength UV/vis detector (usually set at 460 nm). The signal was acquired by an OMS-Tech data processing system run on a 286-microprocessor that logged and plotted the chromatographic data, reported retention times and computed areas under peaks. In one case (*Proto-peridinium* sp.), the data were processed with a Waters model 730 data module, instead of the OMS-Tech system. The Waters data module is not microprocessor interfaced.

To ensure unambiguous detection of animal carotenoids as many as 6 standards were run for each sample analyzed. Standards were usually chromatographed after each sample to ensure the validity of pigment retention times. Many samples were fractionated and chromatographed with and without standards (as internal spikes). Samples of the algae used as food in culture experiments were also chromatographed with and without the addition of astaxanthin and canthaxanthin standards, to identify potential coelution problems. On some occasions astaxanthin seems to have coeluted with diadinoxanthin. Our intensive standardization protocol, however, permitted us to identify astaxanthin in most chromatograms and to detect and reject chromatograms in which co-elution occurred. The chromatographic protocol has subsequently been modified to obviate these problems.

Astaxanthin standards were prepared from live shrimp carapace and muscle tissue. Pigments were extracted in 90 % aqueous acetone and chromato-

graphed as described above. Astaxanthin was collected as it eluted, concentrated to dryness under nitrogen and resuspended in 90 % aqueous acetone. Canthaxanthin standards, in 100 % acetone, were stored in sealed amber vials at -50°C until use. When needed, a vial was opened in a nitrogen atmosphere and the contents were diluted to 90 % aqueous acetone. Fresh astaxanthin was prepared approximately each week. Both pigment standards, once prepared, were stored at -5°C in amber containers, under nitrogen. Standard authenticity was confirmed spectrophotometrically and with published chromatographic data (Table 2).

Quantification of pigments was achieved with empirically-derived relationships between areas under chromatogram peaks and pigment concentrations. The pigment concentrations used in developing these relationships were determined by spectrophotometric analysis of standards with the extinction coefficients given by Herring (1972) for astaxanthin and by Davies (1976) for canthaxanthin.

RESULTS

Pigment screening

Astaxanthin and canthaxanthin were never found in any of the algal food cultures. However, one or both pigments were detected in all of the microzooplankton species examined. *Gymnodinium* sp. contained canthaxanthin (Fig. 1a). Astaxanthin was the predominant carotenoid in the sample of *Protoperdinium* from Port Everglades, Florida (Fig. 1b); a trace of canthaxanthin was also detected. Several photosynthetic accessory pigments were also present in this sample. These would be contributed by phytoplankton cells inadvertently included in the sample during sorting. Both carotenoids, astaxanthin and canthaxanthin, were present in the choreotrichous ciliates *Favella ehrenbergii* and *Strombidinopsis* sp.

(Fig. 1c, d). A small amount of canthaxanthin was detected in the most concentrated sample (12 925 ng C; Table 1) of the oligotrich *Strombidium* sp. (Fig. 1e). Pigments were not detected at lower concentrations in this species. Copepod eggs (Fig. 1f) contained astaxanthin as well as pigments inferred to be xanthophyll esters (peaks 17) that were not detected in the protozoans.

Our ability to detect and quantify pigments in cultured protozoans varied with biomass and taxon (Table 1). Canthaxanthin was detected in approximately 288 ng of *Gymnodinium* sp. carbon-biomass, but was not seen in *Strombidinopsis* sp. below 5800 ng C-biomass. *Strombidium* sp. in our cultures was nearly devoid of carotenoids. The relationship between pigment content and biomass varied with species over more than 2 orders of magnitude. C:carotenoid ratios ranged from ca 100 in copepod eggs and *Gymnodinium* sp., to about 4.7×10^4 in *Strombidium* sp. (Table 1).

Effect of light intensity

The results of the differential light-intensity experiment are shown in Fig. 2. At the high light intensity ($290 \mu\text{E m}^{-2} \text{s}^{-1}$), between-culture differences in cell density, at the time of harvesting, were large. At low light intensity, cell densities were less variable between cultures. In culture HA, the available *Gymnodinium* sp. biomass ($6.6 \text{ cells ml}^{-1} = \text{ca } 100 \text{ ng C per sample}$) was inadequate to provide detectable pigment.

No pigment was detected in cultures maintained at the lower light intensity in this experiment (Fig. 2), though cell densities were sufficient to detect canthaxanthin (Table 1). An additional identical culture (LD) was incubated at $29 \mu\text{E m}^{-2} \text{s}^{-1}$, but harvested 7 (instead of 10) d after feeding, as in the screening studies. *Gymnodinium* sp. density in this culture reached $606 \text{ cells ml}^{-1}$ similar to other low-light cultures. Unlike

Table 2. Spectral and chromatographic characteristics of astaxanthin and canthaxanthin. $E^{1\%}$: published extinction coefficient; -: comparison is made between published absorbance maxima and those observed in the present study; k' : capacity factor, determined from retention time, T_R , and time to elution of the unretained fraction, T_0 , $k' = (T_R - T_0)/T_0$ (Snyder & Kirkland 1979); the number of theoretical plates is determined empirically, and represents the efficiency of the column for resolving a given pigment

Pigment	$E^{1\%}$	Absorbance maxima		Solvent	k'	No. of theoretical plates
		Published	Observed			
Astaxanthin	2099 ^a	480 ^b	479	Acetone	2.47	69 158
Canthaxanthin	2096 ^b	467 ^b	467	Hexane	3.53	70 543

Sources: ^aHerring (1972), ^bDavies (1976)




Fig. 1. Chromatograms of pigments from (a) a cultured heterotrophic dinoflagellate, *Gymnodinium* sp., (b) another heterotrophic dinoflagellate, *Protoperidinium* sp., sorted from plankton tows in a Florida estuary, (c to e) cultures of the ciliates *Favella ehrenbergii*, *Strombidinopsis* sp., and *Strombidium* sp. and (f) the eggs of the copepod *Acartia tonsa*. Note that all chromatograms except (b) were stored to disc using OMS-Tech integrator and processed for presentation using standard spreadsheet software. Chromatogram (b) was traced from hard copy provided by Waters 730 data module. Pigments: 1, unretained fraction; 2, chlorophyll *c* (*c*₁, *c*₂, or both); 3, peridinin-like pigment; 4, peridinin; 5, fucoxanthin; 6, astaxanthin; 7, diadinoxanthin; 8, alloxanthin; 9, diatoxanthin; 10, zeaxanthin; 11, canthaxanthin; 12, unidentified; 13, unidentified; 14, chlorophyll *b*; 15, chlorophyll *a* allomer; 16, chlorophyll *a*; 17, xanthophyll esters; 18, beta-carotene; 19, unidentified; 20, unidentified

the other low-light cultures, canthaxanthin was present in this culture. Cellular canthaxanthin data from the LD culture was combined with canthaxanthin concentrations from *Gymnodinium* sp. in the pigment screening measurements to generate an average, low-light concentration (mean \pm SD) of 1.57 ± 0.60 pg cell⁻¹. This mean was 42 % of the average high-light pigment concentration (3.72 ± 2.61 pg cell⁻¹).

Also shown in Fig. 2 is the concentration of chlorophyll *a* present in the cultures at the time of harvesting. Chlorophyll *a* was used to provide an indication of algal biomass available as food to the heterotrophic *Gymnodinium* sp. Except for culture HA, in which

Gymnodinium sp. cell densities were below the pigment-detection threshold, mean cellular canthaxanthin levels in the dinoflagellate covaried with chlorophyll concentrations in the food supply. Chlorophyll concentrations were higher in cultures exposed to high light intensity than in cultures exposed to low light intensity. Chlorophyll was not detected in those cultures incubated for 10 d at low light intensity (LA, LB, LC), where *Gymnodinium* sp. was abundant but lacked measurable canthaxanthin. Chlorophyll was present in the low-light culture which was harvested after 7 d (LD). Apparently, the food supply was exhausted between Days 7 and 10.




Fig. 2. Mean concentrations of canthaxanthin in *Gymnodinium* sp. exposed to high ($290 \mu\text{E m}^{-2} \text{s}^{-1}$) and low ($29 \mu\text{E m}^{-2} \text{s}^{-1}$) light intensity. Also shown is mean concentration of chlorophyll *a* in each culture. Error bars represent one standard deviation on each side of the mean

DISCUSSION

Pigment screening

It is evident that carotenoid pigments, which are well known in metazoans, also occur in a variety of protozoans. Although astaxanthin and canthaxanthin have been detected in trace amounts in some marine phototrophs (Liaenn-Jensen 1978), these pigments appear to be characteristic of heterotrophic organisms. Astaxanthin is more common than canthaxanthin among metazoans, but both pigments were detected in ciliates and heterotrophic dinoflagellates. Our observation that the heterotrophic dinoflagellates *Gymnodinium* sp. and *Protoperidinium* sp. contain canthaxanthin and astaxanthin are consistent with observations on the occurrence of these pigments in other heterotrophic dinoflagellates [(*Protoperidinium ovatum*: Neveaux & Soyer (1976); *Peridinium foliaceum*: Withers & Haxo (1978); *Noctiluca miliaris*: Balch & Haxo (1984); *Protoperidinium depressum*: Carretto (1985)]. To our knowledge, this is the first report of these pigments in marine ciliates. Xanthophyll esters were detected in copepod eggs but not in the protozoans that we screened for pigments. Further studies will focus on determining whether the presence of xanthophyll esters distinguishes metazoans from protozoans.

Relative to algal cells, most heterotrophs are rather weakly pigmented. Phytoplankton C:carotenoid ratios range from tens to hundreds (Kleppel et al. 1988); ratios for the microzooplankton studied here ranged from hundreds to thousands or higher. The ability to detect pigments in microzooplankton is, naturally, determined by sample size (i.e. amount of biomass in the sample) and by the C:carotenoid ratio. When

cellular pigment levels are relatively high, as in some heterotrophic dinoflagellates and metazoans, such as crustaceans (Hairston 1979, Byron 1982), the dominant pigment can be detected with only a few hundred ng of biomass. Adequate biomass should also be obtained from relatively low numbers of large ciliates (e.g. 200 *Favella ehrenbergii*), despite their characteristically high C:carotenoid ratios. Conversely, small ciliates with high C:carotenoid ratios need to be highly concentrated in order to be detected.

Effects of light intensity

As little as 10% of the incident ambient light is sufficient to cause photo-oxidation of proteins (Krinsky 1971). Carotenoids provide protection against photo-oxidation in a wide variety of organisms (Krinsky 1971, Hairston 1979). However, at the light intensities used in these experiments, the risk of photo-oxidation was minimal (cf. Krinsky & Deneke 1982). Because carotenoids also fulfil a variety of metabolic roles (Cheeseman 1967, Ringleberg 1980, Byron 1982), they might be expected to occur, perhaps at low concentrations, even when not required for photoprotection. In fact, higher average cellular pigment levels occurred in *Gymnodinium* sp. exposed to high light intensity. However, the large variability in pigment content between samples and the occurrence of pigment in cells exposed to low light suggest that factors other than, or in addition to, light intensity influenced the pigment levels in these protozoans.

One such factor may be food availability. Among metazoans, carotenoid biosynthesis is thought to be

mediated by dietary precursors, such as beta-carotene (Katayama et al. 1973, Castillo et al. 1982). In the present study, canthaxanthin was detected in most cultures in which food (i.e. indicated by chlorophyll; Fig. 2) was present (cultures HB, HC, LD). Canthaxanthin was not detected in the relatively dense cultures that were processed after the food supply had been exhausted (cultures LA, LB, LC). The levels of canthaxanthin in *Gymnodinium* sp. covaried with chlorophyll from the algal food supply. It would appear that carotenoid pigment content was dependent upon food availability. In the absence of food, carotenoids were not present in *Gymnodinium* sp. Presumably the pigments were metabolized and could not be replaced.

Carotenoid levels in copepods also seem to be influenced by feeding activity, and carotenoid turnover rates can approach 100 % per day (Hallegraeff et al. 1978, Kleppel et al. 1985). Starvation in cultures LA, LB and LC lasted up to 3 d. This is sufficient time for pigment metabolism to colorless products, given the relatively high weight-specific metabolic rates of protozoans.

CONCLUSION

We have shown that protozoans contain astaxanthin and canthaxanthin, the same carotenoid pigments found in many metazoan zooplankton, but which are rarely reported in phytoplankton. Thus, the carotenoid composition of the protozoans is discernible from that of the phytoplankton. Further work is needed to determine whether carotenoids are ubiquitous among the microzooplankton, as well as to define the magnitude and sources of variability of cellular carbon: carotenoid ratios.

Protozoans and micrometazoans are abundant in the sea and important in marine food webs (Hunter 1981, Porter et al. 1985, Shapiro et al. 1989). Techniques for describing microzooplankton distributions are needed to augment traditional microscopic methods. It is worthwhile exploring the feasibility of using carotenoid pigments as biomarkers for microzooplankton, in much the same way as plant pigments are used to describe phytoplankton distributions. While probably not capable of providing the taxonomic detail of microscopy, carotenoid analysis is relatively rapid and precise. The instrumentation can be used on board ship, and the potential exists for near-real time data acquisition on microzooplankton pigment distributions. It is probable that, in the near future, efforts to describe microzooplankton distributions in the sea will include some form of carotenoid pigment analysis.

Acknowledgements. We are grateful to M. Murrell for assistance, and to R. R. Bidigare for providing canthaxanthin standards. We also appreciate the comments of J. Nelson and E. B. Sherr on the manuscript. The thoughtful comments of 2 anonymous reviewers helped us to avoid several errors in the analysis of the data and generally improved the manuscript. This work was supported by NSF grants OCE 88-22521 and OCE 89-16094 and by the Florida Sea Grant College through NOAA grant NA89 AA-D-SGO53.

LITERATURE CITED

- Balch, W. M., Haxo, F. T. (1984). Spectral properties of *Noctiluca miliaris* Suriray, a heterotrophic dinoflagellate. *J. Plankton Res.* 6: 515-525
- Burkill, P. H., Mantoura, R. F. C., Llewellyn, C. A., Owens, N. J. P. (1987). Microzooplankton grazing and selectivity of phytoplankton in coastal waters. *Mar. Biol.* 93: 581-590
- Byron, E. R. (1982). The adaptive significance of calanoid copepod pigmentation: a comparative experimental analysis. *Ecology* 63: 1871-1886
- Carreto, J. J. (1985). A new keto-carotenoid from the dinoflagellate *Protoperidinium depressum* (Bayley) Balech 1974. *J. Plankton Res.* 7: 421-423
- Castillo, R., Negres-Sadargues, G., Lenel, R. (1982). General survey of carotenoids in Crustacea. In: Britton, G., Goodwin, T. W. (eds.) *Carotenoid chemistry and biochemistry*. Oxford University Press, London, p. 211-224
- Cheeseman, D. F., Lee, W. L., Zegalsky, P. F. (1967). Carotenoids in invertebrates. *Bio. Rev.* 42: 132-160
- Davies, B. H. (1976). Carotenoids. In: Goodwin, T. W. (ed.) *Chemistry and biochemistry of plant pigments*, 2nd edn. Academic Press, London, p. 38-165
- Gifford, D. (1985). Laboratory culture of marine planktonic oligotrichs (Ciliophora, Oligotrichida). *Mar. Ecol. Prog. Ser.* 23: 257-267
- Goodwin, T. W. (ed.) (1976). *Chemistry and biochemistry of plant pigments*, 2nd edn. Academic Press, New York
- Hairston, N. J. Jr (1979). The adaptive significance of color polymorphism in two species of *Diaptomus* (Copepoda). *Limnol. Oceanogr.* 24: 15-37
- Hallegraeff, G. M., Mous, L. J., Veeger, R., Flick, J. G., Ringelberg, J. (1978). A comparative study of carotenoid pigmentation of the zooplankton of Lake Marseveen (Netherlands) and Lac Pavin (Auvergne, France). 2. Diurnal variations in carotenoid content. *Comp. Biochem. Physiol.* 60: 59-62
- Herring, P. J. (1968). The carotenoid pigments of *Daphnia magna* Straus. II. Aspects of pigmentary metabolism. *Comp. Biochem. Physiol.* 24: 205-221
- Herring, P. J. (1972). Depth distribution of carotenoid pigments and lipids of some oceanic animals. I. Mixed zooplankton, copepods and euphausiids. *J. mar. biol. Ass. U.K.* 52: 179-189
- Hunter, J. R. (1981). Feeding ecology and predation of marine fish larvae. In: Lasker, R. (ed.) *Marine fish larvae. Morphology, ecology and relation to fisheries*. Univ. Washington Sea Grant, Seattle, p. 33-77
- Katayama, T., Kunisaki, Y., Shimaya, M., Simpson, K. L., Chichester, C. O. (1973). The biosynthesis of astaxanthin. XIV. The conversion of labeled β -carotene-15- ^{15}C into astaxanthin in the crab *Portunus tuberculatus*. *Comp. Biochem. Physiol.* 24: 205-221
- Kleppel, G. S. (1988). Plant and animal pigments as trophodynamic indicators. In: Soule, D. F., Kleppel, G. S. (eds.)

- Marine organisms as indicators, Springer-Verlag, New York, p. 73–90
- Kleppel, G. S., Frazel, D. W., Pieper, R. E., Holliday, D. V. (1988). Natural diets of zooplankton off southern California. *Mar. Ecol. Prog. Ser.* 49: 231–241
- Kleppel, G. S., Willbanks, L., Pieper, R. E. (1985). Diel variation in body carotenoid content and feeding activity in marine zooplankton assemblages. *J. Plankton Res.* 7: 569–580
- Krinsky, N. I. (1971). Function. In: Isler, O. (ed.) *Carotenoids*. Birkhauser, Basel, p. 669–716
- Krinsky, N. I., Deneke, S. M. (1982). Interaction of oxygen and oxy-radicals with carotenoids. *J. natn. Cancer Inst.* 69: 205–209
- Lessard, E. J. (1991). The trophic role of heterotrophic dinoflagellates in diverse marine environments. *Mar. microb. Fd Webs* 5: 49–58
- Liaaen-Jensen, S. (1979). Carotenoids – a chemosystematic approach. *Pure & appl. Chem.* 51: 661–675
- Mantoura, R. F. C., Llewellyn, C. A. (1983). Rapid determination of algal chlorophyll and carotenoid pigments and their degradation products in natural waters by reverse-phase high performance liquid chromatography. *Analytica chim. Acta* 151: 297–314
- McManus, G. B., Fuhrman, J. A. (1986). Photosynthetic pigments in the ciliate *Laboea strobila* from Long Island Sound, USA. *J. Plankton Res.* 8: 317–327
- Neveux, J., Soyer, M.-O. (1976). Characterisation des pigments et structure fine de *Protoperdinium ovatum* Pouchet (dinoflagellata). *Vie Milieu* 26: 175–199
- Porter, K. G., Sherr, E. B., Sherr, B. F., Pace, M., Sanders, R. W. (1985). Protozoa in planktonic food webs. *J. Protozool.* 32: 409–415
- Putt, M., Stoecker, D. K. (1989). An experimentally determined carbon:volume ratio for marine (oligotrichous) ciliates from estuarine and coast waters. *Limnol. Oceanogr.* 34: 1097–1103
- Ringelberg, J. (1980). Aspects of red pigmentation in zooplankton, especially copepods. In: Kerfoot, W. C. (ed.) *Evolution and ecology of zooplankton communities*. University of New Hampshire Press, Hanover, p. 91–97
- Shapiro, L., Haugen, E. M., Carpenter, E. J. (1989). Occurrence and abundance of green-fluorescing dinoflagellates in surface waters of the Northwest Atlantic and Northeast Pacific Oceans. *J. Phycol.* 25: 189–191
- Sherr, E. B., Sherr, B. F., Paffenhofer, G.-A. (1986). Phagotrophic protozoans as food for metazoans: a missing trophic link in marine pelagic food webs. *Mar. microb. Fd Webs* 1: 61–80
- Smith, R. C., Bidigare, R. R., Prezlin, B. B., Baker, K. S., Brooks, J. M. (1987). Optical characterization of primary productivity across a coastal front. *Mar. Biol.* 96: 575–591
- Snyder, L. R., Kirkland, J. J. (1979). *Introduction to modern liquid chromatography*, 2nd edn. John Wiley, New York
- Stoecker, D. K., Capuzzo, J. M. (1990). Predation on protozoa: its importance to zooplankton. *J. Plankton Res.* 12: 891–908
- Stoecker, D. K., Michaels, A. E., Davis, L. H. (1987). Large proportion of marine planktonic ciliates found to contain functional chloroplasts. *Nature, London* 326: 790–792
- Strain, H. H., Manning, W. M., Harding, G. (1944). Xanthophylls and carotenes of diatoms, brown algae, dinoflagellates and sea anemones. *Biol. Bull. mar. biol. Lab., Woods Hole* 86: 169–191
- Withers, N. W., Cox, E. R., Tomas, R., Haxo, F. T. (1977). Pigments of the dinoflagellate *Peridinium balticum* and its photosynthetic endosymbiont. *J. Phycol.* 13: 354–358
- Withers, N., Haxo, F. T. (1978). Isolation and characterization of carotenoid-rich lipid globules of *Peridinium foliaceum*. *Plant Physiol., Lancaster.* 62: 36–39

This article was presented by E. & B. Sherr, Corvallis, Oregon, USA

*Manuscript first received: March 21, 1992
Revised version accepted: June 4, 1992*