

Effects of ocean acidification on the calcification of otoliths of larval Atlantic cod *Gadus morhua*

R. H. Maneja^{1,*}, A. Y. Frommel¹, A. J. Geffen², A. Folkvord², U. Piatkowski¹,
M. Y. Chang², C. Clemmesen¹

¹GEOMAR, Helmholtz Center for Ocean Research Kiel, Düsternbrooker Weg 20, Kiel 24105, Germany

²Department of Biology, University of Bergen, PO Box 7803, Bergen 5020, Norway

ABSTRACT: The growth and development of the aragonitic CaCO₃ otoliths of teleost fish could be vulnerable to processes resulting from ocean acidification. The potential effects of an increase in atmospheric CO₂ on the calcification of the otoliths were investigated by rearing Atlantic cod *Gadus morhua* L. larvae in 3 pCO₂ concentrations — control (370 µatm), medium (1800 µatm) and high (4200 µatm) — from March to May 2010. Increased otolith growth was observed in 7 to 46 d post hatch (dph) cod larvae at elevated pCO₂ concentrations. The sagittae and lapilli were usually largest in the high pCO₂ treatment followed by the medium and control treatments. The greatest difference in mean otolith surface area (normalized to fish length) was for sagittae at 11 dph, with medium and high treatments being 46 and 43 % larger than the control group, respectively. There was no significant pCO₂ effect on the shape of the otoliths nor were there any trends in the fluctuating asymmetry, defined as the difference between the right and left sides, in relation to the increase in otolith growth from elevated pCO₂.

KEY WORDS: Otolith growth/calcification · Ocean acidification · Atlantic cod larvae · *Gadus morhua* L.

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

The Atlantic cod *Gadus morhua* L. is a northern temperate species, living in a region that is projected to be highly sensitive to future increases in atmospheric CO₂. This is due to higher solubility of CO₂ at low water temperatures, increased air–sea exchange as sea-ice loss increases from global warming, CO₂ remineralization of high organic carbon loads from seasonal primary production and low-alkalinity riverine inputs (Fabry et al. 2009). Upwelling events in coastal waters could enhance the levels of ocean acidification by bringing deeper acidic waters with remineralized CO₂ to the surface, as reported in the bay and inner shelf of the Gulf of Alaska in September 2008 (Fabry et al. 2009) and in Kiel Fjord during summer and autumn in 2008 (Thomsen et al. 2010). The major spawning area of

the Arcto-Norwegian cod, the largest cod population in the world, is located around the Lofoten archipelago in Norway, including Vestfjorden, which is also subject to wind-driven upwelling and downwelling events (Furnes & Sundby 1981). Upwelling caused by winds blowing from the southwest spreads the eggs out toward the central part of Vestfjorden, while downwelling due to winds from the northeast concentrates the older egg stages toward the shore. Wind-driven upwelling and downwelling events along the Norwegian coast also play a role in the dispersal and retention of planktonic organisms and early life stages of fish between the fjords and the Norwegian Coastal Current (Asplin et al. 1999). Accumulation of high-pCO₂ bottom water, which can be upwelled to the surface, along the Norwegian Sea and Barents Sea is also possible. Both regions act as carbon sinks from the first part

*Email: rhmanaja@yahoo.com

of the year until the end of summer (Findlay et al. 2008, Bates et al. 2009).

Teleost fishes produce otolith structures that are mainly composed of aragonitic CaCO_3 (Degens et al. 1969), which may be sensitive to elevated levels of pCO_2 . Munday et al. (2011a) discussed 3 possible consequences of ocean acidification on otolith calcification. First, fish might have more difficulty in precipitating aragonite as the available carbonate ion concentration declines in seawater. Second, otolith hypercalcification might result from increased concentrations of carbonate and bicarbonate ions inside the endolymph due to the maintenance of internal pH by the fish regulatory mechanism with increased ambient pCO_2 . Third, physiological stress from acidic conditions could indirectly influence otolith shape and symmetry. Otolith hypercalcification was reported in white sea bass larvae *Atractoscion nobilis* at 993 and 2558 $\mu\text{atm CO}_2$ (Checkley et al. 2009) and clownfish larvae *Amphiprion percula* at 1721 $\mu\text{atm CO}_2$ (Munday et al. 2011a). In contrast, otolith calcification was not affected in spiny damselfish *Acanthochromis polyacanthus* juveniles grown at 850 $\mu\text{atm CO}_2$ (Munday et al. 2011b).

In the present study, the influence of elevated seawater pCO_2 and the associated shift in carbonate chemistry equilibrium on the calcification of the otoliths of early life stages of Atlantic cod was investigated. The morphology of the otoliths is highly important for the normal acoustic and behavior functions in fish (Gauldie 1988, Lombarte 1992, Aguirre 2003), and abnormal otoliths may ultimately represent an added mortality risk. Understanding the influence of external environmental factors on the growth of the otoliths is also important because otoliths are widely used in fish studies, such as age determination, larval fish ecology and growth studies (Campana 2005).

MATERIALS AND METHODS

Seawater manipulation

The experiment was conducted in land-based mesocosms at the University of Bergen's Espregrennd Marine Station from March to May 2010. Computer-controlled bubbling of CO_2 into the bottom of nine 2650 l experimental tanks was done to achieve 3 pCO_2 levels: control (370 μatm), medium (1800 μatm) and high (4200 μatm), with 3 replicates for each treatment level (see supplementary Table 4 in Frommel et al. 2012). A stable rise of temperature from 5°C

in March to 10°C in May was achieved by placing the experimental tanks inside 2 water baths, which minimized the influence of fluctuations in air temperature. The gradual rise in temperature reflected the natural increase in water temperature of the fjord at 40 m depth, which was the source of the water with mean salinity of 33.3 supplied to the experimental tanks using a flow-through system. For further details, see Frommel et al. (2012).

Larval rearing

Newly fertilized Norwegian coastal cod eggs sourced from Parisvatnet Field Station of the Norwegian Institute of Marine Research were transferred to the floating incubation buckets in the experimental tanks on March 25, 2010. The eggs remained under normal pCO_2 conditions for 3 d, after which the pCO_2 levels were adjusted to the targeted pCO_2 for each treatment. Fifty percent hatching occurred on April 9, 2010, and that date was designated as 0 d post hatch (dph). Redistribution of hatched larvae among replicates within each treatment was carried out on 0 dph to give an equal stocking density of 4 larvae l^{-1} in each tank.

Feeding density of 2000 prey l^{-1} of natural zooplankton was maintained by 24 h filtration of adjacent seawater using a Hydrotech size-selective filter system (Seljeset et al. 2010). The larvae were initially fed with zooplankton with a size fraction of 80–250 μm and this was gradually increased to 350–500 μm .

Handling of animals in the mesocosm experiment was conducted using the animal experimentation permit ID2346 granted by the Animal Welfare Committee as determined under the Norwegian Animal Welfare Act.

Fish sampling and otolith preparation

Fish larvae were sampled 8 times from 7 dph until 46 dph (at 7, 11, 14, 18, 25, 32, 39 and 46 dph). Sampling through the whole water column was done by dropping a PVC pipe with a manual closing mechanism to the bottom of the tanks, obtaining 3 to 12 larvae in each scoop. Point sampling close to the water surface was later implemented when it became harder to catch the larvae using the PVC pipe. Each fish larva was photographed in a thin film of seawater in <30 s and frozen individually in Eppendorf tubes at -80°C . Standard length (SL) of the fish larvae was measured using the calibrated pictures. The

lapilli and sagittae were dissected from the fish larvae using fine needles and mounted on glass slides using Crystalbond™ 509. Pictures were taken of each otolith using an Olympus BX61 compound microscope. Most of the pictures of the otoliths were taken using an oil immersion objective (1000× magnification), while larger otoliths from older larvae were taken at 200 to 600× magnification. The otolith surface area and dimensions were measured from the pictures of the otoliths using ImageJ software (Schneider et al. 2012).

Analysis

The otolith surface area was compared among treatments using analysis of covariance (ANCOVA), with fish SL used as the covariate to account for differences in fish length among fish larvae and dependence of otolith size on fish size (Otterlei et al. 2002). Data from all of the larvae in the 3 replicate tanks of each treatment were treated as one group because there were no significant differences among the replicates within each treatment. Otolith surface area was log-transformed in 2 age groups (32 and 46 dph) to meet the ANCOVA assumption of homogeneity of regression slopes among the 3 pCO₂ treatments, which is required to test for intercept (otolith area) differences. Regression slopes among treatments were not significantly different in the rest of the age group.

In Atlantic cod, an ontogenetic shift in the growth rates of the sagitta and lapillus takes place, with larger lapilli at the earlier larval stages and then a shift to larger sagittae in later stages. Bergstad (1984) reported that the shift in the growth rates of sagitta and lapillus occurred at ~25 dph or 6 mm SL. This transition was monitored in the present study by tracking changes in the ratio of the surface area of the sagitta to that of the lapillus for each fish larva. The sagitta/lapillus surface area ratios were compared among pCO₂ treatments per sampling date using nested ANOVA, with tanks nested in pCO₂ treatments. A non-parametric Kruskal-Wallis test was performed for those sampling dates when the sagitta/lapillus surface area ratios did not satisfy assumptions for normality and homogeneity of variance. The same procedure for nested ANOVA and Kruskal-Wallis test was performed on fish SL data.

Differences in otolith surface area between the right and left sides, referred to as fluctuating asymmetry (Somarakis et al. 1997), were derived for each larva when both right and left otoliths were avail-

able. Higher variability was hypothesized in the high CO₂ treatment. The differences between the 2 sides were normalized by dividing by the mean otolith surface area of the 2 otoliths and then plotted against age of the larvae. The homogeneity of variances of the fluctuating asymmetry with age among the pCO₂ treatments was tested using Hartley *F*-max tests of homogeneity of variances.

The shape of the lapilli and sagittae was also compared using the roundness parameter derived from the otolith pictures using ImageJ. A value of 1 means a perfect circle, while values approaching 0 indicate an increasingly elongated shape (Schneider et al. 2012). A nested ANOVA was used to compare otolith roundness among the pCO₂ treatments with replicate tanks nested in each treatment. The individual roundness value from both the left- and right-side otoliths was included in the analysis.

RESULTS

Increases in the growth of both lapillus and sagittal otoliths were observed in 7 to 46 dph cod larvae (Fig. 1). Larger otolith surface areas were observed in the high, followed by the medium and control treatments. The otolith surface area of larvae in the medium treatment fluctuated between values of the control and high treatments. The greatest difference in mean otolith surface area (ANCOVA adjusted with fish length) was at 11 dph, when the sagittae of larvae from the medium and high treatments were 46 and 43% larger than the control group, respectively.

An ontogenetic shift in the growth rates of the lapilli and sagittae was observed in all larvae (Fig. 2). The lapillus was initially larger than the sagitta, then the growth of the sagitta overtook that of the lapillus. However, the timing of the shift in the sagitta/lapillus surface area ratio occurred earlier, at 32 dph, for larvae from the high treatment compared to the larvae from control and medium treatments. At 11 and 18 dph, the high treatment larvae already showed significantly higher sagitta/lapillus surface area ratios than those in the control and medium treatments even though the shift in growth rates had not yet occurred. At 32 dph, the larvae in the high treatment were >6 mm SL and significantly longer than larvae in the control and medium treatments (Fig. 3). This trend of SL with increasing pCO₂ was seen at 11, 25 and 32 dph.

The left- and right-side otoliths often differed in size (Fig. 4), but there was no bias to indicate directional asymmetry or increase in the magnitude of the




Fig. 1. *Gadus morhua*. Mean (ANCOVA-adjusted) surface area of (a) lapilli and (b) sagittae (both normalized to fish standard length [SL]) from Atlantic cod larvae grown in 3 pCO₂ treatments. Whiskers denote 95% CI. Different letters above bars denote significant differences between pCO₂ treatments based on ANCOVA test with SL as the covariate (at 5% significance level). Numbers inside bars indicate sample sizes; dph: days post-hatch

fluctuation. Variances of the differences in otolith area between right and left sides with age were homogenous amongst pCO₂ treatments (Hartley *F*-max: lapillus 1.28, sagitta 2.33, *p* > 0.05).

The mean roundness values of the lapillus from 7 to 39 dph ranged from 0.91 to 0.94 and did not differ among the pCO₂ treatments (nested ANOVA, *p* > 0.05) (Fig. 5). By 46 dph, the mean roundness values of the lapillus decreased significantly compared to the younger larvae, with otoliths from the medium and high treatments having a significantly more elongated shape than those from the control group (nested ANOVA, *p* < 0.001). For the sagittal otoliths, the roundness values increased from 7 to 32 dph but became more elongated again at 46 dph. There was

no clear pCO₂ treatment effect on the shape of the sagitta except at 25 dph, when more rounded sagittae were associated with increased pCO₂ (nested ANOVA, *p* = 0.016).

DISCUSSION

Our results show that otoliths of Atlantic cod larvae (7 to 46 dph) were larger in elevated pCO₂ conditions. Increase in otolith growth began early in development (7 dph). Several studies have already explored the possibility of enhanced calcification under elevated pCO₂ conditions in marine organisms with active acid/base regulatory mechanisms. High ambient pCO₂ could result in respiratory acidosis where extracellular pCO₂ rises and pH decreases (Melzner et al. 2009). In organisms that regulate acid–base balance, respiratory acidosis is counteracted by elevating the levels of bicarbonate (HCO₃[−]) in the blood and extracellular spaces, which in turn brings the pH closer to the normal level (Marshall & Grosell 2005, Melzner et al. 2009). Cuttlefish minimized the decrease in extracellular blood pH, decreasing by only 0.18 units, when subjected to acute hypercapnia at 0.60 kPa pCO₂ (Gutowska et al. 2010a). This extracellular pH regulation in cuttlefish was achieved through active ion-transport processes, which rapidly increased extra-

cellular HCO₃[−] from 3.38 mM to 9.8 mM within 8 h and stabilized at 10.4 mM after 24 h. Similarly, Atlantic cod (230 to 525 g) showed an increase of 21 mM HCO₃[−] in the blood over 24 h when exposed to 1.10 kPa (Larsen et al. 1997). The increase in HCO₃[−] in the blood of cuttlefish and the partial compensation of extracellular pH could have increased the CaCO₃ saturation state, thereby resulting in elevated calcification rates of the cuttlebone under long-term elevated pCO₂ conditions (Gutowska et al. 2008, 2010b). In teleost fishes, Payan et al. (1998) demonstrated that blood pH in unstarved fish shows a significant positive relationship with the endolymph pH, while Mugiya & Takahashi (1985) reported higher pH and total CO₂ in the endolymph compared




Fig. 2. *Gadus morhua*. Sagitta/lapillus surface area ratio of otoliths from cod larvae grown under 3 pCO₂ concentrations. Whiskers denote 95% CI. Different letters above whiskers denote significant differences between pCO₂ treatments based on nested ANOVA or Kruskal Wallis test, if necessary (at 5% significance level)




Fig. 3. *Gadus morhua*. Standard lengths of cod larvae grown under 3 pCO₂ concentrations. Whiskers denote 95% CI. Different letters above whiskers denote significant differences between pCO₂ treatments based on nested ANOVA or Kruskal Wallis test, if necessary (at 5% significance level)

to plasma. Thus, maintenance of high pH in the blood during hypercapnia could also be translated into higher pH in the endolymph. The hypercalcification observed in larval clownfish *Amphiprion percula* at 1721 µatm pCO₂ (Munday et al. 2011a) and in white seabass larvae at 993 and 2558 µatm pCO₂ (Checkley et al. 2009) was attributed to pH regulation in the otolith endolymph. Payan et al. (1998) provided the evidence that at an endolymph pH of 7.7 to 8, most of the total CO₂ is in HCO₃⁻ form, which acts as the main source for CaCO₃ precipitation onto the otoliths.

Thus, this could be the mechanism causing the increase in otolith growth at elevated pCO₂ levels in our study.

Acid–base regulation by branchial or gill respiration was not yet functional in most of the cod larval stages in our study, especially the younger larvae. Structures that aid in effective gill ventilation, such as the opercular bones, only start to form at 40 to 50 dph (Hunt von Herbing et al. 1996). However, the increase in otolith growth in cod larvae at elevated pCO₂ suggests an already effective acid–base regulation by cutaneous respiration in the early larval stages. Acid–base regulation by raising the HCO₃⁻ concentrations might be just enough for the pH compensation required to counteract acidosis and consequently cause more precipitation of calcium carbonate into the otolith structure. Functional acid–base regulation even at early stages in cod larvae could have been aided by the presence of accessory respiratory structures, such as the pseudo-branch, a structure with a large volume of blood cells and thin epithelial lining positioned in the head of cod larvae, and pigmented hemoglobin in the red blood cells (Mattey et al. 1978, Hunt von Herbing et al. 1996, Marshall & Grosell 2005).

Otoliths are the main organs for detecting acceleration, balance and sound in fish. Increase in the growth of otoliths due to elevated pCO₂ could have consequences on the otolith functions. Huelsenbeck (2010) reported less swimming activity in white seabass larvae grown at pCO₂ 2500 µatm coincident with occurrence of hyper-calcified otoliths. However, cod larvae aged 12 and 27 dph from our experiment showed only subtle changes in their swimming behavior due to increased pCO₂ concentration (Maneja et al. 2012). Changes in otolith growth due to projected and extreme local ocean acidification events might not translate to behavioral changes in fish larvae.




Fig. 4. *Gadus morhua*. Signed difference of otolith surface area ([right–left]/mean) of Atlantic cod larvae grown under 3 pCO₂ concentrations. Signed difference was normalized by dividing by the mean surface area of the 2 sides. Offset in the x-axis was used to display the data per pCO₂ treatment. (a) Lapilli, (b) sagittae

Reports on larvae of tropical marine fishes point to disruptions of neurotransmitter functions in the brain due to elevated levels of HCO₃[−] concentrations in the blood as the main cause of behavioral changes under ocean acidification events (Nilsson et al. 2012). The behavioral changes in fish larvae included damage to olfaction causing loss of homing ability (Munday et al. 2009), loss of detection and avoidance of predators (Dixson et al. 2010, Munday et al. 2010) and loss of predatory reaction to presence of prey (Cripps et al. 2011). The impairment of the neurotransmitter function might also have affected the ability of the larvae to decide between left and right turns (Domenici et al. 2012) and caused a loss of ability to learn anti-predatory

responses (Ferrari et al. 2012). Despite the possible increase in HCO₃[−] concentration in the endolymph as a consequence of pH compensation, implied from increased otolith growth with elevated pCO₂, only subtle changes in the swimming behavior of cod larvae from the same experiment were reported by Maneja et al. (2012). It is important to investigate if the disruption in the neurotransmitter functions in fish larvae due to the effects of ocean acidification (Nilsson et al. 2012) is observable only at the end-points of behavioral cue experiments (e.g. reaction to predator) and not on the specific swimming kinematic variables. For example, a fish larva that is attracted to the smell of a predator might still be swimming normally under acidified seawater conditions. It is worthwhile investigating implications of the subtle behavioral changes in cod larvae with enhanced otolith growth from elevated pCO₂ in behavioral cue experiments, such as reaction to the presence of predators and changes in food densities.

Another interesting result from the present study was the earlier transition of the growth rates of the sagittae and lapilli in the high treatment, which occurred at 32 dph, compared to the medium and control treatments at 39 dph. The differences in the timing of transition in the growth rates of the lapilli and sagittae could have been an indirect effect of the observed differences in the somatic growth rate of the larvae. Bergstad (1984) reported that the lapilli of cod larvae reared at 5 ± 1°C were always larger than the sagittae from hatching until ~6 mm SL or at ~25 dph. Dale (1984) also reported bigger lapilli than sagittae in the early larval stage of cod. In the present study, the mean SL of the larvae from the high treatment had already surpassed 6 mm SL, and was significantly longer than the control and medium treatments, at 32 dph. Cod larvae from the high treatment also had significantly greater dry weights than fish in the control group at 32 dph, as previously reported by Frommel et al. (2012).

Although increased otolith growth was observed in both lapilli and sagittae at elevated pCO₂ levels, there was no systematic pattern of deviation from the normal fluctuating asymmetry and no severe changes in otolith shape. There was a similar and low magnitude of fluctuation of otolith growth between the right and left sides (fluctuating asymmetry) in all treatments. These findings support the earlier report by Munday et al. (2011a) on the maintenance of otolith symmetry and shape despite an increase in otolith growth at 1721 µatm pCO₂. This




Fig. 5. *Gadus morhua*. Mean roundness of (a) lapilli and (b) sagittae from cod larvae grown under 3 pCO₂ concentrations. Whiskers denote 95% CI. Different letters above whiskers denote significant differences between pCO₂ treatments based on nested ANOVA (at 5% significance level). Roundness values range from 0 (highly elongated) to 1 (perfect circle). Numbers inside bars indicate sample sizes

could also mean that changes in the carbonate chemistry inside the endolymph occur at the same magnitude in both left and right sides. The subtle deviations from bilateral symmetry in otolith size have been recommended as an indicator of condition in larval fish, where negative reactions to stress events are recorded as increased levels of fluctuating asymmetry (Somarakis et al. 1997). Lychakov & Rebane (2005) suggested that a specific physicochemical mechanism of the paired otolith growth must exist, allowing teleost fishes to minimize the fluctuation in otolith masses between the left and right otoliths for proper acoustic functionality. Acoustic functionality of the fish, such as sensitivity,

temporal processing and sound localization, were shown through mathematical modeling to be affected by increasing otolith mass asymmetry, particularly those involving large otolith masses.

In summary, the present study showed that under conditions mimicking global and locally enhanced ocean acidification, otolith growth in Atlantic cod larvae increased and size-related transition in lapillus and sagittal growth rates occurred earlier than under normal conditions. However, the magnitude of fluctuating asymmetry did not increase, nor did it lead to directional bias in otolith growth or to severe damage in otolith shape. The increase in otolith calcification due to elevated pCO₂ might be well within the natural variation in otolith sizes maintaining normal functioning.

Acknowledgements. Funding support was provided through the European Community's Seventh Framework Programme (FP7/2007–2013) under the Marie Curie Initial Training Network 'Calcification by Marine Organisms' (CaMarO) and the RTD project 'European Project on Ocean Acidification' (EPOCA, grant agreement N211384). The study was also supported by the project 'Biological Impacts of Ocean ACIDification' (BIOACID), funded by the German Ministry for Education and Research (BMBF). The experiments were conducted at the Norwegian National Mesocosm Centre, Espesgrend, in cooperation with the Department of Biology, University of Bergen.

LITERATURE CITED

- Aguirre WE (2003) Allometric growth of the sulcus in *Cynoscion* spp. (Sciaenidae). *J Fish Biol* 63:1341–1346
- Asplin L, Salvanes AGV, Kristoffersen JB (1999) Nonlocal wind-driven fjord-coast advection and its potential effect on plankton and fish recruitment. *Fish Oceanogr* 8: 255–263
- Bates NR, Mathis JT, Cooper LW (2009) Ocean acidification and biologically induced seasonality of carbonate mineral saturation states in the western Arctic Ocean. *J Geophys Res* 114:C11007, doi:10.1029/2008JC004862
- Bergstad OA (1984) A relationship between the number of growth increments on the otoliths and age of larval and juvenile cod, *Gadus morhua* L. In: Dahl E, Danielssen DS, Moksness E, Solemdal P (eds) The propagation of cod, *Gadus morhua* L. An international symposium.

- Flødevigen rapportser, Part 1. Institute of Marine Research, Arendal, p 251–272
- Campana SE (2005) Otolith science entering the 21st century. *Mar Freshw Res* 56:485–495
- Checkley DM Jr, Dickson AG, Takahashi M, Radich JA, Eisenkolb N, Asch R (2009) Elevated CO₂ enhances otolith growth in young fish. *Science* 324:1683
- Cripps IL, Munday PL, McCormick MI (2011) Ocean acidification affects prey detection by a predatory reef fish. *PLoS ONE* 6:e22736
- Dale T (1984) Embryogenesis and growth of otoliths in the cod (*Gadus morhua* L.). In: Dahl E, Danielssen DS, Moksness E, Solemdal P (eds) The propagation of cod, *Gadus morhua* L. An international symposium. Flødevigen rapportser, Part 1. Institute of Marine Research, Arendal, p 231–250
- Degens ET, Deuser WG, Haedrich RL (1969) Molecular structure and composition of fish otoliths. *Mar Biol* 2: 105–113
- Dixon DL, Munday PL, Jones GP (2010) Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues. *Ecol Lett* 13:68–75
- Domenici P, Allan B, McCormick MI, Munday PL (2012) Elevated carbon dioxide affects behavioral lateralization in a coral reef fish. *Biol Lett* 8(1):78–81
- Fabry VJ, McClintock JB, Mathis JT, Grebmeier JM (2009) Ocean acidification at high latitudes: the bellwether. *Oceanography* (Wash DC) 22:160–171
- Ferrari MCO, Manassa RP, Dixon DL, Munday PL and others (2012) Effects of ocean acidification on learning in coral reef fishes. *PLoS ONE* 7:e31478
- Findlay HS, Tyrrell T, Bellerby RGJ, Merico A, Skjelvan I (2008) Carbon and nutrient mixed layer dynamics in the Norwegian Sea. *Biogeosciences* 5:1395–1410
- Frommel AY, Maneja RH, Lowe D, Malzahn AM and others (2012) Severe tissue damage in Atlantic cod larvae under increasing ocean acidification. *Nature Climate Change* 2:42–46
- Furnes GK, Sundby S (1981) Upwelling and wind induced circulation in Vestfjorden. In: Sætre R, Mork M (eds) The Norwegian Coastal Current. *Proc Norw Coast Curr Symp*. University of Bergen, p 152–177
- Gauldie RW (1988) Function, form and time-keeping properties of fish otoliths. *Comp Biochem Physiol A* 91:395–402
- Gutowska MA, Pörtner HO, Melzner F (2008) Growth and calcification in the cephalopod *Sepia officinalis* under elevated seawater pCO₂. *Mar Ecol Prog Ser* 373:303–309
- Gutowska MA, Melzner F, Langenbuch M, Bock C, Claireaux G, Pörtner HO (2010a) Acid–base regulatory ability of the cephalopod (*Sepia officinalis*) in response to environmental hypercapnia. *J Comp Physiol B* 180:323–335
- Gutowska MA, Melzner F, Pörtner HO, Meier S (2010b) Cuttlebone calcification increases during exposure to elevated seawater pCO₂ in the cephalopod *Sepia officinalis*. *Mar Biol* 157:1653–1663
- Huelsenbeck M (2010) The effects of elevated carbon dioxide concentrations for the larval behavior of white seabass, *Atractoscion nobilis*. MS thesis, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA
- Hunt von Herbing I, Miyake T, Hall BK, Boutilier RG (1996) Ontogeny of feeding and respiration in larval Atlantic cod *Gadus morhua* (Teleostei, Gadiformes): I. Morphology. *J Morphol* 227:15–35
- Larsen BK, Pörtner HO, Jensen FB (1997) Extra- and intracellular acid–base balance and ionic regulation in cod (*Gadus morhua*) during combined and isolated exposures to hypercapnia and copper. *Mar Biol* 128:337–346
- Lombarte A (1992) Changes in otolith area: sensory area ratio with body size and depth. *Environ Biol Fishes* 33:405–410
- Lychakov DV, Rebane YT (2005) Fish otolith mass asymmetry: morphometry and influence on acoustic functionality. *Hear Res* 201:55–69
- Maneja RH, Frommel AY, Browman HI, Clemmesen C and others (2012) The swimming kinematics of larval Atlantic cod, *Gadus morhua* L., are resilient to elevated seawater pCO₂. *Mar Biol* doi:10.1007/s00227-012-2054-y
- Marshall WS, Grosell M (2005) Ion transport and osmoregulation in fish. In: Evans D (ed) *The physiology of fishes*. CRC Press, Boca Raton, FL, p 177–230
- Mattey DL, Moate R, Morgan M (1978) Comparison of 'pseudobranch type' and 'chloride type' cells in the pseudobranch of marine, freshwater and euryhaline teleosts. *J Fish Biol* 13:535–542
- Melzner F, Gutowska MA, Langenbuch M, Dupont S and others (2009) Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences* 6:2313–2331
- Mugiya Y, Takahashi T (1985) Chemical properties of the saccular endolymph in the rainbow trout, *Salmo gairdneri*. *Bull Fac Fish Hokkaido Univ* 36:57–63
- Munday PL, Dixon DL, Donelson JM, Jones GP, Pratchett MS, Devitsina GV, Doving KB (2009) Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proc Natl Acad Sci USA* 106:1848–1852
- Munday PL, Dixon DL, McCormick MI, Meekan M, Ferrari MCO, Chivers DP (2010) Replenishment of fish populations is threatened by ocean acidification. *Proc Natl Acad Sci USA* 107:12930–12934
- Munday PL, Hernaman V, Dixon DL, Thorrold SR (2011a) Effect of ocean acidification on otolith development in larvae of a tropical marine fish. *Biogeosciences* 8: 1631–1641
- Munday PL, Gagliano M, Donelson JM, Dixon DL, Thorrold SR (2011b) Ocean acidification does not affect the early life history development of a tropical marine fish. *Mar Ecol Prog Ser* 423:211–221
- Nilsson GE, Dixon DL, Domenici P, McCormick MI, Sorensen C, Watson SA, Munday PL (2012) Near-future carbon dioxide levels alter fish behavior by interfering with neurotransmitter function. *Nature Climate Change* 2:201–204
- Otterlei E, Folkvord A, Nyhammer G (2002) Temperature dependent otolith growth of larval and early juvenile Atlantic cod (*Gadus morhua*). *ICES J Mar Sci* 59:851–860
- Payan P, Borelli G, Boeuf G, Mayer-Gostan N (1998) Relationship between otolith and somatic growth: consequence of starvation on acid–base balance in plasma and endolymph in the rainbow trout *Oncorhynchus mykiss*. *Fish Physiol Biochem* 19:35–41
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 671–675
- Seljeset O, Vollset K, Folkvord A, Geffen AJ (2010) The role of prey concentration and size range in the growth and survival of larval cod. *Mar Biol Res* 6:251–262
- Somarakis S, Kostikas I, Tsimenides N (1997) Fluctuating asymmetry in the otoliths of larval fish as an indicator of condition: conceptual and methodological aspects. *J Fish Biol* 51(Suppl A):30–38
- Thomsen J, Gutowska MA, Saphörster J, Heinemann A and others (2010) Calcifying invertebrates succeed in a naturally CO₂-rich coastal habitat but are threatened by high levels of future acidification. *Biogeosciences* 7: 3879–3891