

Erratum

Vertical distribution and assemblage structure of leptocephali in the North Equatorial Current region of the western Pacific

Hiroaki Onda, Michael J. Miller, Aigo Takeshige, Yoichi Miyake, Mari Kuroki, Jun Aoyama, Shingo Kimura*

Mar Ecol Prog Ser 575: 119–136, 2017, <https://doi.org/10.3354/meps12198>

*Corresponding author: s-kimura@aori.u-tokyo.ac.jp

- In Table 2 on page 130, the 'N' and '% of total' values for *Gnathophis* at Stn 40 were incorrectly shifted one cell to the left. This has now been corrected in the pdf version online.

The correct table is given below:

Table 2. Number of specimens of leptocephali of 14 taxa in each cluster group and percent contributions (CTB) of taxa to within-group similarity of each cluster group from SIMPER analysis. Only taxa contributing over 4% are shown. No CTB values were calculated for Stn 40 because it did not group with other stations

Family	Taxon	Group A		Group B		Group C		Stn 40	
		N	% CTB	N	% CTB	N	% CTB	N	% of total
Anguillidae	<i>Anguilla japonica</i>	3		7	4.4	2			
	<i>Anguilla marmorata</i>	8		1		41	31.0		
Chlopsidae		34	6.2	13	5.2	3	4.0	3	4.5
Congridae	<i>Ariosoma</i> sp. 4	14	7.3	10	5.4	2		1	1.5
	<i>Ariosoma</i> sp. 5	58	13.2	20	9.5	13	12.1		
	<i>Ariosoma</i> sp. 6	20	5.8	21	8.1	6	7.3	2	3.0
	<i>Ariosoma majus</i>	9		6		3	4.1	8	11.9
	<i>Conger</i>	212	13.5	37	9.3	2			
	<i>Gnathophis</i>	24	10.3	2				31	46.3
Derichthyidae	<i>Derichthys</i>	7	4.1	8	5.3	1			
Muraenidae		50	6.2	17	5.0	5	4.9	10	14.9
Nemichthyidae	<i>Avocettina</i>	2		11	6.1				
	<i>Nemichthys</i>	24	4.9	27	10.0			12	17.9
Serrivomeridae		115	16.3	164	21.6	53	25.6		
Total no. of ind.		584		350		131		67	
No. of tows		6		11		10		1	
Total catch rate (ind. 10 ⁴ m ⁻³)		10.4		3.8		1.6		8.0	