

Corrigendum

Combined effects of temperature and hypoxia shape female brooding behaviors and the early ontogeny of the Chilean kelp crab *Taliepus dentatus*

Simone Baldanzi*, Daniela Storch, Marco Fusi, Nicolas Weidberg, Alexandra Tissot, Sergio A. Navarrete, Miriam Fernández

Mar Ecol Prog Ser 646: 93–107, 2020, <https://doi.org/10.3354/meps13381>

*Corresponding author: simone.baldanzi@uv.cl

-
- An error was discovered in Table S1 of the supplementary material published along with this paper. The Supplement is an environmental data series; wrong dates and incorrect dissolved oxygen values were shown.
 - This error also affects Section 2.2 of the 'Materials and methods' on page 95, where the dates in the series were mentioned. The following sentence was incorrect: 'We analyzed winter and summer mean temperatures and oxygen values for the time period of 1 yr (March 2016 to April 2017). The average daily temperature calculated for austral winter (May to August 2016) and austral summer (November 2016 to February 2017) was 11.02 and 14.62°C, respectively'. This should read: 'We analyzed winter and summer mean temperatures and oxygen values for the time period of 1 yr (October 2015 to October 2016). The average temperature calculated for austral summer (21 December 2015 to 20 March 2016) and austral winter (20 June to 22 September 2016) was 13.85 and 11.97°C, respectively'.
 - We also found a mistake in the beginning of the same paragraph in Section 2.2 on page 95, where the instruments deployed at sea were mentioned: HOBO®Water Temp Pro v2 and HOBO® Dissolved Oxygen Data Loggers were both listed, when in fact only HOBO U26 by Onset (www.onsetcomp.com) was used.
 - Both the amended Supplement and corrected online version of the article are available at www.int-res.com/abstracts/meps/v646/p93-107/
-