

THEME SECTION

Non-breeding distribution and movements of North Atlantic seabirds

Organizers: Hallvard Strøm, Sébastien Descamps, Per Fauchald,
Børge Moe, Morten Ekker

Editors: Kyle H. Elliott, Samantha Patrick, Steve Votier, Rory P. Wilson

Marine Ecology Progress Series Vol. 676, pages 95–276

This Theme Section assembles original research articles based on data collected as part of the SEATRACK project—an international seabird tracking program initiated in 2014 and covering the North East Atlantic (SEATRACK). This program is based on Global Location Sensors (GLS or geolocators) to track individuals during the non-breeding season and better understand this crucial part of the annual life cycle of seabirds. Contributions cover a wide range of topics—from the development of seabird distribution maps for marine spatial planning to the study of the relationships between winter oceanographic conditions, winter movements and demography, and of the individual variability in non-breeding movements.



Photo: Morten Ekker

CONTENTS

**Strøm H, Descamps S, Ekker M, Fauchald P,
Moe B**

INTRODUCTION: REVIEW: Tracking the movements of North Atlantic seabirds: steps towards a better understanding of population dynamics and marine ecosystem conservation 97–116

**Dufour P, Wojczulanis-Jakubas K, Lavergne S,
Renaud J, Jakubas D, Descamps S**

A two-fold increase in migration distance does not have breeding consequences in a long-distance migratory seabird with high flight costs 117–126

**Amélineau F, Merkel B, Tarroux A, Descamps S,
Anker-Nilssen T, Bjørnstad O, Bråthen VS,
Chastel O, Christensen-Dalsgaard S,**

**Danielsen J, Daunt F, Dehnhard N, Ekker M,
Erikstad KE, Ezhov A, Fauchald P, Gavrilo M,
Hallgrímsson GT, Hansen ES, Harris MP,
Helberg M, Helgason HH, Johansen MK,
Jónsson JE, Kolbeinsson Y, Krasnov Y,
Langset M, Lorentsen SH, Lorentzen E,**

**Melnikov MV, Moe B, Newell MA, Olsen B,
Reiertsen T, Systad GH, Thompson P,**

**Thórarinsson TL, Tolmacheva E, Wanless S,
Wojczulanis-Jakubas K, Åström J, Strøm H**

Six pelagic seabird species of the North Atlantic engage in a fly-and-forage strategy during their migratory movements 127–144

**Moe B, Daunt F, Bråthen VS, Barrett RT,
Ballesteros M, Bjørnstad O, Bogdanova MI,
Dehnhard N, Erikstad KE, Follestad A,
Gíslason S, Hallgrímsson GT, Lorentsen SH,
Newell M, Petersen A, Phillips RA,
Ragnarsdóttir SB, Reiertsen TK, Åström J,
Wanless S, Anker-Nilssen T**

Twilight foraging enables European shags to survive the winter across their latitudinal range 145–157

**Dupuis B, Amélineau F, Tarroux A, Bjørnstad O,
Bråthen VS, Danielsen J, Descamps S,
Fauchald P, Hallgrímsson GT, Hansen ES,
Helberg M, Helgason HH, Jónsson JE,
Kolbeinsson Y, Lorentzen E, Thompson P,
Thórarinsson TL, Strøm H**

Light-level geolocators reveal spatial variations in interactions between northern fulmars and fisheries 159–172

**Merkel B, Descamps S, Yoccoz NG, Grémillet D,
Fauchald P, Danielsen J, Daunt F, Erikstad KE,
Ezhov AV, Harris MP, Gavrilo M, Lorentsen SH,
Reiertsen TK, Systad GH, Thórarinsson TL,
Wanless S, Strøm H**

Strong migratory connectivity across meta-populations of sympatric North Atlantic seabirds 173–188

Ezhov AV, Gavrilo MV, Krasnov YV, Bråthen VS, Moe B, Baranskaya AV, Strøm H Transpolar and bi-directional migration strategies of black-legged kittiwakes <i>Rissa tridactyla</i> from a colony in Novaya Zemlya, Barents Sea, Russia	189–203
Léandri-Breton DJ, Tarroux A, Elliott KH, Legagneux P, Angelier F, Blévin P, Bråthen VS, Fauchald P, Goutte A, Jouanneau W, Tartu S, Moe B, Chastel O Long-term tracking of an Arctic-breeding seabird indicates high fidelity to pelagic wintering areas	205–218
Reiertsen TK, Layton-Matthews K, Erikstad KE, Hodges K, Ballesteros M, Anker-Nilssen T, Barrett RT, Benjaminsen S, Bogdanova M, Christensen-Dalsgaard S, Daunt F, Dehnhard N, Harris MP, Langset M, Lorentsen SH, Newell M, Bråthen VS, Støyle-Brinsvor I, Systad GH, Wanless S Inter-population synchrony in adult survival and effects of climate and extreme weather in non-breeding areas of Atlantic puffins.....	219–231
Descamps S, Merkel B, Strøm H, Choquet R, Steen H, Fort J, Gavrilo M, Grémillet D, Jakubas D, Jerstad K, Karnovsky NJ, Krasnov YV, Moe B, Welcker J, Wojczulanis-Jakubas K Sharing wintering grounds does not synchronize annual survival in a high Arctic seabird, the little auk.....	233–242
Albert C, Bråthen VS, Descamps S, Anker-Nilssen T, Cherenkov A, Christensen-Dalsgaard S, Danielsen J, Erikstad KE, Gavrilo M, Hanssen SA, Helgason HH, Jónsson JE, Kolbeinsson Y, Krasnov Y, Langset M, Lorentzen E, Olsen B, Reiertsen TK, Strøm H, Systad GH, Tertitski G, Thompson PM, Thórarinsson TL, Bustamante P, Moe B, Fort J Inter-annual variation in winter distribution affects individual seabird contamination with mercury	243–254
Fauchald P, Tarroux A, Amélineau F, Bråthen VS, Descamps S, Ekker M, Helgason HH, Johansen MK, Merkel B, Moe B, Åström J, Anker-Nilssen T, Bjørnstad O, Chastel O, Christensen-Dalsgaard S, Danielsen J, Daunt F, Dehnhard N, Erikstad KE, Ezhov A, Gavrilo M, Hallgrímsson GT, Hansen ES, Harris M, Helberg M, Jónsson JE, Kolbeinsson Y, Krasnov Y, Langset M, Lorentsen SH, Lorentzen E, Newell M, Olsen B, Reiertsen TK, Systad GH, Thompson P, Thórarinsson TL, Wanless S, Wojczulanis-Jakubas K, Strøm H Year-round distribution of Northeast Atlantic seabird populations: applications for population management and marine spatial planning	255–276