



Corrigendum

Long-term data reveal greater intertidal oyster biomass in predicted suitable habitat

Rachel S. Smith*, Sara Hogan, Kinsey N. Tedford, Bo Lusk, Matthew A. Reidenbach, Max C. N. Castorani

Mar Ecol Prog Ser 683: 221–226, 2022, <https://doi.org/10.3354/meps13949>

*Corresponding author: rsmith218@gmail.com

- A unit conversion error of oyster counts and length to oyster biomass resulted in data presented per 0.25 m² instead of per 1 m² as stated in the y-axis scales of Fig. 2 and the body text. This conversion error does not affect the interpretation of our results.
- This error affects Section 3 'Results' on page 224. The following sentence was incorrect: 'From 2006 to 2019, adult oyster biomass on reefs in suitable habitat was 1.5 greater than adult oyster biomass on reefs in predicted less suitable habitat (Fig. 2A; mean ± SE; 89.03 ± 6.52 g ash free dry weight [AFDW] m⁻², vs. 54.86 ± 10.57 g AFDW m⁻²; F_{1,60} = 8.99, p = 0.004)'. This should read: 'From 2006 to 2019, adult oyster biomass on reefs in suitable habitat was 1.5 greater than adult oyster biomass on reefs in predicted less suitable habitat (Fig. 2A; mean ± SE; 356.11 ± 26.06 g ash free dry weight [AFDW] m⁻², vs. 219.45 ± 42.27 g AFDW m⁻²; F_{1,60} = 8.99, p = 0.004)'.
- In Fig. 2 on page 225, this error affected the y-axis scales and data presented in both panels A and B. Both y-axis scales and data presented in Fig. 2 need to be multiplied by 4 to be correctly presented as per 1 m² as indicated in the y-axis title. The corrected figure is given below.
- This error also affects the values presented in Table S1 in the Supplement in the column 'Adult oyster biomass per m² (mean ± SE)'. The data in this column need to be multiplied by 4 to be correctly presented as per 1 m².
- Both the corrected online version of the article and the corrected Supplement are available at <https://www.int-res.com/abstracts/meps/v683/p221-226/>.




Fig. 2. (A) Adult oyster biomass on suitable and less suitable oyster reefs. Each point represents mean adult oyster biomass (averaged across quadrats) for a given sampling event (*p < 0.05; linear mixed effects model). Boxplots show median (bold lines) and interquartile range (boxes), with outliers greater than 1.5 × IQR (whiskers). AFDW: ash-free dry weight. (B) Mean ± SE adult oyster biomass over the 14 yr sampling period. Dashed lines indicate mean adult oyster biomass, averaged across the 14 yr sampling period, for reefs in predicted suitable (black line) and less suitable (grey line) habitat