

Future potential distribution of the emerging amphibian chytrid fungus under anthropogenic climate change

Dennis Rödder^{1,2}, Jos Kielgast^{3,*}, Stefan Lötters²

¹Zoologisches Forschungsmuseum Alexander Koenig, Adenauerallee 160, 53113 Bonn, Germany

²Department of Biogeography, Trier University, Am Wissenschaftspark 25-27, 54296 Trier, Germany

³Natural History Museum of Denmark, Zoological Museum, Universitetsparken 15, 2100 Copenhagen, Denmark

*Corresponding author. Email: jkielgast@snm.ku.dk

Diseases of Aquatic Organisms: 92: 201–207 (2010)

Supplement

Maxent species distribution models for the A2a and B2a anthropogenic future climate change scenarios for the 2080 CCCMA, CSIRO and HADCM3 models and corresponding standard deviations




Fig. S1. Standard deviation (SD) per grid cell of each of 100 Maxent models projected onto (A) A2a and (B) B2a scenarios




Fig. S2. Species distribution model projection onto the A2a family. Warmer colours indicate higher climatic suitability for *Batrachochytrium dendrobatidis* (*Bd*)




Fig. S3. Species distribution model projection onto the B2a family. Warmer colours indicate higher climatic suitability for *Batrachochytrium dendrobatidis* (*Bd*)