

The following supplement accompanies the article

Food web constraints on larval growth in subtropical coral reef and pelagic fishes

Miram R. Gleiber*, Su Sponaugle, Kelly L. Robinson, Robert K. Cowen

*Corresponding author: mirangleiber@gmail.com

Marine Ecology Progress Series [https://doi: 10.3354/meps13217](https://doi.org/10.3354/meps13217)






Figure S1. Model results of the partial effect of (a) prey background density (*BD*), (b) prey frequency elevated (*FE*), and (c) temperature on recent growth of *Thalassoma bifasciatum*. Predicted recent growth is expressed as a detrended growth index. Fitted lines, 95% confidence intervals (grey shaded areas) and partial residuals (dots) are shown; whiskers on x-axes are field observations for that covariate.








Figure S2. Model results of the partial effect of (a) prey background density (*BD*), (b) prey frequency elevated (*FE*), (c) predator *BD*, (d) predator *FE*, and (e) temperature on recent growth of *Xyrichtys novacula*. Predicted recent growth is expressed as a detrended growth index. Fitted lines, 95% confidence intervals (grey shaded areas) and partial residuals (dots) are shown; whiskers on x-axes are field observations for that covariate.






Figure S3. Model results of the partial effect of (a) prey background density (*BD*), (b) prey frequency elevated (*FE*), and (c) temperature on recent growth of *Katsuwonus pelamis*. Predicted recent growth is expressed as a detrended growth index. Fitted lines, 95% confidence intervals (grey shaded areas) and partial residuals (dots) are shown; whiskers on x-axes are field observations for that covariate.