

SUPPORTING INFORMATION

Table overview

Table S1. List of studies on trophodynamics of larvae from small pelagic fish compiled and analyzed in this review

Table S2. Mean proportion (%) of each prey group for 20 different small pelagic fish larval species.

Table S3. Mean proportion (%) of each prey group for 10 different small pelagic fish larval species and three size classes (mm).

Table S4. Number of studies that used different techniques to get specific information on trophodynamics.

Table S5. Summary of trophic studies on small pelagic fish larvae before and after 2013 in 16 different biogeographic regions worldwide.

Figure overview

Fig. S1. Temporal changes in prey taxonomic resolution reporting level of larval diet studies published.

Table overview**Table S1.** List of studies on trophodynamics of larvae from small pelagic fish compiled and analyzed in this review. For further details on the methods, please see the main text.

Family	Species	System	Method	Reference
Alosidae	<i>Alosa sapidissima</i>	NW Atlantic	Lab. experiment	Johnson et al. (1995)
	<i>Alosa sapidissima</i>	NW Atlantic	Lab. experiment	Riley et al. (2012)
	<i>Brevoortia patronus</i>	Gulf of Mexico	Stomach content	Chen et al. (1992)
	<i>Brevoortia patronus</i>	Gulf of Mexico	Stomach content	Hoover et al. (2022)
	<i>Sardinops melanostictus</i>	NW Pacific	Stomach content	Fukami et al. (1999)
	<i>Sardinops melanostictus</i>	NW Pacific	Stomach content	Yasue et al. (2011)
	<i>Sardinops melanostictus</i>	NW Pacific	Stomach content	Okazaki et al. (2019)
	<i>Sardinops melanostictus</i>	NW Pacific	Stomach content	Matsushita et al. (1988)
	<i>Sardinops melanostictus</i>	NW Pacific	Biomarkers	Yasue et al. (2014)
	<i>Sardinops melanostictus</i>	NW Pacific	Molecular methods	Hirai et al. (2017)
Clupeidae	<i>Sardinops ocellata</i>	Benguela EBUS	Lab. experiment	Brownell (1983)
	<i>Sardinops sagax</i>	Australia	Biomarkers	Uehara et al. (2005)
	<i>Sardinops sagax</i>	California EBUS	Stomach content	Arthur (1976)
	<i>Sardinops sagax</i>	Humboldt EBUS	Stomach content	Muck et al. (1989)
	<i>Sardinops sagax</i>	Humboldt EBUS	Stomach content	Balbontin et al. (1992)
	<i>Sardinops sagax</i>	Humboldt EBUS	Stomach content	Llanos-Rivera et al. (1996)
	<i>Sardinops sagax</i>	Humboldt EBUS	Stomach content	Balbontin et al. (1997)
	<i>Sardinops sagax</i>	Humboldt EBUS	Stomach content	Llanos-Rivera et al. (2004)
	<i>Clupea harengus</i>	Baltic Sea	Stomach content	Schnak (1972)
	<i>Clupea harengus</i>	Baltic Sea	Stomach content	Arula et al. (2012)
Clupeidae	<i>Clupea harengus</i>	Baltic Sea	Lab. experiment	Rosenthal (1969)
	<i>Clupea harengus</i>	Baltic Sea	Lab. experiment	Busch (1996)
	<i>Clupea harengus</i>	Baltic Sea	Lab. experiment	Illing et al. (2015)
	<i>Clupea harengus</i>	Baltic Sea	Lab. experiment	Illing et al. (2018)
	<i>Clupea harengus</i>	Baltic Sea	Biomarkers	Paulsen et al. (2014)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Lebour (1920)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Lebour (1921)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Marshall et al. (1937)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Blaxter (1965)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Schnak (1972)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Munk (1992)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Thiel et al. (1996)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Fox et al. (1999)
Clupeidae	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Figuereido et al. (2005)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Lusseau et al. (2014)

Family	Species	System	Method	Reference
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Denis et al. (2017)
	<i>Clupea harengus</i>	NE Atlantic	Stomach content	Denis et al. (2018)
	<i>Clupea harengus</i>	NE Atlantic	Biomarkers	Denis et al. (2016)
	<i>Clupea harengus</i>	NE Atlantic	Biomarkers	Bils et al. (2022)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Blaxter (1968)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Rosenthal & Hempel (1970)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Geffen (1982)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Checkley (1982)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Checkley (1984)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Kiorboe et al. (1985)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Munk & Kiorboe (1985)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Batty (1987)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Kiorboe et al. (1987)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Pedersen et al. (1990)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	McKenzie & Kiorboe (1995)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Morley & Batty (1996)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Geffen (1996)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Fox et al. (2003)
	<i>Clupea harengus</i>	NE Atlantic	Lab. experiment	Catalan et al. (2011)
	<i>Clupea harengus</i>	NW Atlantic	Stomach content	Cohen & Lough (1983)
	<i>Clupea harengus</i>	NW Atlantic	Stomach content	Courtois & Dodson (1986)
	<i>Clupea harengus</i>	NW Atlantic	Stomach content	Pepin & Penney (1997)
	<i>Clupea harengus</i>	NW Atlantic	Stomach content	Wilson et al. (2018)
	<i>Clupea harengus</i>	NW Atlantic	Stomach content	Pepin (2023)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Stomach content	Pedersen & Fossheim (2008)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Lab. experiment	Folkvord et al. (1997)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Lab. experiment	Folkvord et al. (2000)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Lab. experiment	Utne-Palm & Stiansen (2002)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Lab. experiment	Utne-Palm (2004)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Lab. experiment	Folkvord et al. (2009)
	<i>Clupea harengus</i>	Norwegian/Barents Sea	Lab. experiment	Allan et al. (2021)
	<i>Clupea pallasi</i>	NE Pacific	Stomach content	Hay (1981)
	<i>Clupea pallasi</i>	NE Pacific	Stomach content	Purcell & Grover (1990)
	<i>Clupea pallasi</i>	NE Pacific	Lab. experiment	Boehlert & Yoklavich (1984)
	<i>Clupea pallasi</i>	NE Pacific	Lab. experiment	McGurk (1984)
	<i>Clupea pallasi</i>	NE Pacific	Lab. experiment	Friedenberg et al. (2012)
	<i>Clupea pallasi</i>	California EBUS	Molecular methods	Jungbluth et al. (2021)
	<i>Sardina pilchardus</i>	Canary EBUS	Stomach content & biomarkers	Chicharo et al. (2012)
	<i>Sardina pilchardus</i>	Canary EBUS	Biomarkers	Riveiro et al. (2003)
	<i>Sardina pilchardus</i>	Canary EBUS	Lab. experiment	Caldeira et al. (2014)
Clupeidae	<i>Sardina pilchardus</i>	Canary EBUS	Lab. experiment	Garrido et al. (2016)

Family	Species	System	Method	Reference
	<i>Sardina pilchardus</i>	Canary EBUS	Lab. experiment	Garrido et al. (2021)
	<i>Sardina pilchardus</i>	Canary EBUS	Biomarkers	Becognee et al. (2009)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Stomach content	Rasoanarivo et al. (1991)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Stomach content	Morote et al. (2010)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Stomach content	Costalago et al. (2014)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Biomarkers	Laiz-Carrión et al. (2011)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Biomarkers	Costalago et al. (2012)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Biomarkers	Quintanilla et al. (2015)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Biomarkers	Quintanilla et al. (2020)
	<i>Sardina pilchardus</i>	Mediterranean Sea	Molecular methods	Yebra et al. (2019)
	<i>Sardina pilchardus</i>	NE Atlantic	Stomach content	Lebour (1920)
	<i>Sardina pilchardus</i>	NE Atlantic	Stomach content	Lebour (1921)
	<i>Sardina pilchardus</i>	NE Atlantic	Stomach content	Munuera-Fernandez et al. (2006)
	<i>Sardina pilchardus</i>	NE Atlantic	Stomach content	Voss et al. (2009)
	<i>Sardina pilchardus</i>	NE Atlantic	Lab. experiment	Blaxter (1969)
	<i>Sprattus fuegensis</i>	Patagonia	Stomach content	Contreras et al. (2014)
	<i>Sprattus fuegensis</i>	Patagonia	Stomach content	Bernal et al. (2020)
	<i>Sprattus sprattus</i>	Baltic Sea	Stomach content	Voss et al. (2003)
	<i>Sprattus sprattus</i>	Baltic Sea	Stomach content	Dickmann et al. (2007)
	<i>Sprattus sprattus</i>	NE Atlantic	Stomach content	Lebour (1920)
	<i>Sprattus sprattus</i>	NE Atlantic	Stomach content	Lebour (1921)
	<i>Sprattus sprattus</i>	NE Atlantic	Stomach content	Fortier & Harris (1989)
	<i>Sprattus sprattus</i>	NE Atlantic	Stomach content	Figuereido et al. (2005)
	<i>Sprattus sprattus</i>	NE Atlantic	Stomach content	Figuereido et al. (2007)
	<i>Sprattus sprattus</i>	North Sea	Stomach content	Voss et al. (2009)
	<i>Sprattus sprattus</i>	Baltic Sea	Biomarkers	Peters et al. (2015)
	<i>Sprattus sprattus</i>	Baltic Sea	Lab. experiment	Baumann et al. (2005)
	<i>Sprattus sprattus</i>	Baltic Sea	Lab. experiment	Peck et al. (2015)
	<i>Strangomerina bentincki</i>	Patagonia	Stomach content	Contreras et al. (2014)
	<i>Strangomerina bentincki</i>	Patagonia	Stomach content	Bernal et al. (2020)
	Unidentified clupeidae	SW Pacific	Stomach content	Quah et al. (2022)
Dorosomatidae	<i>Sardinella aurita</i>	Mediterranean Sea	Stomach content	Morote et al. (2008)
	<i>Sardinella brasiliensis</i>	SW Atlantic	Lab. experiment	Del Bianco Rossi-Wongtschow et al. (2003)
	<i>Sardinella zunasi</i>	NW Pacific	Stomach content	Fukami et al. (1999)
Dorosomatidae	<i>Etrumeus teres</i>	Gulf of Mexico	Stomach content	Chen et al. (1992)
	<i>Etrumeus teres</i>	NW Pacific	Stomach content	Yasue et al. (2011)
	<i>Etrumeus teres</i>	NW Pacific	Biomarkers	Yasue et al. (2014)
	<i>Etrumeus teres</i>	NW Pacific	Molecular methods	Hirai et al. (2017)

Family	Species	System	Method	Reference
Ehiravidae	<i>Gilchristella aestuaria</i>	South Africa	Stomach content	Strydom et al. (2014)
	<i>Gilchristella aestuaria</i>	South Africa	Biomarkers	Costalago et al. (2016)
Engraulidae	<i>Anchoa mitchilli</i>	NW Atlantic	Lab. experiment	Houde & Schechter (1980)
	<i>Anchoa mitchilli</i>	NW Atlantic	Lab. experiment	Tucker (1989)
	<i>Anchoa mitchilli</i>	NW Atlantic	Lab. experiment	Cowan & Houde (1990)
	<i>Anchoa mitchilli</i>	NW Atlantic	Lab. experiment	Chesney (2008)
	<i>Anchoa mitchilli</i>	NW Atlantic	Lab. experiment	Munelly et al. (2021)
<i>Anchovia clupeoides</i>	<i>Anchovia clupeoides</i>	SW Atlantic	Stomach content	Lima & Barletta (2016)
	<i>Engraulis anchoita</i>	SW Atlantic	Stomach content	Ciechomski (1963)
	<i>Engraulis anchoita</i>	SW Atlantic	Stomach content	Vasconcellos et al. (1998)
	<i>Engraulis anchoita</i>	SW Atlantic	Stomach content	Sato et al. (2011)
	<i>Engraulis encrasiculus</i>	Benguela EBUS	Lab. experiment	Brownell (1983)
	<i>Engraulis encrasiculus</i>	Canary EBUS	Stomach content	Ferreira & Ré (1993)
	<i>Engraulis encrasiculus</i>	Canary EBUS	Stomach content & biomarkers	Chicharo et al. (2012)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Stomach content	Conway (1998)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Stomach content	Tudela (2002)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Stomach content	Borme et al. (2009)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Stomach content	Catalan et al. (2010)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Stomach content	Morote et al. (2010)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Stomach content	Borme et al. (2013)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Biomarkers	Rossi et al. (2006)
	<i>Engraulis encrasiculus</i>	Mediterranean Sea	Biomarkers	Costalago et al. (2011)
<i>Engraulis japonicus</i>	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Uotani (1985)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Matsushita et al. (1988)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Fukami et al. (1999)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Islam et al. (2006)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Islam & Tanaka (2009)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Yasue et al. (2010)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Yasue et al. (2011)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Choi et al. (2018)
	<i>Engraulis japonicus</i>	NW Pacific	Stomach content	Okazaki et al. (2019)
	<i>Engraulis japonicus</i>	NW Pacific	Biomarkers	Lindsay et al. (1998)
Engraulidae	<i>Engraulis japonicus</i>	NW Pacific	Biomarkers	Yasue et al. (2014)
	<i>Engraulis mordax</i>	California EBUS	Stomach content	Arthur (1976)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Lasker et al. (1970)

Family	Species	System	Method	Reference
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Theilacker & McMaster (1971)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Hunter (1972)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Hunter & Thomas (1974)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Lasker (1975)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Hunter (1976a)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Hunter (1976b)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Scura and Jerde (1977)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Moffat (1991)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Schmitt (1986)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Theilacker (1987)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Huntley (1989)
	<i>Engraulis mordax</i>	California EBUS	Lab. experiment	Rodriguez-Murillo et al. (1989)
	<i>Engraulis mordax</i>	California EBUS	Biomarkers	Hakanson (1993)
	<i>Engraulis mordax</i>	California EBUS	Molecular methods	Ohman et al. (1991)
	<i>Engraulis ringens</i>	Humboldt EBUS	Biomarkers	Castro et al. (2020)
	<i>Engraulis ringens</i>	Humboldt EBUS	Stomach content	Muck et al. (1989)
	<i>Engraulis ringens</i>	Humboldt EBUS	Stomach content	Balbontin et al. (1992)
	<i>Engraulis ringens</i>	Humboldt EBUS	Stomach content	Llanos-Rivera et al. (1996)
	<i>Engraulis ringens</i>	Humboldt EBUS	Stomach content	Balbontin et al. (1997)
	<i>Engraulis ringens</i>	Humboldt EBUS	Stomach content	Llanos-Rivera et al. (2004)
	<i>Engraulis ringens</i>	Humboldt EBUS	Stomach content	Yanez-Rubio et al. (2011)
	<i>Engraulis ringens</i>	Patagonia	Stomach content	Landaeta et al. (2014)
	<i>Engraulis ringens</i>	Patagonia	Stomach content	Bernal et al. (2020)
	<i>Unidentified Engraulidae</i>	SW Pacific	Stomach content	Quah et al. (2022)

References

- Allan BJM, Browman HI, Shema S, Skiftesvik A-B, Folkvord A, Durif CMF, Kjesbu OS (2022) Increasing temperature and prey availability affect the growth and swimming kinematics of Atlantic herring (*Clupea harengus*) larvae. J Plankton Res 44:401–413.
- Arthur DK (1976) Food and feeding of larvae of three fishes occurring in the California Current, *Sardinops sagax*, *Engraulis mordax*, and *Trachurus symmetricus*. Fish Bull 74:517–530.
- Arula T, Kotta J, Lankov A, Simm M, Pölmé S (2012) Diet composition and feeding activity of larval spring-spawning herring: Importance of environmental variability. J Sea Res 68:33–40.
- Balbontín F (1992) Growth and survival of Chilean sardine *Sardinops sagax*, larvae reared at different densities of food. Revista de Biología Marina Y Oceanografía 27:101–112.
- Balbontín F, Llanos A, Valenzuela V (1997) Trophic overlap and feeding incidence in fish larvae from central Chile. Revista Chilena de Historia Natural 70:381–390.
- Batty RS (1987) Effect of light intensity on activity and food-searching of larval herring, *Clupea harengus*: a laboratory study. Mar Biol 94:323–327.
- Baumann H, Peck MA, Herrmann J-P (2005) Short-term decoupling of otolith and somatic growth induced by food level changes in postlarval Baltic sprat, *Sprattus sprattus*. Mar Freshwater Res 56:539.
- Bécognée P, Moyano M, Almeida C, Rodríguez JM, Fraile-Nuez E, Hernández-Guerra A, Hernández-León S (2009) Mesoscale distribution of clupeoid larvae in an upwelling filament trapped by a quasi-permanent cyclonic eddy off Northwest Africa. Deep Sea Research Part I: Oceanographic Research Papers 56:330–343.
- Bernal A, Castro LR, Soto S, Cubillos LA (2020) Ichthyoplankton distribution and feeding habits of fish larvae at the inshore zone of northern Patagonia, Chile. Mar Biodivers 50:56.
- Bils F, Aberle N, van Damme CJG, Peck MA, Moyano M (2022) Role of protozooplankton in the diet of North Sea autumn spawning herring (*Clupea harengus*) larvae. Mar Biol 169:90.
- Blaxter JHS (1969) Experimental rearing of pilchard larvae, *Sardina pilchardus*. J Mar Biol Ass 49:557–575.
- Blaxter JHS (1965) The feeding of herring larvae and their ecology in relation to feeding. CalCOFI Report X 79-88.
- Blaxter JHS (1968) Visual Thresholds and Spectral Sensitivity of Herring Larvae. J Exp Biol 48:39–53.

- Boehlert GW, Yoklavich MM (1984) Carbon assimilation as a function of ingestion rate in larval pacific herring, *Clupea harengus pallasi* Valenciennes. J Exp Mar Biol Ecol 79:251–262.
- Borme D, Tirelli V, Brandt S, Fonda Umani S, Arneri E (2009) Diet of *Engraulis encrasicolus* in the northern Adriatic Sea (Mediterranean): ontogenetic changes and feeding selectivity. Mar Ecol Prog Ser 392:193–209.
- Borme D, Tirelli V, Palomera I (2013) Feeding habits of European pilchard late larvae in a nursery area in the Adriatic Sea. J Sea Res 78:8–17.
- Brownell CL (1983) Laboratory rearing of Cape anchovy *Engraulis capensis* and South African pilchard *Sardinops ocellata* through metamorphosis. African J Mar Sci 1:181–188.
- Busch A (1996) Transition from endogenous to exogenous nutrition: larval size parameters determining the start of external feeding and size of prey ingested by Ruegen spring herring *Clupea harengus*. Mar Ecol Prog Ser 130:39–46.
- Caldeira C, Santos A, Ré P, Peck M, Saiz E, Garrido S (2014) Effects of prey concentration on ingestion rates of European sardine *Sardina pilchardus* larvae in the laboratory. Mar Ecol Prog Ser 517:217–228.
- Castro LR, González V, Claramunt G, Barrientos P, Soto S (2020) Stable isotopes ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) seasonal changes in particulate organic matter and in different life stages of anchoveta (*Engraulis ringens*) in response to local and large scale oceanographic variations in north and central Chile. Progress in Oceanography 186:102342.
- Catalán IA, Folkvord A, Palomera I, Quílez-Badía G, Kallianoti F, Tselepidis A, Kallianotis A (2010) Growth and feeding patterns of European anchovy (*Engraulis encrasicolus*) early life stages in the Aegean Sea (NE Mediterranean). Estuarine, Coastal and Shelf Science 86:299–312.
- Catalán IA, Vollset KW, Morales-Nin B, Folkvord A (2011) The effect of temperature gradients and stomach fullness on the vertical distribution of larval herring in experimental columns. J Exp Mar Biol Ecol 404:26–32.
- Checkley DM (1984) Relation of growth to ingestion for larvae of Atlantic herring *Clupea harengus* and other fish. Mar Ecol Prog Ser 18:215–224.
- Checkley DM (1982) Selective feeding by Atlantic herring (*Clupea harengus*) larvae on zooplankton in natural assemblages. Mar Ecol Prog Series 9:245–253.
- Chen W, Govoni J, Warlen SM (1992) Comparison of feeding and growth of larval round herring *Etrumeus teres* and gulf menhaden *Brevoortia patronus*. Fish Bull 90:183–189.
- Chesney EJ (2008) Foraging behavior of bay anchovy larvae, *Anchoa mitchilli*. J Exp Mar Biol Ecol 362:117–124.
- Chícharo MA, Amaral A, Faria A, Morais P, Mendes C, Piló D, Ben-Hamadou R, Chícharo L (2012) Are tidal lagoons ecologically relevant to larval recruitment of small pelagic fish? An approach using nutritional condition and growth rate. Estuarine, Coastal and Shelf Science 112:265–279.
- Choi HC, Youn SH, Huh S-H, Park JM (2018) Diet Composition and Feeding Habits of Two Engraulid Fish Larvae (*Engraulis japonicus* and *Coilia nasus*) in the Nakdong River Estuary, Korea. J Coastal Res 85:346–350.
- Ciechomski JD (1963) Investigations of food and feeding habits of larvae and juveniles of the Argentine anchovy *Engraulis anchoita*. CalCOFI Report XI.
- Cohen RE, Lough RG (1983) Prey field of larval herring *Clupea harengus* on a continental shelf spawning area. Mar Ecol Prog Series 10:211–222.
- Contreras T, Castro LR, Montecinos S, Gonzalez HE, Soto S, Muñoz MI, Palma S (2014) Environmental conditions, early life stages distributions and larval feeding of patagonian sprat *Sprattus fuegensis* and common sardine *Strangomeria bentincki* in fjords and channels of the northern Chilean patagonia. Progress in Oceanography 129:136–148.
- Conway D, Coombs S, Smith C (1998) Feeding of anchovy *Engraulis encrasicolus* larvae in the northwestern Adriatic Sea in response to changing hydrobiological conditions. Mar Ecol Prog Ser 175:35–49.
- Costalago D, Miranda N, Strydom N, Perissinotto R (2016) Using stable isotope analysis to study the diet of *Gilchristella aestuaria* larvae: preliminary insights into the foodwebs of six South African estuaries. African J Aquatic Sci 41:389–398.
- Costalago D, Navarro J, Álvarez-Calleja I, Palomera I (2012) Ontogenetic and seasonal changes in the feeding habits and trophic levels of two small pelagic fish species. Mar Ecol Prog Ser 460:169–181.
- Costalago D, Palomera I (2014) Feeding of European pilchard (*Sardina pilchardus*) in the northwestern Mediterranean: from late larvae to adults. Sci Mar 78:41–54.
- Costalago D, Tecchio S, Palomera I, Álvarez-Calleja I, Ospina-Álvarez A, Raicevich S (2011) Ecological understanding for fishery management: Condition and growth of anchovy late larvae during different seasons in the Northwestern Mediterranean. Estuarine, Coastal and Shelf Science 93:350–358.
- Courtois R, Dodson JJ (1986) Régime alimentaire et principaux facteurs influençant l'alimentation des larves de capelan (*Mallotus villosus*), d'éperlan (*Osmerus mordax*) et de hareng (*Clupea harengus harengus*) dans un estuaire partiellement mélangé. Can. J. Fish. Aquat. Sci: 43.
- Cowan JJH, Houde ED (1990) Growth and survival of bay anchovy *Anchoa mitchilli* larvae in mesocosm enclosures. Mar Ecol Prog Ser 68:47–57.
- Del Bianco Rossi-Wongtschow CL, Clemmesen C, Ueberschär B, Ferraz Dias J (2003) Larval condition and growth of *Sardinella brasiliensis* (Steindachner, 1879): preliminary results from laboratory studies. Sci Mar 67:13–23.
- Denis J, Mahe K, Tavernier E, Monchy S, Vincent D, Vallet C, Marchal P, Antajan E, Caboche J, Lefebvre V, Cordier R, Loots C (2017) Ontogenetic changes in the larval condition of Downs herring: use of a multi-index approach at an individual scale. Mar Biol 164:154.
- Denis J, Vallet C, Courcot L, Lefebvre V, Caboche J, Antajan E, Marchal P, Loots C (2016) Feeding strategy of Downs herring larvae (*Clupea harengus* L.) in the English Channel and North Sea. J Sea Res 115:33–46.
- Denis J, Vincent D, Antajan E, Vallet C, Mestre J, Lefebvre V, Caboche J, Cordier R, Marchal P, Loots C (2018) Gut fluorescence technique to quantify pigment feeding in Downs herring larvae. Mar Ecol Prog Ser 607:129–142.
- Dickmann M, Möllmann C, Voss R (2007) Feeding ecology of Central Baltic sprat *Sprattus sprattus* larvae in relation to zooplankton dynamics: implications for survival. Mar Ecol Prog Ser 342:277–289.
- Ferreira PL, Re P (1993) Feeding of larval anchovy, *Engraulis encrasicolus* (L.), in the Mira Estuary (Portugal). Portugaliaae zoologica 2:1–37.
- de Figueiredo GM, Nash RDM, Montagnes DJS (2007) Do protozoa contribute significantly to the diet of larval fish in the Irish Sea? J Mar Biol Ass 87:843–850.
- de Figueiredo GM, Nash RDM, Montagnes DJS (2005) The role of the generally unrecognised microprey source as food for larval fish in the Irish Sea. Mar Biol 148:395–404.
- Folkvord A, Blom G, Johannessen A, Moksness E (2000) Growth-dependent age estimation in herring (*Clupea harengus* L.) larvae. Fish Res 46:91–103.

- Folkvord A, Høie H, Johannessen A, Solbakken T (2009) Effects of prey concentration, light regime, and parental origin on growth and survival of herring larvae under controlled experimental conditions. ICES J Mar Sci 66:1702–1709.
- Folkvord A, Rukan K, Johannessen A, Moksness E (1997) Early life history of herring larvae in contrasting feeding environments determined by otolith microstructure analysis. J Fish Biol 51:250–263.
- Fortier L, Harris RP (1989) Optimal foraging and density-dependent competition in marine fish larvae. Mar Ecol Prog Ser 51:19–33.
- Fox CJ, Folkvord A, Geffen AJ (2003) Otolith micro-increment formation in herring *Clupea harengus* larvae in relation to growth rate. 264:83–94.
- Fox CJ, Harrop R, Wimpenny A (1999) Feeding ecology of herring (*Clupea harengus*) larvae in the turbid Blackwater Estuary. Mar Biol 134:353–365.
- Friedenberg LE, Bollens SM, Rollwagen-Bollens G (2012) Feeding dynamics of larval Pacific herring (*Clupea pallasi*) on natural prey assemblages: the importance of protists: Larval herring feeding on protists. Fish Oceanogr 21:95–108.
- Fukami K, Watanabe A, Fujita S, Yamaoka K, Nishijima T (1999) Predation on naked protozoan microzooplankton by fish larvae. Mar Ecol Prog Ser 185.
- Garrido S, Cristóvão A, Caldeira C, Ben-Hamadou R, Baylina N, Batista H, Saiz E, Peck M, Ré P, Santos A (2016) Effect of temperature on the growth, survival, development and foraging behaviour of *Sardina pilchardus* larvae. Mar Ecol Prog Ser 559:131–145.
- Garrido S, Ferreira S, Soares C, Meneses I, Baylina N, Batista H, Teodósio MA, Piecho-Santos AM, Ré P (2021) Effect of food availability on the growth and age determination of European sardine (*Sardina pilchardus* Walbaum, 1792) larvae. J Mar Biol Ass 101:609–619.
- Garrido S, Saiz E, Peters J, Ré P, Alvarez P, Cotano U, Herrero DL, de Murguía AM, Irigoien X (2012) Effect of food type and concentration on growth and fatty acid composition of early larvae of the anchovy (*Engraulis encrasicolus*) reared under laboratory conditions. J Exp Mar Biol Ecol 434–435:16–24.
- Geffen A (1996) Effect of Experimental Manipulation of Feeding Conditions on the Population Structure of Larval Cod (*Gadus morhua*) and Herring (*Clupea harengus*). Mar Freshwater Res 47:291.
- Geffen AJ (1982) Otolith ring deposition in relation to growth rate in herring (*Clupea harengus*) and turbot (*Scophthalmus maximus*) larvae. Mar Biol 71:317–326.
- Håkanson JL (1993) Nutritional condition and growth rate of anchovy larvae (*Engraulis mordax*) in the California Current: two contrasting years. Mar Biol 115:309–316.
- Hay DE (1981) Effects of capture and fixation on gut contents and body size of Pacific herring larvae. Rapport et proces-verbaux des réunions, Conseil permanent international pour l'exploration de la mer 178:395–400.
- Hirai J, Hidaka K, Nagai S, Ichikawa T (2017) Molecular-based diet analysis of the early post-larvae of Japanese sardine *Sardinops melanostictus* and Pacific round herring *Etrumeus teres*. Mar Ecol Prog Ser 564:99–113.
- Hoover A, Chiaverano L, Deary AL, Hernandez F (2022) Variation in larval Gulf Menhaden diet, growth and condition during an atypical winter freshwater-discharge event in the northern Gulf of Mexico. Estuarine, Coastal and Shelf Science 265:107692.
- Houde ED, Schekter RC (1980) Feeding by marine fish larvae: developmental and functional responses. Environ Biol Fish 5:315–334.
- Hunter JR (1976) Culture and growth of Northern anchovy, *Engraulis mordax*, larvae. Fish Bull 74:81–88.
- Hunter JR (1972) Swimming and feeding behaviour of larval anchovy, *Engraulis mordax*. Fishery Bulletin 70:821–838.
- Hunter JR, Thomas GL (1974) Effect of Prey Distribution and Density on the Searching and Feeding Behaviour of Larval Anchovy *Engraulis mordax* Girard. In: *The Early Life History of Fish*. Blaxter JHS (ed) Springer Berlin Heidelberg, Berlin, Heidelberg, p 559–574.
- Huntley ME (1989) Larval feeding of northern anchovy, *Engraulis mordax*, on dinoflagellates: implications for year-class strength. Sci Mar 53:239–245.
- Illing B, Moyano M, Berg J, Hufnagl M, Peck MA (2018) Behavioral and physiological responses to prey match-mismatch in larval herring. Estuarine, Coastal and Shelf Science 201:82–94.
- Illing B, Moyano M, Niemax J, Peck MA (2015) Direct Effects of Microalgae and Protists on Herring (*Clupea harengus*) Yolk Sac Larvae. PLoS ONE 10:e0129344.
- Intxausti L, Villate F, Motos L, Uriarte I, Iriarte A (2017) Diet variability in European anchovy: a comparative analysis between larval populations of the inner Bay of Biscay and the NW Mediterranean. Hydrobiologia 790:49–65.
- Islam MdS, Hibino M, Tanaka M (2006) Distribution and diets of larval and juvenile fishes: Influence of salinity gradient and turbidity maximum in a temperate estuary in upper Ariake Bay, Japan. Estuarine, Coastal and Shelf Science 68:62–74.
- Islam MdS, Tanaka M (2009) Diet and prey selection in larval and juvenile Japanese anchovy *Engraulis japonicus* in Ariake Bay, Japan. Aquat Ecol 43:549–558.
- Johnson JH, Dropkin DS (1995) Effects of prey density and short term food deprivation on the growth and survival of American shad larvae. Journal of Fish Biology 46:872–879.
- Jungbluth MJ, Burns J, Grimaldo L, Slaughter A, Katla A, Kimmerer W (2021) Feeding habits and novel prey of larval fishes in the northern San Francisco Estuary. Environmental DNA 3:1059–1080.
- Kiørboe T, Munk P, Richards K (1987) Respiration and growth of larval herring *Clupea harengus*: relation between specific dynamic action and growth efficiency. Mar Ecol Prog Ser 40:1–10.
- Kiørboe T, Munk P, Støtrup JG (1985) First feeding by larval herring *Clupea harengus* L. Dana:95–107.
- Laiz-Carrión R, Quintanilla JM, Mercado JM, García A (2011) Combined study of daily growth variability and nitrogen-carbon isotopic signature analysis of schooling *Sardina pilchardus* larvae. Journal of Fish Biology 79:896–914.
- Landaeta MF, Ochoa-Muñoz MJ, Bustos CA (2014) Feeding success and selectivity of larval anchoveta *Engraulis ringens* in a fjord-type inlet from northern Patagonia (Southeast Pacific). Rev biol mar oceanogr 49:461–475.
- Lasker R (1975) Field criteria for survival of anchovy larvae: the relation between inshore chlorophyll maximum layers and successful first feeding. Fish Bull 73:453–462.
- Lasker R, Feder HM, Theilacker GH, May RC (1970) Feeding, growth, and survival of *Engraulis mordax* larvae reared in the laboratory. Mar Biol 5:345–353.
- Lebour MV (1920) The Food of Young Fish No. III (1919). J Mar Biol Ass UK 12:261–324.
- Lebour MV (1921) The food of young clupeoids. J Mar Biol Ass UK 12:458–467.
- Lima ARA, Barletta M (2016) Lunar influence on prey availability, diet shifts and niche overlap between Engraulidae larvae in tropical mangrove creeks. J Fish Biol 89:2133–2152.

- Lindsay DJ, Minagawa M, Mitani I, Kawaguchi K (1998) Trophic Shift in the Japanese Anchovy *Engraulis japonicus* in its Early Life History Stages as Detected by Stable Isotope Ratios in Sagami Bay, Central Japan. *Fisheries science* 64:403–410.
- Llanos-Rivera A, Herrera G, Bernal P (1996) Análisis del tamaño de las presas en la dieta de las larvas de cuatro clupeiformes en un área costera de Chile central. *Scientia Marina* 60:435–442.
- Llanos-Rivera A, Herrera GA, Bernal P (2004) Food size selectivity and diet overlap in larvae of Clupeiform species from central Chile. *Cahiers De Biologie Marine* 45:1–8.
- Lusseau SM, Gallego A, Rasmussen J, Hatfield EMC, Heath M (2014) North Sea herring (*Clupea harengus L.*) recruitment failure may be indicative of poor feeding success. *ICES J Mar Sci* 71:2026–2041.
- MacKenzie BR, Kiørboe T (1995) Encounter rates and swimming behavior of pause-travel and cruise larval fish predators in calm and turbulent laboratory environments. *Limnol Oceanogr* 40:1278–1289.
- Marshall SM, Nicholls AG, Orr AP (1937) On The Growth and Feeding of the Larval and Post-Larval Stages of the Clyde Herring. *J Mar Biol Ass* 22:245–267.
- Matsushita K, Shimizu M, Nose Y (1988) Food density and rate of feeding larvae of anchovy and sardine in patchy distribution. *Nippon Suisan Gakkaishi* 54:401–411.
- McGurk MD (1984) Effects of delayed feeding and temperature on the age of irreversible starvation and on the rates of growth and mortality of Pacific herring larvae. *Mar Biol* 84:13–26.
- Moffat NM (1981) Survival and growth of Northern anchovy larvae on low zooplankton densities by the presence of a Chlorella bloom. *Rapp P-V Reun Cons Int Explor Mer* 178:475–480.
- Morley SA, Batty RS (1996) The effects of temperature on "S-Strike" feeding of larval herring, *Clupea Harengus L.* *Marine and Freshwater Behaviour and Physiology* 28:123–136.
- Morote E, Olivari MP, Villate F, Uriarte I (2010) A comparison of anchovy (*Engraulis encrasicolus*) and sardine (*Sardina pilchardus*) larvae feeding in the Northwest Mediterranean: influence of prey availability and ontogeny. *ICES J Mar Sci* 67:897–908.
- Morote E, Olivari MP, Villate F, Uriarte I (2008) Diet of round sardinella, *Sardinella aurita*, larvae in relation to plankton availability in the NW Mediterranean. *J Plankton Res* 30:807–816.
- Muck PE, Rojas de Mendiola B, Antonietti EM (1989) Comparative studies on feeding in larval anchoveta (*Engraulis ringens*) and sardine (*Sardinops sagax*). In: *ICLARM Conference Proceedings*. p 86–96
- Munk P (1992) Foraging behaviour and prey size spectra of larval herring *Clupea harengus*. *Mar Ecol Prog Ser* 80:149–158.
- Munk P, Kiørboe T (1985) Feeding behaviour and swimming activity of larval herring (*Clupea harengus*) in relation to density of copepod nauplii. *Mar Ecol Prog Ser* 24:15–21.
- Munnely RT, Windecker CC, Reeves DB, Rieucau G, Portier RJ, Chesney EJ (2021) Effects of short-duration oil exposure on bay anchovy (*Anchoa mitchilli*) embryos and larvae: mortality, malformation, and foraging. *Aquatic Toxicology* 237:105904.
- Munuera-Fernandez I, Gonzalez-Quiros R (2006) Analysis of feeding of *Sardina pilchardus* (Walbaum, 1792) larval stages in the central Cantabrian Sea. *Sci Mar* 70:131–139.
- Ohman MD, Theilacker GH, Kaupp SE (1991) Immunochemical Detection of Predation on Ciliate Protists by Larvae of the Northern Anchovy (*Engraulis mordax*). *The Biological Bulletin* 181:500–504.
- Okazaki Y, Tadokoro K, Kubota H, Kamimura Y, Hidaka K (2019) Dietary overlap and optimal prey environments of larval and juvenile sardine and anchovy in the mixed water region of the western North Pacific. *Mar Ecol Prog Ser* 630:149–160.
- Paulsen M, Clemmesen C, Malzahn AM (2014) Essential fatty acid (docosahexaenoic acid, DHA) availability affects growth of larval herring in the field. *Mar Biol* 161:239–244.
- Peck MA, Ewest B, Herrmann J-P, Al-Misned FA, Mortuza MG, Stäcker S, Temming A (2015) Relationships between feeding, growth and swimming activity of European sprat (*Sprattus sprattus L.*) post-larvae in the laboratory. *Environ Biol Fish* 98:1117–1127.
- Pedersen BH, Ugelstad I, Hjelmeland K (1990) Effects of a transitory, low food supply in the early life of larval herring (*Clupea harengus*) on mortality, growth and digestive capacity. *Mar Biol* 107:61–66.
- Pedersen T, Fossheim M (2008) Diet of 0-group stages of capelin (*Mallotus villosus*), herring (*Clupea harengus*) and cod (*Gadus morhua*) during spring and summer in the Barents Sea. *Mar Biol* 153:1037–1046.
- Pepin P, Penney RW (1997) Patterns of prey size and taxonomic composition in larval fish: are there general size-dependent models? *J Fish Biol* 51:84–100.
- Peters J, Diekmann R, Clemmesen C, Hagen W (2015) Lipids as a proxy for larval starvation and feeding condition in small pelagic fish: a field approach on match-mismatch effects on Baltic sprat. *Mar Ecol Prog Ser* 531:277–292.
- Purcell JE, Grover JJ (1990) Predation and food limitation as causes of mortality in larval herring at a spawning ground in British Columbia. *Mar Ecol Prog Ser* 59:55–61.
- Quah WC, Chew LL, Chong VC, Chu C, Teoh CY, Ooi AL (2022) Does structural change in the zooplankton community affect larval fish feeding in anthropogenically disturbed tropical waters? *Environ Biol Fish* 105:55–76.
- Quintanilla J, Laiz-Carrión R, Uriarte A, García A (2015) Influence of trophic pathways on daily growth patterns of western Mediterranean anchovy *Engraulis encrasicolus* larvae. *Mar Ecol Prog Ser* 531:263–275.
- Quintanilla JM, Laiz-Carrión R, García A, Quintanilla LF, Cortés D, Gómez-Jakobsen F, Yebra L, Salles S, Putzeys S, León P, Mercado JM (2020) Early life trophodynamic influence on daily growth patterns of the Alboran Sea sardine (*Sardina pilchardus*) from two distinct nursery habitats (bays of málaga and almería) in the western Mediterranean Sea. *Mar Environment Res* 162:105195.
- Rasoanarivo R, Folack J, Champalbert G, Becker B (1991) Relations entre les communautés phytoplanctoniques et l'alimentation des larves de *Sardina pilchardus* Walb. dans le golfe de Fos (Méditerranée occidentale): influence de la lumière sur l'activitéalimentaire des larves. *J Exp Mar Biol Ecol* 151:83–92.
- Riley KL, Binion SM, Overton AS (2012) Estimating the Food Requirements and Prey Size Spectra of Larval American Shad. *Marine and Coastal Fisheries* 4:228–238.
- Riveiro I, Guisande C, Franco C, Lago de Lanzós A, Maneiro I, Vergara A (2003) Egg and larval amino acid composition as indicators of niche resource partitioning in pelagic fish species. *Mar Ecol Prog Ser* 260:255–262.
- Rodriguez-Murillo JA, Carrillo-Barrios-Gomez E, Chagoya-Guzman A (1989) Growth and survival of larvae of the northern anchovy *Engraulis mordax* (Pisces: Engraulidae) reared under starvation conditions. *Revista de Biología Tropical* 37:169–179.
- Rosenthal H (1969) Untersuchungen über das Beutefangverhalten bei larven des herings *Clupea harengus*. *Marine Biology* 3:208–221.
- Rosenthal H, Hempel G (1970) Experimental studies in feeding and food requirements of herring larvae (*Clupea harengus L.*). In: *Marine food chains*. Steele JH (ed) University of California Press, Berkeley, p 344–364

- Rossi S, Sabatés A, Latasa M, Reyes E (2006) Lipid biomarkers and trophic linkages between phytoplankton, zooplankton and anchovy (*Engraulis encrasicolus*) larvae in the NW Mediterranean. *J Plankton Res* 28:551–562.
- Sato NE, Hernandez D, Viñas MD (2011) Feeding habits of *Engraulis anchoita* (Hubbs & Marini, 1935) larvae in coastal waters off Buenos Aires Province, Argentina. *Latin American Journal of Aquatic Research* 39:16–24.
- Schmitt PD (1986) Prey size selectivity and feeding rate of larvae of Northern anchovy, *Engraulis mordax* Girard. CalCOFI Report XXVII.
- Schnack D (1972) Studies on the food ecology of herring larvae. *Ber dt wiss Kommn Meeresforsch* 22:273–343.
- Scura ED, Jerde CW (1977) Various species of phytoplankton as food for larval Northern anchovy, *Engraulis mordax*, and relative nutritional value of the dinoflagellates *Gymnodinium splendens* and *Gonyaulax polyedra*. *Fish Bull* 75:577–583.
- Strydom N, Sutherland K, Wooldridge T (2014) Diet and prey selection in late-stage larvae of five species of fish in a temperate estuarine nursery. *African J Mar Sci* 36:85–98.
- Theilacker GH (1987) Feeding ecology and growth energetics of larval Northern anchovy, *Engraulis mordax*. *Fish Bull* 85:213–228.
- Theilacker GH, McMaster MF (1971) Mass culture of the rotifer *Brachionus plicatilis* and its evaluation as a food for larval anchovies. *Mar Biol* 10:183–188.
- Thiel R, Mehner T, Kopcke B, Kafemann R (1996) Diet Niche Relationships among Early Life Stages of Fish in German Estuaries. *Mar Freshwater Res* 47:123.
- Tucker JW (1989) Energy utilization in Bay anchovy, *Anchoa mitchilli*, and Black sea bass, *Centropristes striata striata*, eggs and larvae. *Fish Bull* 78:279–298.
- Tudela S, Palomera I, Quílez G (2002) Feeding of anchovy *Engraulis encrasicolus* larvae in the north-west Mediterranean. *J Mar Biol Ass* 82:349–350.
- Uehara S, Syahailatua A, Suthers IM (2005) Recent growth rate of larval pilchards *Sardinops sagax* in relation to their stable isotope composition, in an upwelling zone of the East Australian Current. *Mar Freshwater Res* 56:549.
- Uotani I (1985a) The relation between feeding mode and feeding habit of the anchovy larvae. *Bulletin of the Japanese Society for the Science of Fish* 51:1057–1065.
- Uotani I (1985b) The relation between feeding mode and feeding habit of the anchovy larvae. *Nippon Suisan Gakkaishi* 51:1057–1065.
- Utne-Palm AC (2004) Effects of larvae ontogeny, turbidity, and turbulence on prey attack rate and swimming activity of Atlantic herring larvae. *J Exp Mar Biol Ecol* 310:147–161.
- Utne-Palm AC, Stiansen JE (2002) Effect of larval ontogeny, turbulence and light on prey attack rate and swimming activity in herring larvae. *J Exp Mar Biol Ecol* 268:147–170.
- Vasconcellos MC, Freire KF, Castello JP (1998) Distribution patterns and feeding success of anchovy, *Engraulis anchoita*, larvae off southern Brazil. *Sci Mar* 62:385–392.
- Voss R, Dickmann M, Schmidt JO (2009) Feeding ecology of sprat (*Sprattus sprattus* L.) and sardine (*Sardina pilchardus* W.) larvae in the German Bight, North Sea. *Oceanologia* 51:117–138.
- Voss R, Köster FW, Dickmann M (2003) Comparing the feeding habits of co-occurring sprat (*Sprattus sprattus*) and cod (*Gadus morhua*) larvae in the Bornholm Basin, Baltic Sea. *Fisheries Research* 63:97–111.
- Wilson CJ, Murphy HM, Bourne C, Pepin P, Robert D (2018) Feeding ecology of autumn-spawned Atlantic herring (*Clupea harengus*) larvae in Trinity Bay, Newfoundland: Is recruitment linked to main prey availability? *J Plankton Res* 40:255–268.
- Yañez-Rubio A, Llanos-Rivera A, Castro LR, Claramunt G, Herrera L (2011) Variations in type, width, volume and carbon content of anchoveta *Engraulis ringens* food items during the early larval stages. *J Mar Biol Ass* 91:1207–1213.
- Yasue N, Doiuchi R, Takasuka A (2014) Trophodynamic similarities of three sympatric clupeoid species throughout their life histories in the Kii Channel as revealed by stable isotope approach. *ICES J Mar Sci* 71:44–55.
- Yasue N, Doiuchi R, Yoshimoto Y, Takeuchi T (2010) Diet of late larval Japanese anchovy *Engraulis japonicus* in the Kii Channel, Japan. *Fish Sci* 76:63–73.
- Yasue N, Takasuka A, Shirakihara K (2011) Interspecific comparisons of growth and diet among late larvae of three co-occurring clupeoid species in the Kii Channel, Japan. *Mar Biol* 158:1709–1720.
- Yebra L, Hernández de Rojas A, Valcárcel-Pérez N, Castro M, García-Gómez C, Cortés D, Mercado J, Laiz-Carrión R, García A, Gómez-Jakobsen F, Uriarte A, Rodríguez J, Quintanilla J (2019) Molecular identification of the diet of *Sardina pilchardus* larvae in the SW Mediterranean Sea. *Mar Ecol Prog Ser* 617–618:41–52.

Table S2. Mean proportion (%) of each prey group for 20 different small pelagic fish larval species obtained from 48 studies reporting 60 diets. For each species the number (n) of studies reporting diet are indicated.

Small pelagic fish larval species	<i>C. harengus</i>	<i>E. mordax</i>	<i>S. pilchardus</i>	<i>E. encrasicolus</i>	<i>S. sprattus</i>	<i>E. japonicus</i>	<i>S. sagax</i>	<i>E. ringens</i>	<i>C. pallasi</i>	<i>S. melanostictus</i>	<i>E. anchoita</i>	<i>E. teres</i>	<i>S. aurita</i>	<i>B. patronus</i>	<i>S. bentincki</i>	<i>S. fuengensis</i>	<i>G. aestuaria</i>	<i>A. clupeoides</i>	<i>C. edentulus</i>
Prey group / n	13	1	5	8	3	6	4	5	1	2	2	2	1	2	1	1	1	1	
Copepod	17.3	62.0	35.3	34.5	53.8	11.0	65.3	37.9	76.5	4.4	28.1	30.7	48.2	21.3	92.3	72.0	-	25.4	15.8
Calanoid	41.9	7.4	34.1	7.9	34.4	49.6	-	-	-	41.0	37.7	35.7	-	55.7	-	-	25.8	45.9	40.0
Cyclopoid	7.7	-	2.9	10.8	-	11.9	-	-	-	40.1	5.9	14.0	-	6.9	-	-	17.2	-	-
Harpacticoid	-	-	2.1	9.8	-	-	-	-	-	4.1	-	-	2.6	-	-	-	-	-	-
Cladocera	-	-	3.1	8.1	2.9	13.3	-	-	-	4.6	-	23.3	3.8	-	-	-	-	-	22.5
Euphausiids	-	-	-	-	-	-	-	-	-	3.4	-	-	-	-	-	-	-	-	-
Cirripedia	-	-	-	-	-	-	-	12.0	-	-	-	-	-	-	-	-	-	-	-
Ostracoda	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5.6	-	-
Crustacea	6.1	-	-	-	-	-	-	3.1	-	6.3	-	3.5	2.4	-	7.6	15.1	-	24.7	12.1
Bivalvia	2.4	-	2.1	-	-	-	-	-	2.4	-	-	-	-	-	-	-	-	-	3.8
Gastropod	6.5	-	-	-	-	-	-	-	-	-	3.2	2.6	-	-	-	-	-	-	-
Polychaeta	-	-	-	-	-	-	-	-	-	-	-	4.3	-	-	-	-	-	-	-
Appendicularia	-	-	-	-	-	-	-	-	-	-	-	3.6	-	-	-	-	-	-	-
Tintinnid	-	2.5	9.3	9.8	-	3.8	-	-	-	2.0	2.6	-	2.1	-	-	-	-	-	-
Amphioxus	-	-	-	-	-	-	-	-	-	-	-	3.0	-	-	-	-	-	-	-
Insecta	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	50.8	-	-	-
Invertebrate	-	-	-	-	-	-	23.6	13.7	-	-	5.8	-	5.7	-	-	-	-	-	-
Protozoa	-	-	-	-	-	-	-	-	6.4	-	-	-	-	-	-	-	-	-	-
Rotifer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ciliophora	-	3.0	-	-	-	-	-	2.2	-	-	-	-	-	-	-	-	-	-	-
Algae	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4.9
Diatom	-	-	-	-	-	-	-	-	14.4	-	-	-	-	-	-	-	-	-	-
Dinoflagellates	-	4.8	3.1	3.6	-	-	3.0	11.6	-	-	-	-	-	-	-	-	-	-	-
Phytoplankton	2.24	3.3	-	-	-	-	-	3.2	-	-	-	-	3.2	-	-	12.8	-	-	-
Other	15.8	16.8	7.9	15.3	8.9	10.3	8.2	16.2	0.2	8.2	14.2	4.3	9.5	2.0	0.1	-	0.6	4.0	0.9

Table S3. Mean proportion (%) of each prey group for 10 different small pelagic fish larval species and three size classes (mm) obtained from 23 studies. If available, the numerical percentage (%N) was used. For each species the number (n) of studies reporting ontogenetic diet are indicated.

	Cladocera	Copepodite	Copepoda	Copepoda egg	Crustacea	Invertebrate egg	Copepoda Nauplii	Other	Microplankton	Copepoda Postnauplii	Polychaeta	Tunicata
<i>C. harengus</i> (n=3)												
<10 mm	-	8.8	0.0	-	0.0	-	87.3	-	-	-	-	-
7-20 mm	-	21.0	13.5	-	5.9	-	50.5	-	-	-	-	-
20-53 mm	-	54.41	19.8	-	13.1	-	5.2	-	-	-	-	-
<i>E. mordax</i> (n=1)												
<4.5 mm	-	-	-	15.3	-	-	42.9	28.5	13.2	-	-	-
5-6.5 mm	-	-	-	15.3	-	-	42.9	28.5	13.2	-	-	-
7-9 mm	-	-	-	7.2	-	-	79.4	8.1	5.4	-	-	-
<i>S. pilchardus</i> (n=1)												
4.5-7.6 mm	0.0	-	6.6	-	-	74.0	19.5	-	-	-	-	-
7.7-13 mm	7.5	-	5.0	-	-	82.0	5.5	-	-	-	-	-
13.1-22.1 mm	100.0	-	0.0	-	-	0.0	0.0	-	-	-	-	-
<i>E. encrasiculus</i> (n=3)												
2-4 mm	-	0.0	-	14.2	-	-	45.3	34.8	5.8	0.0	-	0.0
3-9.5 mm	-	26.0	-	8.4	-	-	31.3	19.3	0.0	15.1	-	0.0
8-16 mm	-	32.3	-	11.1	-	-	11.3	20.9	0.0	18.5	-	5.8
<i>S. sprattus</i> (n=2)												
4-10 mm	0.0	8.6	-	0.0	-	-	70.5	20.9	-	-	-	-
9-18 mm	0.0	70.9	-	5.7	-	-	18.5	4.9	-	-	-	-
>18 mm	24.3	70.9	-	0.0	-	-	0.0	4.9	-	-	-	-
<i>E. japonicas</i> (n=2)												
2.5-7.5 mm	14.0	-	43.8	-	-	-	0.0	11.4	30.9	-	-	0.0
8-25 mm	0.0	-	76.9	-	-	-	0.0	11.6	0.0	-	-	11.6
25-40 mm	5.0	-	75.3	-	-	-	5.3	6.0	0.0	-	-	8.3
<i>S. sagax</i> (n=3)												
3-9.5 mm	-	0.0	0.0	44.5	0.0	-	40.5	15.0	-	-	-	-
8-14 mm	-	14.3	22.5	10.8	12.5	-	32.4	7.5	-	-	-	-
10-25 mm	-	36.0	0.0	19.0	0.0	-	38.7	6.3	-	-	-	-
<i>E. ringens</i> (n=3)												
3-8 mm	-	0.0	0.0	31.4	0.0	-	28.1	12.2	28.3	-	-	-
6-14 mm	-	8.3	6.3	15.8	5.4	-	33.6	13.8	11.4	-	-	-
9-20 mm	-	34.5	0.0	7.8	0.0	-	34.7	8.8	5.3	-	-	-
<i>E. anchoita</i> (n=1)												
3-5 mm	0.0	0.0	0.0	0.0	0.0	7.2	47.8	0.0	-	-	-	-
12-22 mm	17.6	10.4	60.8	5.6	0.0	5.6	0.0	0.0	-	-	-	-
22-42 mm	27.8	0.0	45.6	45.1	12.7	6.0	0.0	7.9	-	-	-	-
<i>S. aurita</i> (n=2)												
4.5-7.6 mm	0.0	-	-	-	-	-	0.0	0.0	-	0.0	0.0	-
3.9-8 mm	18.7	-	-	-	-	-	33.3	39.8	-	8.1	0.0	-
8-14.7 mm	27.9	-	-	-	-	-	25.6	16.3	-	23.3	7.0	-

Table S4. Summary of the number of studies* that use each technique to get specific

Information obtained	Techniques				
	Biomarkers	Experiments	Gut analysis	SIA	Molecular
Diet composition_type of prey	11		91		5
Diet composition_size of prey			37		
Diel feeding intensity			27		1
Selectivity	5	13	25	7	2
Foraging behaviour			18		
Ingestion rates			17		
Digestion			4		
Growth and/or survival - Food level			40		
Growth and/or survival - temperature			4		
$\delta^{15}\text{N}$ - Trophic level					12
Trophic position					2
Food source					13
Resource partitioning	4				5

* Numbers reflect the combination of studies and species, therefore if one study reports two species, it counts as two studies.
information on trophodynamics.

Table S5. Summary of trophic studies on small pelagic fish larvae before and after 2013 in 16 different biogeographic regions worldwide.

Region	1920-2012 (# studies)	2013-2022 (# studies)
NE Atlantic	29	7
California EBUS	16	1
Mediterranean Sea	11	5
NW Atlantic	9	2
NW Pacific	8	3
Baltic Sea	7	5
Norwegian/Barents Sea	6	1
Humboldt EBUS	6	1
Alaska/Bering Sea	5	0
Canary EBUS	4	2
SW Atlantic	4	1
Australia	1	2
Gulf of Mexico	1	1
Benguela EBUS	1	0
Patagonia	0	3
South Africa	0	2

Figure overview

Figure S1 - Temporal changes in prey taxonomic resolution reporting level of larval diet studies published. Each circle corresponds to a publishing period (1920-1990; 1991-2012; 2013-2022), and each colour shows the proportion (%) of papers reporting the diet at; species (dark blue), genus (light blue), order (grey), qualitative (yellow) or kingdom (orange) level.

