

Functional diversity of the Barents Sea fish community

Magnus Aune Wiedmann^{1*}, Michaela Aschan¹, Grégoire Certain², Andrey Dolgov³,
Michael Greenacre^{1,4}, Edda Johannesen⁵, Benjamin Planque², Raul Primicerio⁶

¹Norwegian College of Fishery Science, University of Tromsø (UiT), 9037 Tromsø, Norway

²Institute of Marine Research (IMR), PO Box 6404, 9294 Tromsø, Norway

³Knipovich Polar Research Institute of Marine Fisheries and Oceanography (PINRO), 6 Knipovich Street, 183038 Murmansk, Russian

⁴Universitat Pompeu Fabra and Barcelona Graduate School of Economics, Ramon Trias Fargas 25–27, 08005 Barcelona, Spain

⁵Institute of Marine Research (IMR), PO Box 1870, Nordnes, 5817 Bergen, Norway

⁶Department of Marine and Arctic Biology, University of Tromsø (UiT), 9037 Tromsø, Norway

*Corresponding author: magnus.a.wiedmann@uit.no

Marine Ecology Progress Series 495: 205–218 (2014)

Supplement 2. Literature cited for functional trait matrix presented in Supplement 1

- Albert OT (1993) Distribution, population structure and diet of silvery pout (*Gadiculus argenteus thori* J. Schmidt), poor cod (*Trisopterus minutus minutus* (L.)), four-bearded rockling (*Rhinonemus cimbrius* (L.)), and Vahl's eelpout (*Lycodes vahlii gracilis* Reinhardt) in the Norwegian Deep. *Sarsia* 78:141–154
- Albert OT, Nilssen EM, Stene A, Gundersen AC, Nedreaas KH (2001) Maturity classes and spawning behavior of Greenland halibut (*Reinhardtius hippoglossoides*). *Fish Res* 51:217–228
- Andriyashev AP (1954) Fishes of northern seas of the U.S.S.R. Academy of Science Press, Moscow-Leningrad. Translated from Russian by the Israel Program for Scientific Translations, Jerusalem 1964
- Atkinson EG, Percy JA (1992) Diet comparison among demersal marine fish from the Canadian Arctic. *Polar Biol* 11:567–573
- Bagenal TB (1963) The fecundity of witches in the Firth of Clyde. *J Mar Biol Assoc UK* 43:401–407
- Beare DJ, Moore PG (1997) The contribution of amphipoda to the diet of certain inshore fish species in the Kames Bay, Millport. *J Mar Biol Assoc UK* 77:907–910
- Berestovskiy YG (1995) Feeding habitats and strategy of American plaice, *Hippoglossoides platessoides limandoides*, in the Barents and Norwegian Seas. *J Ichthyol* 35:40–54
- Bjelland O, Bergstad OA, Skjæraasen JE, Meland K (2000) Trophic ecology of deep-water fishes associated with the continental slope of the eastern Norwegian Sea. *Sarsia* 85:101–117
- Breder CM, Rosen DE (1966) Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey
- Burnett J, Ross MR, Clark SH (1992) Several biological aspects of the witch flounder (*Glyptocephalus cynoglossus* (L.)) in the Gulf of Maine-Georges Bank region. *J Northwest Atl Fish Sci* 12:15–25

- Calis E, Jackson EH, Nolan CP, Jeal F (2005) Preliminary age and growth estimates of the rabbitfish, *Chimara monstrosa*, with implications for future resource management. *J Northwest Atl Fish Sci* 35:15–26
- Casey JM, Myers RA (1998) Near extinction of a large, widely distributed fish. *Science* 281:690–692
- Christiansen SJ, Fevolden SE, Karamuschko OV, Karamuschko LI (1998) Maternal output in polar fish reproduction. In: Di Prisco G, Pisano E, Clarke A (eds) *Fishes of Antarctica. A biological overview. Part 1 Ecology*. Springer Verlag, Milano, p 41–51
- Clarke MW, Kelly CJ, Connolly PL, Molloy JP (2003) A life history approach to the assessment and management of deep-water fisheries in the Northeast Atlantic. *J Northwest Atl Fish Sci* 31:401–411
- Coad BW, Power G (1973) Observations on the ecology and phenotypic variation of the threespine stickleback, *Gasterosteus aculeatus* L., 1758, and the blackspotted stickleback, *G. wheatlandi* Putnam, 1867 (Osteichthyes: Gasterosteidae) in Amory Cove, Quebec. *Can Field Nat* 87:113–122
- Dalpadado P, Ellertsen B, Melle W, Dommasnes A (2000) Food and feeding conditions of Norwegian spring-spawning herring (*Clupea harengus*) through its feeding migrations. *ICES J Mar Sci* 57:843–857
- Dolgov AV (2005) Feeding and food consumption by the Barents Sea skates. *J Northwest Atl Fish Sci* 35:495–503
- Dolgov AV (2009) Trophic structure of the Barents Sea fish assemblage with special reference to the cod stock recoverability. *Prog Oceanogr* 81:165–173
- Drevetnyak KV, Kluev AI (2005) On fecundity of *Sebastes viviparus* from the North East Arctic. *ICES CM* 2005/Q:31
- Ebert DA, Bizzarro JJ (2007) Standardized diet compositions and trophic levels of skates (Chondrichthyes: Rajiformes: Rajoidei). *Environ Biol Fishes* 80:221–237
- Eliassen JE, Falk-Petersen IB (1985) Reproductive biology of the roughhead grenadier (*Macrourus berglax* Lacepède) (Pisces, Gadiformes) from the continental slope of northern Norway. *Sarsia* 70:59–67
- Eliassen JE, Jobling M (1985) Food of roughhead grenadier, *Macrourus berglax*, Lacepede in north Norwegian waters. *J Fish Biol* 26:367–376
- Ennis GP (1970) Age, growth and sexual maturity of the shorthorn sculpin, *Myoxocephalus scorpius*, in Newfoundland waters. *J Fish Res Board Can* 27:2155–2158
- Erzini K (1994) An empirical study of variability in length-at-age of marine fishes. *J Appl Ichthyology* 10:17–41
- Falk-Petersen IB, Frivoll V, Gulliksen B, Haug T, Vader W (1988) Age/size relations and food of two snailfishes, *Liparis gibbus* and *Careproctus reinhardii* (Teleostei, Liparididae) from Spitsbergen coastal waters. *Polar Biol* 8:353–358
- Ferry-Graham LA, Drazen JC, Franklin V (2007) Laboratory observations of reproduction in the deep-water zoarcids *Lycodes cortezianus* and *Lycodapus mandibularis* (Teleostei: Zoarcidae). *Pac Sci* 61:129–139
- Froese R, Pauly D (eds) (2011) FishBase. Available at www.fishbase.org
- Gjøsæter J (1973) Preliminary results of Norwegian polar cod investigation 1970-1972. *ICES CM* F:8
- Gjøsæter J (1981a) Life history and ecology of *Maurolicus muelleri* (Gonostomatidae) in Norwegian waters. *Fiskeridir Skr Ser Havunders* 17:109–131
- Gjøsæter J (1981b) Growth, production and reproduction of the myctophid fish *Benthoosema glaciale* from western Norway and adjacent seas. *FiskDir Skr Ser Havunders* 17:79–108
- Gjøsæter H (1998) The population biology and exploitation of capelin (*Mallotus villosus*) in the Barents Sea. *Sarsia* 83:453–496

- Goodson MS, Giske J, Rosland R (1995) Growth and ovarian development of *Maurolicus muelleri* during spring. *Mar Biol* 124:185–195
- Gundersen AC, Kjesbu OS, Nedreaas KH, Stene A (1999) Fecundity of Northeast Arctic Greenland halibut (*Reinhardtius hippoglossoides*). *J Northwest Atl Fish Sci* 25:29–36
- Haug T (1990) Biology of the Atlantic halibut, *Hippoglossus hippoglossus* (L., 1758). *Adv Mar Biol* 26:3–70
- Haug T, Gulliksen B (1988) Fecundity and oocyte sizes in ovaries of female Atlantic halibut. *Sarsia* 73:259–261
- Hildebrandt N, Bergmann M, Knust R (2011) Longevity and growth efficiency of two deep-dwelling Arctic zoarcids and comparison with eight other zoarcid species from different climatic regions. *Polar Biol* 34:1523–1533
- Hovde SC, Albert OT, Nilssen EM (2002) Spatial, seasonal and ontogenetic variation in diet of Northeast Arctic Greenland halibut (*Reinhardtius hippoglossoides*). *ICES J Mar Sci* 59:421–437
- Huse G, Gjørseter H (1997) Fecundity of the Barents Sea capelin (*Mallotus villosus*). *Mar Biol* 130:309–313
- Husebø Å, Nøttestad L, Fosså JH, Furevik DM, Jørgensen SB (2002) Distribution and abundance of fish in deep-sea coral habitats. *Hydrobiologia* 471:91–99
- Il'inskiy EN, Balanov AA, Ivanov OA (1995) Rare mesopelagic fishes *Scopelosaurus harryi*, *Arctozenus rissoi*, *Magnisudis atlantica*, and *Tactostoma macropus* from the Northwest Pacific. 2. Spatial distribution and biology. *J Ichthyol* 35:1–19
- Jákupsstovu SH, Haug T (1988) Growth, sexual maturation, and spawning season of Atlantic halibut, *Hippoglossus hippoglossus*, in Faroese waters. *Fish Res* 6:201–215
- Jennings S, Reynolds JD, Mills SC (1998) Life history correlates of responses to fisheries exploitation. *Proc Biol Sci* 265:333–339
- Jennings S, Greenstreet SPR, Reynolds JD (1999) Structural change in an exploited fish community: a consequence of differential fishing effects on species with contrasting life histories. *J Anim Ecol* 68:617–627
- Karamushko OV (2008) Species composition and structure of the ichthyofauna of the Barents Sea. *J Ichthyol* 48:277–291
- Link JS, Bolles K, Milliken CG (2002) The feeding ecology of flatfish in the northwest Atlantic. *J Northwest Atl Fish Sci* 30:1–17
- Link JS, Fogarty MJ, Langton RW (2005) The trophic ecology of flatfishes. In: Gibson RN (ed) *Flatfishes: biology and exploitation*. Blackwell Publishing, Oxford, p 185–212
- Lønne OJ, Gulliksen B (1989) Size, age and diet of polar cod, *Boreogadus saida* (Lepechin 1773), in ice covered waters. *Polar Biol* 9:187–191
- MacPherson E (1980) Food and feeding of *Chimaera monstrosa*, Linnaeus, 1758, in the western Mediterranean. *J Cons Int Explor Mer* 39:26–29
- Magnussen E (2007) Interpopulation comparison of growth patterns of 14 fish species on Faroe Bank: Are all fishes on the bank fast-growing? *J Fish Biol* 71:453–475
- Marshall NB (1965) Systematic and biological studies of the macrourid fishes (Anacanthini-Teleostii). *Deep-Sea Res* 12:299–322
- McAllister DE, Anderson ME, Hunter JG (1981) Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska. *Can J Fish Aquat Sci* 38:821–839
- McMeans BC, Svavarsson J, Dennard S, Fisk AT (2010) Diet and resource use among Greenland sharks (*Somniosus microcephalus*) and teleosts sampled in Icelandic waters, using $\delta^{13}\text{C}$, $\delta^{15}\text{C}$, and mercury. *Can J Fish Aquat Sci* 67:1428–1438
- Møller PR, Jørgensen OA (2000) Distribution and abundance of eelpouts (Pisces, Zoarcidae) off west Greenland. *Sarsia* 85:23–48

- Moura T, Figueiredo I, Machado PB, Gordo LS (2004) Growth pattern and reproductive strategy of the holocephalan *Chimaera monstrosa* along the Portuguese continental slope. *J Mar Biol Assoc UK* 84:801–804
- Munk P, Nielsen JG (2005) Eggs and larvae of North Sea fishes. *Biofolia*, Frederiksberg
- Murua H (2003) Population structure, growth and reproduction of roughhead grenadier on the Flemish Cap and Flemish Pass. *J Fish Biol* 63:356–373
- Nash RDM (1986) Aspects of the general biology of Wahl's eelpout, *Lycodes vahlii gracilis* M. Sars, 1867 (Pisces: Zoarcidae) in Oslofjorden, Norway. *Sarsia* 71:289–296
- Nedreaas K (1990) Age determination of Northeast Atlantic *Sebastes* species. *J Cons Int Explor Mer* 47:208–230
- Newton AW, Armstrong DW (1974) A note on the fecundity of lemon sole. *ICES CM F:34*
- Oldham WS (1972) Biology of the Scotian Shelf cusk, *Brosme brosme*. *ICNAF Res Bull* 9:85–98
- Olsen E, Aanes S, Mehl S, Holst JC, Aglen A, Gjøsæter H (2010) Cod, haddock, saithe, herring, and capelin in the Barents Sea and adjacent waters: a review of the biological value of the area. *ICES J Mar Sci* 67:87–101
- Ottesen C (2004) Taxonomy, morphology and biology of *Triglops murrayi* and *Triglops nybelini* (family Cottidae) obtained at Svalbard and Jan Mayen. *Cand Scient thesis*, University of Tromsø
- Pauly D (1978) A preliminary compilation of fish length growth parameters. *Ber Inst Meereskd Christian-Albrechts-Univ Kiel* 55
- Pauly D (1980) On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *J Cons Int Explor Mer* 39:175–192
- Penney R (1987) Development of deep-water redfish (*Sebastes mentella*) larvae, with comparative notes on newly extruded Acadian redfish (*Sebastes fasciatus*) from the Flemish Cap. *Can J Zool* 65:1167–1180
- Pethon P (2005) *Aschehougs store fiskebok*. Aschehoug, Oslo [In Norwegian]
- Raciborski K (1984) Migrations, reproduction, growth and feeding of *Myoxocephalus scorpius* (L) in Gdansk Bay (South Baltic). *Pol Arch Hydrobiol* 41:109–118
- Raitt DFS, Hall WB (1967) On the fecundity of the redfish. *J Cons Int Explor Mer* 31:237–245
- Rasmussen OI, Giske J (1994) Life-history parameters and vertical distribution of *Maurolicus muelleri* in Masfjorden in summer. *Mar Biol* 120:649–664
- Robertson DA (1976) Planktonic stages of *Maurolicus muelleri* (Teleostei: Sternoptychidae) in New Zealand waters. *NZ J Mar Freshw Res* 10:311–328
- Salvanes AGV, Stockley BM (1996) Spatial variation of growth and gonadal developments of *Maurolicus muelleri* in the Norwegian Sea and in a Norwegian fjord. *Mar Biol* 126:321–332
- Sameoto DD (1988) Feeding of lantern fish *Benthosema glaciale* off the Nova Scotia Shelf. *Mar Ecol Prog Ser* 44:113–129
- Singh-Renton S, Bromley PJ (1999) Feeding of small whiting (*Merlangius merlangus*) in the central and southern North Sea. *J Mar Biol Assoc UK* 79:957–960
- Snyder RJ (1991) Migration and life histories of the threespine stickleback: evidence for adaptive variation in growth rate between populations. *Environ Biol Fishes* 31:381–388
- Stokesbury MJW, Hervey-Clark C, Gallant J, Block BA, Myers RA (2005) Movement and environmental preferences of Greenland sharks (*Somniosus microcephalus*) electronically tagged in the St. Lawrence Estuary, Canada. *Mar Biol* 148:159–165
- Tåning ÅV (1961) Larval and postlarval stages of *Sebastes* species and *Helicolenus dactylopterus*. *Rapp PV Reun Cons Int Explor Mer* 150:234–240
- Templeman W (1982) Development, occurrence and characteristics of egg capsules of the thorny skate, *Raja radiata*, in the Northwest Atlantic. *J Northwest Atl Fish Sci* 3:47–56

- von Dorrien CF (1993) Ökologie und Respiration ausgewählter arktischer Bodenfischarten. Ber Polarforsch 125:1–99
- Walker PA, Hislop JRG (1998) Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. ICES J Mar Sci 55:392–402
- Walsh SJ (1994) Life history traits and spawning characteristics in populations of long rough dab (American plaice) *Hippoglossoides platessoides* (Fabricius) in the North Atlantic. Neth J Sea Res 32:241–254
- Wienerroither R, Johannesen E, Dolgov A, Byrkjedal I and others (2011) Atlas of the Barents Sea fishes. IMR/PINRO Joint Report Series 1-2011