

The following supplement accompanies the article

## First representation of the trophic structure and functioning of the Portuguese continental shelf ecosystem: insights into the role of sardine

T. Veiga-Malta\*, D. Szalaj, M. M. Angélico, M. Azevedo, I. Farias, S. Garrido, S. Lourenço, A. Marçalo, V. Marques, A. Moreno, P. B. Oliveira, V. H. Paiva, N. Prista, C. Silva, L. Sobrinho-Gonçalves, J. Vingada, A. Silva

\*Corresponding author: timat@aqua.dtu.dk

Marine Ecology Progress Series 617-618: 323–340 (2019)

Table S1 Sources of the parameters used by functional group for the Portuguese continental shelf ecosystem model. Percentages before each species scientific names represent the relative abundance of each species within a group. When no relative abundance is given, such information was not available or the functional group has only one species.

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sup>∞</sup> and L-W	Diet
<b>1. Seabirds</b> [2.7% <i>Alca torda</i> (Linnaeus, 1758), 7.9% <i>Calonectris diomedea borealis</i> (Cory, 1881), 86.8% <i>Morus bassanus</i> (Linnaeus, 1758), 1.6% <i>Puffinus mauretanicus</i> (Lowe, 1921), 0.9% <i>Uria aalge</i> (Pontoppidan, 1763)]	0.009	0.1	57.64	0		
	From 2010 to 2012, Araújo et al. 2014	Christensen et al. 2009	Nilsson & Nilsson 1976	Estimated by the model	Spring 1971, Wanless et al. 1985, Granadeiro 1991, Genovart et al. 2003, Langston & Teuten 2013	Beja 1990, Käkälä et al. 2010, Paiva et al. 2010
<b>2. Minkie whale</b> [ <i>Balaenoptera acutorostrata</i> (Lacépède, 1804)]	0.061	0.03	35.113	0.371		
	From 2010 to 2012, Araújo et al. 2014	Christensen et al. 2009	From Markussen et al. 1992 daily food consumption estimates and mean weight in the area	Estimated by model	MarPro database	MarPro database

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sup>∞</sup> and L-W	Diet
<b>3. Common dolphin</b> [ <i>Delphinus delphis</i> (Linnaeus, 1758)]	0.038	0.08	31.293	0.2		
	From 2010 to 2012, Araújo et al. 2014	Christensen et al. 2009	From Begona Santos et al. 2014 daily food consumption estimates and mean weight in the area	Estimated by model	Fernández 2003	Stomach content analysis from stranded animals
<b>4. Striped dolphin</b> [ <i>Stenella coeruleoalba</i> (Meyen, 1833)]	0.005	0.080	30.977	0.105		
	From 2010 to 2012, Araújo et al. 2014	Christensen et al. 2009	From Begona Santos et al. 2014 daily food consumption estimates and mean weight in the area	Estimated by model	Fernández 2003	Stomach content analysis from stranded animals
<b>5. Bottlenose dolphin</b> [ <i>Tursiops truncatus</i> (Montagu, 1821)]	0.006	0.080	32.056	0.868		
	From 2010 to 2012, Araújo et al. 2014	Christensen et al. 2009	From Begona Santos et al. (2014) daily food consumption estimates and mean weight in the area	Estimated by model	Fernández (2003)	Stomach content analysis from stranded animals
<b>6. Harbor porpoise</b> [ <i>Phocoena phocoena</i> (Linnaeus, 1758)]	0.004	0.080	14.166	0.013		
	From 2010 to 2012, Araújo et al. 2014	Christensen et al. 2009	From Begona Santos et al. (2014) daily food consumption estimates and mean weight in the area	Estimated by model	MarPro Stranding database (2015)	Stomach content analysis from stranded animals
<b>7. Tunas</b> [ <i>Auxis rochei</i> (Risso, 1810), <i>Sarda sarda</i> (Bloch, 1793), <i>Thunnus thynnus</i> (Linnaeus, 1758)]	0.200	0.420	4.211	0.096		
	Sanchez & Olaso 2004, Coll et al. 2006 (considering that only a	Pauly 1980, F assumed 0.1	Pauly et al. 1990	Estimated by model	Santos et al. 2002, Relini et al. 2005, Rooker et al. 2007,	Campo et al. 2006, Coll et al. 2006, Logan et al. 2011,

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sub>∞</sub> and L-W	Diet
<b>8. Rays</b> [ <i>Leucoraja naevus</i> (Müller & Henle, 1841), <i>Raja brachyuran</i> (Günther, 1880), <i>Raja clavata</i> (Linnaeus, 1758), <i>Raja montagui</i> (Fowler, 1910)]	0.136	0.729	4.826	0.850	Zaboukas & Megalofonou 2007, Kahraman et al. 2011, ICCAT 2011	Bănanu et al. 2013, Fletcher et al. 2013
	portion of this tuna migrates through the Portuguese continental shelf ecosystem) Estimated by model	ICES 2012, Pauly 1980	Pauly et. 1990	Sanchez & Olaso 2004, Torres et al. 2013	ICES 2012	Farias et al. 2006
<b>9. Hake</b> [ <i>Merluccius merluccius</i> (Linnaeus, 1758)]	0.389	1.300	6.070	0.934		
	Bottom trawl surveys 2006-2009	ICES 2013	Pauly et. 1990	Estimated by model	Mendes et al. 2004, Mellon-Duval et al. 2010	Cabral & Murta 2002
<b>10. Squids</b> [ <i>Alloteuthis</i> sp. <i>Illex coindetii</i> (Vérany, 1839), <i>Loligo vulgaris</i> (Lamarck, 1798), <i>Todaropsis eblanae</i> (Ball, 1841)]	0.962	2.800	8.000	0.950		
	Estimated by model	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013		Bidder 1950, Rocha et al. 1994, Rasero et al. 1996, Coelho et al. 1997, Sanchez & Olaso 2004, Torres et al. 2013
<b>11. Benthic cephalopods</b> [ <i>Eledone cirrhosa</i> (Lamarck, 1798), <i>Octopus vulgaris</i> (Cuvier, 1797), <i>Sepia officinalis</i> (Linnaeus, 1758), <i>Sepioloa</i> sp.]	1.154	2.300	6.500	0.950		
	Estimated by model	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013		Sanchez & Olaso 2004, Lassalle et al. 2011, Torres et al. 2013
<b>12. Horse mackerel</b> [ <i>Trachurus trachurus</i> (Linnaeus, 1758)]	2.841	0.512	7.607	0.935		
	Acoustic survey 2008-	Pauly 1980, F	Pauly et al. 1990	Estimated by	Costa 2004,	Cabral & Murta

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sub>∞</sub> and L-W	Diet
	2009	from official landings information for the study area and period (2006-2009)		model	Mendes et al. 2004	2002
<b>13. Blue jack mackerel</b> [ <i>Trachurus picturatus</i> (Bowdich, 1825)]	0.784	0.536	7.208	0.901		
	Bottom trawl surveys 2006-2009	Pauly 1980, F from official landings information for the study area and period (2006-2009)	Pauly et al. 1990	Estimated by model	Mendes et al. 2004, Vasconcelos et al. 2006	Deudero, 2001, Hirsch, 2009, www.fishbase.org
<b>14. Chub mackerel</b> [ <i>Scomber colias</i> (Gmelin, 1789)]	6.578	0.655	6.816	0.915		
	Acoustic surveys 2007-2009	Pauly 1980, F from official landings information for the study area and period (2006-2009)	Pauly et al. 1990	Estimated by model	Mendes et al. 2004, Velasco et al. 2011	Garrido et al. 2015
<b>15. Mackerel</b> [ <i>Scomber scombrus</i> (Linnaeus, 1758)]	0.448	0.804	7.137	0.942		
	Bottom trawl surveys 2006-2009	Pauly 1980, F from official landings information for the study area and period (2006-2009)	Pauly et al. 1990	Estimated by model	Mendes et al. 2004, Villamor et al. 2004	Cabral & Murta 2002
<b>16. Demersal piscivorous fish</b> [ <i>Chelidonichthys lucernus</i> (Linnaeus, 1758), <i>Conger conger</i> (Linnaeus, 1758), <i>Dicentrarchus labrax</i> (Linnaeus, 1758), <i>Helicolenus dactylopterus</i> (Delaroche,	0.531	0.644	3.575	0.950		

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sup>∞</sup> and L-W	Diet
1809), <i>Lophius budegassa</i> (Spinola, 1807), <i>Lophius piscatorius</i> (Spinola, 1807)]	Estimated by model	Pauly 1980 and considering the models from neighbour areas (Lassalle et al. 2011, Sanchez & Olaso 2004, Torres et al. 2013)	Pauly et al. 1990	Sanchez & Olaso 2004, Torres et al. 2013	Gonçalves et al. 1997, Landa et al. 2001, Mendes et al. 2004, García-Rodríguez et al. 2005, Olim & Borges 2006, Cicek et al. 2008, Correia et al. 2009, Sequeira et al. 2009, ICES 2013, <a href="http://www.fishbase.org">www.fishbase.org</a>	Sanchez & Olaso 2004, Preciado et al. 2006, Xavier et al. 2010, Neves et al. 2012, Castro et al. 2013, Spitz et al. 2013, Torres et al. 2013
<b>17. Demersal invertivorous fish</b> [ <i>Callanthias ruber</i> , <i>Callionymus lyra</i> , <i>Cepola macrophthalma</i> , <i>Chelidonichthys cuculus</i> , <i>Lepidotrigla cavillone</i> , <i>Mullus barbatus</i> , <i>Mullus surmuletus</i> , <i>Serranus hepatus</i> , <i>Synchiropus phaeton</i> , <i>Trigla lyra</i> , Ammodytidae, Gobiidae]	5.773	1.221	6.677	0.950		
	Estimated by model	Pauly 1980, F assumed 0.5	Pauly et al. 1990	Sanchez & Olaso 2004, Torres et al. 2013	Reñones et al. 1995, Borges et al. 2003, Mendes et al. 2004, Olim & Borges 2006, Dulčić et al. 2007, <a href="http://www.fishbase.org">www.fishbase.org</a>	Terrats et al. 2000, Mackinson & Daskalov 2007, Castro 2008, Hirsch 2009, Lopez-Lopez et al. 2011, Castro et al. 2013, Torres 2013, <a href="http://www.fishbase.org">www.fishbase.org</a>
<b>18. Benthopelagic piscivorous fish</b> [ <i>Belone belone</i> (Linnaeus, 1760), <i>Lepidopus caudatus</i> (Euphrasen, 1788), <i>Phycis blennoides</i> (Brünnich, 1768), <i>Phycis phycis</i> (Linnaeus, 1766), <i>Zeus faber</i> (Linnaeus, 1758)]	0.336	0.833	4.098	0.950		
	Estimated by model	Pauly 1980,	Pauly et. 1990	Sanchez &	Demestre et al.	Demestre et al.

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sub>∞</sub> and L-W	Diet
		Sanchez & Olaso 2004, Torres et al. 2013 and Official landings information for the study area and period (2006-2009)		Olaso 2004, Torres et al. 2013	1993, Casas & Piñeiro 2000, Borges et al. 2003, Mendes et al. 2004, Vieira et al. 2014, www.fishbase.org	1993, Morato et al. 1999, Silva 1999, Morte et al. 2002, Klimpel et al. 2006, Sever et al. 2009, Torres 2013
<b>19. Benthopelagic invertivorous fish</b> [ <i>Anthias anthias</i> (Linnaeus, 1758), <i>Argentina sphyraena</i> (Linnaeus, 1758), <i>Capros aper</i> (Linnaeus, 1758), <i>Gadiculus argenteus</i> (Guichenot, 1850), <i>Macroramphosus</i> spp. <i>Micromesistius poutassou</i> (Risso, 1827), <i>Trisopterus luscus</i> (Linnaeus, 1758)]	6.622	0.954	7.096	0.950		
	Estimated by model	Pauly 1980, F assumed 0.3	Pauly et al. 1990	Sanchez & Olaso 2004, Torres et al. 2013	Merayo & Villegas 1994, Borges 2000, Borges et al. 2003, Mendes et al. 2004, Hüseyin et al. 2012, www.fishbase.org	Cabral & Murta 2002, Oliveira 2002, Lopes et al. 2006, Hirsch 2009, Sever et al. 2013, www.fishbase.org
<b>20. Flatfish</b> [ <i>Arnoglossus laterna</i> (Walbaum, 1792), <i>Citharus linguatula</i> (Linnaeus, 1758), <i>Dicologlossa cuneata</i> (Moreau, 1881), <i>Lepidorhombus boscii</i> (Risso, 1810), <i>Lepidorhombus whiffiagonis</i> (Walbaum, 1792), <i>Microchirus azevia</i> (Capello, 1867), <i>Microchirus variegatus</i> (Donovan, 1808), <i>Pegusa lascaris</i> (Risso, 1810), <i>Platichthys flesus</i> (Linnaeus, 1758), <i>Scophthalmus maximus</i> (Linnaeus, 1758), <i>Scophthalmus rhombus</i> (Linnaeus, 1758), <i>Solea senegalensis</i> (Kaup, 1858), <i>Solea solea</i> (Linnaeus, 1758)]	0.908	0.940	5.687	0.950		
	Estimated by model	Pauly 1980, F assumed 0.3	Pauly et al. 1990	Sanchez & Olaso 2004,	Santos 1995, Andrade 1998,	Cabral 2000, Cabral et al. 2002,

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sup>∞</sup> and L-W	Diet
				Torres et al. 2013	Landa & Piñeiro 2000, Mendes et al. 2004, Teixeira et al. 2009, Teixeira & Cabral 2010, Teixeira et al. 2010, Félix et al. 2011, ICES 2013, www.fishbase.org	Amezcuca et al. 2003, Šantić et al. 2009, Teixeira et al. 2009, Teixeira et al. 2010, Vinagre et al. 2011, Torres 2013, Torres et al. 2013
<b>21. Sparids</b> [0.01% <i>Diplodus annularis</i> (Linnaeus, 1758), 0.05% <i>Diplodus Bellottii</i> (Steindachner, 1882), 0.01% <i>Diplodus sargus</i> (Linnaeus, 1758), 26.4% <i>Diplodus vulgaris</i> (Geoffroy Saint-Hilaire, 1817), 55.9% <i>Pagellus acarne</i> (Risso, 1827), 0.73% <i>Pagellus bogaraveo</i> (Brünnich, 1768), 3% <i>Pagellus Erythrinus</i> (Linnaeus, 1758), 0.2% <i>Pagrus auriga</i> (Valenciennes, 1843), 1.1% <i>Pagrus pagrus</i> (Linnaeus, 1758), 12.5% <i>Spondylisoma cantharus</i> (Linnaeus, 1758)]	0.599	0.725	5.425	0.858		
	Bottom trawl surveys 2006-2009	Pauly 1980, F from official landings information for the study area and period (2006-2009)	Pauly et al. 1990	Estimated by model	Gonçalves et al. 1997, Pajuelo & Lorenzo 2000, Erzini et al. 2001, Santos et al. 2002, Borges et al. 2003, Gonçalves et al. 2003, Mendes et al. 2004, Pajuelo et al. 2006, Matić-Skoko et al. 2007, Coelho et al. 2010, Lorance 2011, www.fishbase.org	Chakroun-Marzouk & Kartas 1987, Gonçalves & Erzini 1998, Pita et al. 2002, Torres 2013
<b>22. Anchovy</b> [ <i>Engraulis encrasicolus</i> (Linnaeus, 1758)]	0.305	1.170	12.220	0.919		

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sub>∞</sub> and L-W	Diet
	Acoustic surveys 2006-2009	Pauly 1980, F from official landings information for the study area and period (2006-2009)	Pauly et al. 1990	Estimated by model	Basilone et al. 2004, www.fishbase.org	Sanchez & Olaso 2004, Lassalle et al. 2011, Castro et al. 2013, Torres et al. 2013
<b>23. Sardine</b> [ <i>Sardina pilchardus</i> (Walbaum, 1792)]	14.760	0.891	9.677	0.956		
	Acoustic surveys 2006-2009	Pauly 1980, F from official landings information for the study area and period (2006-2009)	Pauly et al. 1990	Estimated by model	Mendes et al. 2004, Silva et al. 2008	Garrido et al. 2008
<b>24. Bogue</b> [ <i>Boops boops</i> (Linnaeus, 1758)]	0.794	1.246	6.201	0.950		
	Estimated by model	Pauly 1980, F from Monteiro et al. 2006	Pauly et al. 1990	Sanchez & Olaso 2004, Torres et al. 2013	Mendes et al. 2004, Monteiro et al. 2006	Derbal & Kara 2008, Bănarău et al. 2013, Dobrosłavić et al. 2013
<b>25. Henslow's crab</b> [ <i>Polybius henslowii</i> (Leach, 1820)]	0.526	2.100	10.640	0.950		
	Estimated by model	Coll et al. 2006, Torres et al. 2013	Cammen 1980	Sanchez & Olaso 2004, Torres et al. 2013	Acoustic and bottom trawl surveys 2006-2009	Signa et al. 2008, Torres et al. 2013
<b>26. Shrimps</b> [Crustacea natantia]	3.619	4.200	11.670	0.950		
	Estimated by model	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013		Sanchez & Olaso 2004, Torres et al. 2013
<b>27. Macrozoobenthos</b> [Starfishes, Annelids, Sea urchins, Sea cucumbers, Bivalves, Crustaceans, Sea anemone, other benthic invertebrates]	14.540	3.555	9.727	0.842		
	López-Jamar et al.	Sanchez & Olaso	Sanchez & Olaso	Estimated by		Sanchez & Olaso



<b>Species / Parameters</b>	<b>B (t·km<sup>-2</sup>)</b>	<b>P/B (year<sup>-1</sup>)</b>	<b>Q/B (year<sup>-1</sup>)</b>	<b>EE</b>	<b>Mean weight/L<sub>∞</sub> and L-W</b>	<b>Diet</b>
	1992, Martins et al. 2013	2004, Torres et al. 2013	2004, Torres et al. 2013	model		2004, Torres et al. 2013
<b>28. Suprabenthic invertebrates</b> [Euphausiids, Mysids, Isopods, Amphipods]	3.599	16.000	42.000	0.950		
	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013		Sanchez & Olaso 2004, Torres et al. 2013
<b>29. Macrozooplankton</b> [Zooplankton length > 3 mm, width > 1 mm]	5.253	17.000	44.000	0.781		
	Spring and Autumn surveys 2006-2009, Conversion factors were used to estimate wet weight of those samples (Cushing et al. 1958, Wiebe et al. 1975, Wiebe 1988, Alcaraz et al. 2003). The conversion to t/m <sup>2</sup> from t/m <sup>3</sup> was made based on information about vertical distribution of zooplankton in the western Portuguese continental shelf (L. Sobrinho-Gonçalves unpubl. Data, Sobrinho-Gonçalves et al. 2013)	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Estimated by model		Sanchez & Olaso 2004, Torres et al. 2013
<b>30. Meso e Microzooplankton</b> [Zooplankton length ≤ 3 mm, width ≤ 1 mm]	25.29	30.100	152.500	0.635		
	Mesozooplankton: Spring and Autumn surveys 2006-2009,	Sanchez & Olaso 2004, Torres et al. 2013	Sanchez & Olaso 2004, Torres et al. 2013	Estimated by model		Sanchez & Olaso 2004, Torres et al. 2013

Species / Parameters	B (t·km <sup>-2</sup> )	P/B (year <sup>-1</sup> )	Q/B (year <sup>-1</sup> )	EE	Mean weight/L <sub>∞</sub> and L-W	Diet
	<p>Microzooplankton: L. Sobrinho-Gonçalves unpubl. data, 2010-2011</p> <p>Conversion factors were used to estimate wet weight of those samples (Cushing et al. 1958, Wiebe et al. 1975, Wiebe 1988, Alcaraz et al. 2003). The conversion to t/m<sup>2</sup> from t/m<sup>3</sup> was made based on information about vertical distribution of zooplankton in the western Portuguese continental shelf (L. Sobrinho-Gonçalves unpubl. Data, Sobrinho-Gonçalves et al. 2013)</p>					
<b>31. Phytoplankton</b>	<p>18.720</p> <p>chlorophyll-a satellite data from the Globcolour project (algorithm by Morel &amp; Brethon 1989; www.globcolour.info)</p> <p>Conversion factors (Jorgensen et al. 1991, Dalsgaard &amp; Pauly 1997)</p>	<p>350.800</p> <p>Vertically Generalized Production Model (VGPM) (Behrenfeld &amp; Falkowski 1997)</p>		<p>0.432</p> <p>Estimated by model</p>		
<b>32. Discards</b>	<p>1.003</p> <p>Borges et al. 2001, Batista et al. 2009, Prista et al. 2012, Feijó</p>			<p>0.933</p> <p>Estimated by model</p>		

<b>Species / Parameters</b>	<b>B (t·km<sup>-2</sup>)</b>	<b>P/B (year<sup>-1</sup>)</b>	<b>Q/B (year<sup>-1</sup>)</b>	<b>EE</b>	<b>Mean weight/L<sup>∞</sup> and L-W</b>	<b>Diet</b>
	2013, Fernandes et al. 2015					
<b>33. Detritus</b>	45.230 Pauly et al. 1993			0.068 Estimated by model		

## References

- Alcaraz M, Saiz E, Calbet A, Trepas I, Broglio E (2003) Estimating zooplankton biomass through image analysis. *Mar Biol* 143:307–315 [doi:10.1007/s00227-003-1094-8](https://doi.org/10.1007/s00227-003-1094-8)
- Amezcuá FNRDM, Nash RDM, Veale L (2003) Feeding habits of the Order Pleuronectiformes and its relation to the sediment type in the north Irish Sea. *J Mar Biol Assoc UK* 83:593–601 [doi:10.1017/S0025315403007525h](https://doi.org/10.1017/S0025315403007525h)
- Andrade JP (1998) Age and growth of the bastard sole, *Microchirus azevia* (Capello, 1868) (Pisces, Soleidae) from the south coast of Portugal. *Fish Res* 34:205–208 [doi:10.1016/S0165-7836\(97\)00093-3](https://doi.org/10.1016/S0165-7836(97)00093-3)
- Araújo H, Santos J, Rodrigues P, Vingada J, Eira C, Rainho A, Arriegas I, Leonardo T, Nunes M, Sequeira M (2014) First Technical proposal of new SCI sites dedicated to the conservation of cetaceans in Portugal for inclusion in the National List of Sites. Annex to the 2014 Report of project LIFE MarPro NAT/PT/00038, pp. 285.
- Bănaru D, Mellon-Duval C, Roos D, Bigot JL and others (2013) Trophic structure in the Gulf of Lions marine ecosystem (north-western Mediterranean Sea) and fishing impacts. *J Mar Syst* 111-112:45–68 [doi:10.1016/j.jmarsys.2012.09.010](https://doi.org/10.1016/j.jmarsys.2012.09.010)
- Basilone G, Patti B, Mazzola S, Cuttitta A, Bonanno A, Sposito P, Kallianiotis A (2004) Growth of the European anchovy (*Engraulis encrasicolus*) in the Strait of Sicily. *MedSudMed Tech. Doc.* 5:79–89
- Behrenfeld MJ, Falkowski PG (1997) Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnol Oceanogr* 42:1–20 [doi:10.4319/lo.1997.42.1.0001](https://doi.org/10.4319/lo.1997.42.1.0001)
- Beja PR (1990) A note on the diet of Razorbills *Alca torda* wintering off Portugal. *Seabird* 12:11–13
- Bidder AM (1950) The digestive mechanism of the European squids *Loligo vulgaris*, *Loligo forbesii*, *Alloteuthis media* and *Alloteuthis subulata*. *J Cell Sci* 3:1–43
- Borges L (2000) Age and growth of the snipefish, *Macrorhamphosus* spp., in the Portuguese continental waters. *J Mar Biol Assoc UK* 80:147–153 [doi:10.1017/S0025315499001666](https://doi.org/10.1017/S0025315499001666)
- Borges TC, Olim S, Erzini K (2003) Weight–length relationships for fish species discarded in commercial fisheries of the Algarve (southern Portugal). *J Appl Ichthyology* 19:394–396 [doi:10.1111/j.1439-0426.2003.00480.x](https://doi.org/10.1111/j.1439-0426.2003.00480.x)
- Cabral HN (2000) Comparative feeding ecology of sympatric *Solea solea* and *S. senegalensis*, within the nursery areas of the Tagus estuary, Portugal. *J Fish Biol* 57:1550–1562 [doi:10.1111/j.1095-8649.2000.tb02231.x](https://doi.org/10.1111/j.1095-8649.2000.tb02231.x)
- Cabral HN, Murta AG (2002) The diet of blue whiting, hake, horse mackerel and mackerel off Portugal. *J Appl Ichthyology* 18:14–23 [doi:10.1046/j.1439-0426.2002.00297.x](https://doi.org/10.1046/j.1439-0426.2002.00297.x)
- Cabral HN, Lopes M, Loeper R (2002) Trophic niche overlap between flatfishes in a nursery area on the Portuguese coast. *Sci Mar* 66:293–300 [doi:10.3989/scimar.2002.66n3293](https://doi.org/10.3989/scimar.2002.66n3293)
- Cammen LM (1979) Ingestion rate: an empirical model for aquatic deposit feeders and detritivores. *Oecologia* 44:303–310 [doi:10.1007/BF00545232](https://doi.org/10.1007/BF00545232)
- Campo D, Mostarda E, Castriota L, Scarabello MP, Andaloro F (2006) Feeding habits of the Atlantic bonito, *Sarda sarda* (Bloch, 1793) in the southern Tyrrhenian sea. *Fish Res* 81:169–175 [doi:10.1016/j.fishres.2006.07.006](https://doi.org/10.1016/j.fishres.2006.07.006)

- Casas JM, Piñeiro C (2000) Growth and age estimation of greater fork-beard (*Phycis blennoides* Brünnich, 1768) in the north and northwest of the Iberian Peninsula (ICES Division VIIIc and IXa). *Fish Res* 47:19–25 [doi:10.1016/S0165-7836\(00\)00108-9](https://doi.org/10.1016/S0165-7836(00)00108-9)
- Castro N (2008) Estrutura e Ecologia alimentar das comunidades piscícolas em uma área de viveiro para espécies de interesse comercial da região centro de Portugal. MSc thesis, University of Lisboa, Lisboa.
- Castro N, Costa JL, Domingos I, Angélico MM (2013) Trophic ecology of a coastal fish assemblage in Portuguese waters. *J Mar Biol Assoc UK* 93:1151–1161 [doi:10.1017/S0025315412001853](https://doi.org/10.1017/S0025315412001853)
- Chakroun-Marzouk N, Kartas F (1987) Denture et régime alimentaire des espèces du genre *Pagrus* (Pisces, Sparidae) des côtes Tunisiennes. *Cybium* 11:3–19
- Christensen V, Walters CJ, Ahrens R, Alder J and others (2009) Database-driven models of the world's Large Marine Ecosystems. *Ecol Modell* 220:1984–1996 [doi:10.1016/j.ecolmodel.2009.04.041](https://doi.org/10.1016/j.ecolmodel.2009.04.041)
- Cicek E, Avsar D, Ozyurt CE, Yeldan H, Manasirli M (2008) Age, growth, reproduction and mortality of Tub Gurnard (*Chelidonichthys lucernus* (Linnaeus, 1758)) inhabiting in Babadillimani Bight (northeastern Mediterranean coast of Turkey). *J Biol Sci* 8:155–160 [doi:10.3923/jbs.2008.155.160](https://doi.org/10.3923/jbs.2008.155.160)
- Coelho M, Domingues P, Balguerias E, Fernandez M, Andrade JP (1997) A comparative study of the diet of *Loligo vulgaris* (Lamarck, 1799)(Mollusca: Cephalopoda) from the south coast of Portugal and the Saharan Bank (Central-East Atlantic). *Fish Res* 29:245–255 [doi:10.1016/S0165-7836\(96\)00540-1](https://doi.org/10.1016/S0165-7836(96)00540-1)
- Coelho R, Bentes L, Correia C, Gonçalves J and others (2010) Life history of the common pandora, *Pagellus erythrinus* (Linnaeus, 1758)(Actinopterygii: Sparidae) from southern Portugal. *Braz J Oceanogr* 58:233–245 [doi:10.1590/S1679-87592010000300006](https://doi.org/10.1590/S1679-87592010000300006)
- Coll M, Palomera I, Tudela S, Sardà F (2006) Trophic flows, ecosystem structure and fishing impacts in the South Catalan Sea, Northwestern Mediterranean. *J Mar Syst* 59:63–96 [doi:10.1016/j.jmarsys.2005.09.001](https://doi.org/10.1016/j.jmarsys.2005.09.001)
- Correia AT, Manso S, Coimbra J (2009) Age, growth and reproductive biology of the European conger eel (*Conger conger*) from the Atlantic Iberian waters. *Fish Res* 99:196–202 [doi:10.1016/j.fishres.2009.06.002](https://doi.org/10.1016/j.fishres.2009.06.002)
- Costa AM (2004) Idade e crescimento do carapau (*Trachurus trachurus* L.) da costa portuguesa no período de 1992 a 1998. *Relatórios Científicos e Técnicos, IPIMAR, Série digital 18*, 25 pp.
- Cushing DH, Humphrey GF, Banse K, Laevastu T (1958) Report of the committee on terms and equivalents. *Rapp P-V Reun Cons Int Explor Mer* 144:15–16
- Dalsgaard J, Pauly D (1997) Preliminary mass-balance model of Prince William Sound, Alaska for the pre-spill period, 1980-1989. Fisheries Centre, University of British Columbia, Vancouver. *British Columbia, Canada*.
- Demestre M, Moli B, Recasens L, Sánchez P (1993) Life history and fishery of *Lepidopus caudatus* (Pisces: Trichiuridae) in the Catalan Sea (northwestern Mediterranean). *Mar Biol* 115:23–32 [doi:10.1007/BF00349382](https://doi.org/10.1007/BF00349382)
- Derbal F, Kara MH (2008) Composition du régime alimentaire du bogue *Boops boops* (Sparidae) dans le golfe d'Annaba (Algérie). *Cybium* 32:325–333

- Deudero S (2001) Interspecific trophic relationships among pelagic fish species underneath FADs. *J Fish Biol* 58:53–67 [doi:10.1111/j.1095-8649.2001.tb00498.x](https://doi.org/10.1111/j.1095-8649.2001.tb00498.x)
- Dobrosravić T, Zlatović A, Bartulović V, Lučić D, Glamuzina B (2013) Diet overlap of juvenile salema (*Sarpa salpa*), bogue (*Boops boops*) and common two-banded sea bream (*Diplodus vulgaris*) in the south-eastern Adriatic. *J Appl Ichthyology* 29:181–185 [doi:10.1111/j.1439-0426.2012.02046.x](https://doi.org/10.1111/j.1439-0426.2012.02046.x)
- Dulčić J, Matić-Skoko S, Paladin A, Kraljević M (2007) Age, growth and mortality of brown comber, *Serranus hepatus* (Linnaeus, 1758)(Pisces: Serranidae), in the eastern Adriatic (Croatian coast). *J Appl Ichthyology* 23:195–197 [doi:10.1111/j.1439-0426.2006.00840.x](https://doi.org/10.1111/j.1439-0426.2006.00840.x)
- Erzini K, Bentes L, Lino PG, Ribeiro J and others (2001) Age and growth of seven sparid species of the South coast of Portugal. In Tenth European Congress on Ichthyology ECI X, Prague, Czech Republic.
- Farias I, Figueiredo I, Moura T, Gordo LS, Neves A, Serra-Pereira B (2006) Diet comparison of four ray species (*Raja clavata*, *Raja brachyura*, *Raja montagui* and *Leucoraja naevus*) caught along the Portuguese continental shelf. *Aquat Living Resour* 19:105–114 [doi:10.1051/alr:2006010](https://doi.org/10.1051/alr:2006010)
- Félix PM, Vinagre C, Cabral HN (2011) Life-history traits of flatfish in the Northeast Atlantic and Mediterranean Sea. *J Appl Ichthyology* 27:100–111 [doi:10.1111/j.1439-0426.2010.01623.x](https://doi.org/10.1111/j.1439-0426.2010.01623.x)
- Fernández AL (2003) Estatus dos pequenos cetáceos da plataforma de Galicia. PhD dissertation, University of Santiago de Compostela, Santiago de Compostela, 373 pp.
- Fletcher N, Batjakas IE, Pierce GJ (2013) Diet of the Atlantic bonito *Sarda sarda* (Bloch, 1793) in the Northeast Aegean Sea. *J Appl Ichthyology* 29:1030–1035 [doi:10.1111/jai.12164](https://doi.org/10.1111/jai.12164)
- García-Rodríguez M, Pereda P, Landa J, Esteban A (2005) On the biology and growth of the anglerfish *Lophius budegassa* Spinola, 1807 in the Spanish Mediterranean: a preliminary approach. *Fish Res* 71:197–208 [doi:10.1016/j.fishres.2004.08.033](https://doi.org/10.1016/j.fishres.2004.08.033)
- Garrido S, Ben-Hamadou R, Oliveira PB, Cunha ME, Chícharo MA, van der Lingen CD (2008) Diet and feeding intensity of sardine *Sardina pilchardus*: correlation with satellite-derived chlorophyll data. *Mar Ecol Prog Ser* 354:245–256 [doi:10.3354/meps07201](https://doi.org/10.3354/meps07201)
- Garrido S, Silva A, Pastor J, Dominguez R, Silva A, Santos A (2015) Trophic ecology of pelagic fish species off the Iberian coast: diet overlap, cannibalism and intraguild predation. *Mar Ecol Prog Ser* 539:271–285. [doi:10.3354/meps11506](https://doi.org/10.3354/meps11506)
- Genovart M, McMinn M, Bowler D (2003) A discriminant function for predicting sex in the Balearic Shearwater. *Waterbirds* 26:72–76 [doi:10.1675/1524-4695\(2003\)026\[0072:ADFFPS\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2003)026[0072:ADFFPS]2.0.CO;2)
- Gonçalves JMS, Erzini K (1998) Feeding habits of the two-banded sea bream (*Diplodus vulgaris*) and the black sea bream (*Spondyllosoma gantharus*) (Sparidae) from the south-west coast of Portugal. *Cybium* 22:245–254
- Gonçalves JMS, Bentes L, Lino PG, Ribeiro J, Canario AV, Erzini K (1997) Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. *Fish Res* 30:253–256 [doi:10.1016/S0165-7836\(96\)00569-3](https://doi.org/10.1016/S0165-7836(96)00569-3)

- Gonçalves JMS, Bentes L, Coelho R, Correia C and others (2003) Age and growth, maturity, mortality and yield-per-recruit for two banded bream (*Diplodus vulgaris* Geoffr.) from the south coast of Portugal. *Fish Res* 62:349–359 [doi:10.1016/S0165-7836\(02\)00280-1](https://doi.org/10.1016/S0165-7836(02)00280-1)
- Granadeiro JP (1991) The breeding biology of Cory's Shearwater *Calonectris diomedea borealis* on Berlenga Island, Portugal. *Seabird* 13:30–39
- Hirsch S (2009) Trophic interactions at seamounts. PhD Dissertation, University of Hamburg, Hamburg, 211 pp.
- Hüssy K, Coad JO, Farrell ED, Clausen LW, Clarke MW (2012) Sexual dimorphism in size, age, maturation, and growth characteristics of boarfish (*Capros aper*) in the Northeast Atlantic. *ICES J Mar Sci* 69:1729–1735 [doi:10.1093/icesjms/fss156](https://doi.org/10.1093/icesjms/fss156)
- ICCAT (2011) Report of the 2010 Atlantic Bluefin tuna stock assessment session (Madrid, Spain, September 6 to 12, 2010). ICCAT, Collect. Vol. Sci. Pap. 66:505–714
- ICES (2012) Report of the Working Group on Elasmobranch Fishes (WGEF), 19–26 June 2012, Lisbon, Portugal. ICES CM 2012/ACOM:19, 547 pp.
- ICES (2013) Report of the Working Group on the Assessment of Southern Shelf Stocks of Hake, Monk and Megrin (WGHMM), 10 - 16 May 2013, ICES Headquarters, Copenhagen. ICES CM 2013/ACOM:11A, 11 pp.
- Jørgensen SE, Nielsen SN, Jørgensen LA (1991) Handbook of ecological parameters and ecotoxicology. Elsevier Science Publishers, Amsterdam, The Netherlands, pp.1-126
- Kahraman AE, Göktürk D, Karakulak FS (2011) Age and growth of bullet tuna, *Auxis rochei* (Risso), from the Turkish Mediterranean coasts. *Afr J Biotechnol* 10:3009–3013 [doi:10.5897/AJB10.2021](https://doi.org/10.5897/AJB10.2021)
- Käkelä R, Käkelä A, Martínez-Abraín A, Sarzo B and others (2010) Fatty acid signature analysis confirms foraging resources of a globally endangered Mediterranean seabird species: calibration test and application to the wild. *Mar Ecol Prog Ser* 398:245–258 [doi:10.3354/meps08291](https://doi.org/10.3354/meps08291)
- Klimpel S, Rückert S, Piatkowski U, Palm HW, Hanel R (2006) Diet and metazoan parasites of silver scabbard fish *Lepidopus caudatus* from the Great Meteor Seamount (North Atlantic). *Mar Ecol Prog Ser* 315:249–257 [doi:10.3354/meps315249](https://doi.org/10.3354/meps315249)
- Landa J, Piñeiro C (2000) Megrin (*Lepidorhombus whiffiagonis*) growth in the North-eastern Atlantic based on back-calculation of otolith rings. *ICES J Mar Sci* 57:1077–1090 [doi:10.1006/jmsc.2000.0702](https://doi.org/10.1006/jmsc.2000.0702)
- Landa J, Pereda P, Duarte R, Azevedo M (2001) Growth of anglerfish (*Lophius piscatorius* and *L. budegassa*) in Atlantic Iberian waters. *Fish Res* 51:363–376 [doi:10.1016/S0165-7836\(01\)00260-0](https://doi.org/10.1016/S0165-7836(01)00260-0)
- Langston RH, Teuten E, Butler A (2013) Foraging ranges of northern gannets *Morus bassanus* in relation to proposed offshore wind farms in the UK: 2010-2012. *RSPB report to DECC*.
- Lassalle G, Lobry J, Le Loc'H F, Bustamanten P and others (2011) Lower trophic levels and detrital biomass control the Bay of Biscay continental shelf food web: implications for ecosystem management. *Prog Oceanogr* 91:561–575 [doi:10.1016/j.pocean.2011.09.002](https://doi.org/10.1016/j.pocean.2011.09.002)
- Logan JM, Rodríguez-Marín E, Goni N, Barreiro S, Arrizabalaga H, Golet W, Lutcavage M (2011) Diet of young Atlantic bluefin tuna (*Thunnus thynnus*) in eastern and western Atlantic foraging grounds. *Mar Biol* 158:73–85 [doi:10.1007/s00227-010-1543-0](https://doi.org/10.1007/s00227-010-1543-0)

- Lopes M, Murta AG, Cabral HN (2006) The ecological significance of the zooplanktivores, snipefish *Macroramphosus* spp. and boarfish *Capros aper*, in the food web of the south-east North Atlantic. *J Fish Biol* 69:363–378 [doi:10.1111/j.1095-8649.2006.01093.x](https://doi.org/10.1111/j.1095-8649.2006.01093.x)
- López-Jamar E, Cal RM, González G, Hanson RB, Rey J, Santiago G, Tenore KR (1992) Upwelling and outwelling effects on the benthic regime of the continental shelf off Galicia, NW Spain. *J Mar Res* 50:465–488 [doi:10.1357/002224092784797584](https://doi.org/10.1357/002224092784797584)
- López-López L, Preciado I, Velasco F, Olaso I, Gutiérrez-Zabala JL (2011) Resource partitioning amongst five coexisting species of gurnards (Scorpaeniforme: Triglidae): Role of trophic and habitat segregation. *J Sea Res* 66:58–68 [doi:10.1016/j.seares.2011.04.012](https://doi.org/10.1016/j.seares.2011.04.012)
- Lorance P (2011) History and dynamics of the overexploitation of the blackspot sea bream (*Pagellus bogaraveo*) in the Bay of Biscay. *ICES J Mar Sci* 68:290–301 [doi:10.1093/icesjms/fsq072](https://doi.org/10.1093/icesjms/fsq072)
- Mackinson S, Daskalov G (2007) An ecosystem model of the North Sea to support an ecosystem approach to fisheries management: description and parameterisation. *Cefas Sci. Ser. Tech. Rep.* 142, p. 196.
- Markussen NH, Ryg M, Lydersen C (1992) Food consumption of the NE Atlantic minke whale (*Balaenoptera acutorostrata*) population estimated with a simulation model. *ICES J Mar Sci* 49:317–323 [doi:10.1093/icesjms/49.3.317](https://doi.org/10.1093/icesjms/49.3.317)
- Martins R, Quintino V, Rodrigues AM (2013) Diversity and spatial distribution patterns of the soft-bottom macrofauna communities on the Portuguese continental shelf. *J Sea Res* 83:173–186 [doi:10.1016/j.seares.2013.03.001](https://doi.org/10.1016/j.seares.2013.03.001)
- Matić-Skoko S, Kraljević M, Dulčić J, Jardas I (2007) Age, growth, maturity, mortality, and yield-per-recruit for annular sea bream (*Diplodus annularis* L.) from the eastern middle Adriatic Sea. *J Appl Ichthyology* 23:152–157 [doi:10.1111/j.1439-0426.2006.00816.x](https://doi.org/10.1111/j.1439-0426.2006.00816.x)
- Mellon-Duval C, De Pontual H, Métral L, Quemener L (2010) Growth of European hake (*Merluccius merluccius*) in the Gulf of Lions based on conventional tagging. *ICES J Mar Sci* 67:62–70 [doi:10.1093/icesjms/fsp215](https://doi.org/10.1093/icesjms/fsp215)
- Mendes B, Fonseca P, Campos A (2004) Weight–length relationships for 46 fish species of the Portuguese west coast. *J Appl Ichthyology* 20:355–361 [doi:10.1111/j.1439-0426.2004.00559.x](https://doi.org/10.1111/j.1439-0426.2004.00559.x)
- Merayo CR, Villegas ML (1994) Age and growth of *Trisopterus luscus* (Linnaeus, 1758)(Pisces, Gadidae) off the coast of Asturias. *Hydrobiologia* 281:115–122 [doi:10.1007/BF00006440](https://doi.org/10.1007/BF00006440)
- Monteiro P, Bentes L, Coelho R, Correia C and others (2006) Age and growth, mortality, reproduction and relative yield per recruit of the bogue, *Boops boops* Linné, 1758 (Sparidae), from the Algarve (south of Portugal) longline fishery. *J Appl Ichthyology* 22:345–352 [doi:10.1111/j.1439-0426.2006.00756.x](https://doi.org/10.1111/j.1439-0426.2006.00756.x)
- Morato T, Solà E, Grós MP, Menezes GM (1999) Diets of forkbeard (*Phycis phycis*) and conger eel (*Conger conger*) off the Azores during spring of 1996 and 1997. *Arquipel. Life Mar. Sci. (Suppl. 17A)*:51–64 ISSN 0873-4704
- Morte MS, Redón MJ, Sanz-Brau A (2002) Diet of *Phycis blennoides* (Gadidae) in relation to fish size and season in the western Mediterranean (Spain). *Mar Ecol* 23:141–155 [doi:10.1046/j.1439-0485.2002.02763.x](https://doi.org/10.1046/j.1439-0485.2002.02763.x)



- Neves A, Sequeira V, Paiva RB, Vieira AR, Gordo LS (2012) Feeding habits of the bluemouth, *Helicolenus dactylopterus dactylopterus* (Delaroche, 1809)(Pisces: Sebastidae) in the Portuguese coast. *Helgol Mar Res* 66:189-197 [doi:10.1007/s10152-011-0260-5](https://doi.org/10.1007/s10152-011-0260-5)
- Nilsson SG, Nilsson IN (1976) Numbers, food consumption, and fish predation by birds in Lake Möckeln, southern Sweden. *Ornis Scand* 7:61–70 [doi:10.2307/3676175](https://doi.org/10.2307/3676175)
- Olim S, Borges TC (2006) Weight–length relationships for eight species of the family Triglidae discarded on the south coast of Portugal. *J Appl Ichthyology* 22:257–259 [doi:10.1111/j.1439-0426.2006.00644.x](https://doi.org/10.1111/j.1439-0426.2006.00644.x)
- Oliveira MIS (2002) Dieta alimentar da faneca, *Trisopterus luscus* (Linnaeus, 1758) de Angeiras e da Aguda. MSc Thesis, University of Porto, Porto, 170 pp.
- Paiva VH, Xavier J, Geraldés P, Ramirez I, Garthe S, Ramos JA (2010) Foraging ecology of Cory's shearwaters in different oceanic environments of the North Atlantic. *Mar Ecol Prog Ser* 410:257–268 [doi:10.3354/meps08617](https://doi.org/10.3354/meps08617)
- Pajuelo BJ, Lorenzo JM (2000) Reproduction, age, growth and mortality of axillary seabream, *Pagellus acarne* (Sparidae), from the Canarian archipelago. *J Appl Ichthyology* 16:41–47 [doi:10.1046/j.1439-0426.2000.00154.x](https://doi.org/10.1046/j.1439-0426.2000.00154.x)
- Pajuelo JG, Socorro J, González JA, Lorenzo JM, Pérez-Peñalvo JA, Martínez I, Hernández-Cruz CM (2006) Life history of the red-banded seabream *Pagrus auriga* (Sparidae) from the coasts of the Canarian archipelago. *J Appl Ichthyology* 22:430–436 [doi:10.1111/j.1439-0426.2006.00748.x](https://doi.org/10.1111/j.1439-0426.2006.00748.x)
- Pauly D (1980) On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES J Mar Sci* 39:175–192 [doi:10.1093/icesjms/39.2.175](https://doi.org/10.1093/icesjms/39.2.175)
- Pauly D, Christensen V, Sambilay V Jr (1990) Some features of fish food consumption estimates used by ecosystem modelers. International Council for the Exploration of the Sea (ICES) C.M.1990/G: 17, Session O, 9 p.
- Pauly D, Soriano-Bartz M, Palomares ML (1993) Improved construction, parameterization and interpretation of steady-state ecosystem models. In: Christensen, V., Pauly, D. (Eds.), *Trophic Models of Aquatic Ecosystems*. ICLARM Conference Proceedings 26:1–13.
- Pita C, Gamito S, Erzini K (2002) Feeding habits of the gilthead seabream (*Sparus aurata*) from the Ria Formosa (southern Portugal) as compared to the black seabream (*Spondylisoma cantharus*) and the annular seabream (*Diplodus annularis*). *J Appl Ichthyology* 18:81–86 [doi:10.1046/j.1439-0426.2002.00336.x](https://doi.org/10.1046/j.1439-0426.2002.00336.x)
- Preciado I, Velasco F, Olaso I, Landa J (2006) Feeding ecology of black anglerfish *Lophius budegassa*: seasonal, bathymetric and ontogenetic shifts. *J Mar Biol Assoc UK* 86:877–884 [doi:10.1017/S0025315406013816](https://doi.org/10.1017/S0025315406013816)
- Rasero M, Gonzalez AF, Castro BG, Guerra A (1996) Predatory relationships of two sympatric squid, *Todaropsis eblanae* and *Illex coindetii* (Cephalopoda: Ommastrephidae) in Galician waters. *J Mar Biol Assoc UK* 76:73–87 [doi:10.1017/S0025315400029027](https://doi.org/10.1017/S0025315400029027)
- Relini LO, Garibaldi F, Cima C, Palandri G, Lanteri L, Relini M (2005) Biology of Atlantic bonito, *Sarda sarda* (Bloch, 1793), in the western and central Mediterranean a summary concerning a possible stock unit. *Collect. Vol. Sci. Pap. ICCAT* 58:575–588

- Reñones O, Massuti E, Morales-Nin B (1995) Life history of the red mullet *Mullus surmuletus* from the bottom-trawl fishery off the Island of Majorca (north-west Mediterranean). *Mar Biol* 123:411–419 [doi:10.1007/BF00349219](https://doi.org/10.1007/BF00349219)
- Rocha F, Castro BG, Gil MS, Guerra A (1994) The diets of *Loligo vulgaris* and *L. forbesi* (Cephalopoda: Loliginidae) in northwestern Spanish Atlantic waters. *Sarsia* 79:119–126 [doi:10.1080/00364827.1994.10413552](https://doi.org/10.1080/00364827.1994.10413552)
- Rooker JR, Alvarado Bremer JR, Block BA, Dewar H and others (2007) Life history and stock structure of Atlantic bluefin tuna (*Thunnus thynnus*). *Rev Fish Sci* 15:265–310 [doi:10.1080/10641260701484135](https://doi.org/10.1080/10641260701484135)
- Sánchez F, Olaso I (2004) Effects of fisheries on the Cantabrian Sea shelf ecosystem. *Ecol Modell* 172:151–174 [doi:10.1016/j.ecolmodel.2003.09.005](https://doi.org/10.1016/j.ecolmodel.2003.09.005)
- Šantić M, Podvinski M, Pallaoro A, Jardas I, Kirinčić M (2009) Feeding habits of megrim, *Lepidorhombus whiffiagonis* (Walbaum, 1792), from the central Adriatic Sea. *J Appl Ichthyology* 25:417–422 [doi:10.1111/j.1439-0426.2009.01257.x](https://doi.org/10.1111/j.1439-0426.2009.01257.x)
- Santos P (1995) Growth, mortality and maturation of *Lepidorhombus boscii* in Portuguese waters. ICES Demersal Fish Committee, CM. 1995/G:38, 20 pp.
- Santos MN, Gaspar MB, Vasconcelos P, Monteiro CC (2002) Weight–length relationships for 50 selected fish species of the Algarve coast (southern Portugal). *Fish Res* 59:289–295 [doi:10.1016/S0165-7836\(01\)00401-5](https://doi.org/10.1016/S0165-7836(01)00401-5)
- Begoña Santos M, Saavedra C, Pierce GJ (2014) Quantifying the predation on sardine and hake by cetaceans in the Atlantic waters of the Iberian peninsula. *Deep Sea Res II* 106:232–244 [doi:10.1016/j.dsr2.2013.09.040](https://doi.org/10.1016/j.dsr2.2013.09.040)
- Sequeira V, Neves A, Vieira AR, Figueiredo I, Gordo LS (2009) Age and growth of bluemouth, *Helicolenus dactylopterus*, from the Portuguese continental slope. *ICES J Mar Sci* 66:524–531 [doi:10.1093/icesjms/fsp010](https://doi.org/10.1093/icesjms/fsp010)
- Sever TM, Bayhan B, Bilge G, Taşkavak E (2009) Diet composition of *Belone belone* (Linnaeus, 1761)(Pisces: Belonidae) in the Aegean Sea. *J Appl Ichthyology* 25:702–706 [doi:10.1111/j.1439-0426.2009.01368.x](https://doi.org/10.1111/j.1439-0426.2009.01368.x)
- Sever TM, Bayhan B, Filiz H, Taşkavak E, Bilge G (2015) Diet composition of the five deep sea fish from the Aegean Sea. *Ege. J. Fish. Aqua. Sci.* 30:61–67
- Signa G, Cartes JE, Solé M, Serrano A, Sánchez F (2008) Trophic ecology of the swimming crab *Polybius henslowii* Leach, 1820 in Galician and Cantabrian Seas: Influences of natural variability and the Prestige oil spill. *Cont Shelf Res* 28:2659–2667 [doi:10.1016/j.csr.2008.08.008](https://doi.org/10.1016/j.csr.2008.08.008)
- Silva A (1999) Feeding habits of John Dory, *Zeus faber*, off the Portuguese continental coast. *J Mar Biol Assoc UK* 79:333–340 [doi:10.1017/S002531549800037X](https://doi.org/10.1017/S002531549800037X)
- Silva A, Carrera P, Masse J, Uriarte A and others (2008) Geographic variability of sardine growth across the northeastern Atlantic and the Mediterranean Sea. *Fish Res* 90:56–69 [doi:10.1016/j.fishres.2007.09.011](https://doi.org/10.1016/j.fishres.2007.09.011)
- Sobrinho-Gonçalves L, Moita MT, Garrido S, Cunha ME (2013) Environmental forcing on the interactions of plankton communities across a continental shelf in the Eastern Atlantic upwelling system. *Hydrobiologia* 713:167–182 [doi:10.1007/s10750-013-1500-2](https://doi.org/10.1007/s10750-013-1500-2)
- Spitz J, Chouvelon T, Cardinaud M, Kostecki C, Lorance P (2013) Prey preferences of adult sea bass *Dicentrarchus labrax* in the northeastern Atlantic: implications for bycatch of

- common dolphin *Delphinus delphis*. ICES J Mar Sci 70:452–461  
[doi:10.1093/icesjms/fss200](https://doi.org/10.1093/icesjms/fss200)
- Spring L (1971) A comparison of functional and morphological adaptations in the common murre (*Uria aalge*) and thick-billed murre (*Uria lomvia*). Condor 73:1–27  
[doi:10.2307/1366120](https://doi.org/10.2307/1366120)
- Teixeira CM, Cabral HN (2010) Comparative analysis of the diet, growth and reproduction of the soles, *Solea solea* and *Solea senegalensis*, occurring in sympatry along the Portuguese coast. J Mar Biol Assoc UK 90:995–1003 [doi:10.1017/S0025315410000238](https://doi.org/10.1017/S0025315410000238)
- Teixeira CM, Pinheiro A, Cabral HN (2009) Feeding ecology, growth and sexual cycle of the sand sole, *Solea lascaris*, along the Portuguese coast. J Mar Biol Assoc UK 89:621–627  
[doi:10.1017/S0025315409002562](https://doi.org/10.1017/S0025315409002562)
- Teixeira CM, Batista MI, Cabral HN (2010) Diet, growth and reproduction of four flatfishes on the Portuguese coast. Sci Mar 74:223–233 [doi:10.3989/scimar.2010.74n2223](https://doi.org/10.3989/scimar.2010.74n2223)
- Terrats A, Petrakis G, Papaconstantinou C (2000) Feeding habits of *Aspitrigla cuculus* (L., 1758)(red gurnard), *Lepidotrigla cavillone* (Lac., 1802)(large scale gurnard) and *Trigloporus lastoviza* (Brunn., 1768)(rock gurnard) around Cyclades and Dodecanese Islands (E. Mediterranean). Mediterr. Mar. Sci. 1108-393X 1(1):91-104.
- Torres MÁ (2013) Modelización ecológica del Golfo de Cádiz: relaciones tróficas, análisis de la estructura de la comunidad e impacto de la pesca en el ecosistema, PhD Dissertation, University of Cádiz, Cádiz, 177 pp.
- Torres MÁ, Coll M, Heymans JJ, Christensen V, Sobrino I (2013) Food-web structure of and fishing impacts on the Gulf of Cadiz ecosystem (South-western Spain). Ecol Modell 265:26–44 [doi:10.1016/j.ecolmodel.2013.05.019](https://doi.org/10.1016/j.ecolmodel.2013.05.019)
- Vasconcelos J, Alves A, Gouveia E, Faria G (2006) Age and growth of the blue jack mackerel, *Trachurus picturatus* Bowdich, 1825 (Pisces: Teleostei) off Madeira archipelago. Arquipel. Life Mar. Sci. (Suppl. 23):47–57
- Velasco EM, Del Arbol J, Baro J, Sobrino I (2011) Age and growth of the Spanish chub mackerel *Scomber colias* off southern Spain: a comparison between samples from the NE Atlantic and the SW Mediterranean. Rev Biol Mar Oceanogr 46:27–34  
[doi:10.4067/S0718-19572011000100004](https://doi.org/10.4067/S0718-19572011000100004)
- Vieira AR, Neves A, Sequeira V, Paiva RB, Gordo LS (2014) Age and growth of forkbeard, *Phycis phycis*, in Portuguese continental waters. J Mar Biol Assoc UK 94:623–630  
[doi:10.1017/S0025315413001549](https://doi.org/10.1017/S0025315413001549)
- Villamor B, Abaunza, Celso Fariña A (2004) Growth variability of mackerel (*Scomber scombrus*) off north and northwest Spain and a comparative review of the growth patterns in the northeast Atlantic. Fish Res 69:107–121 [doi:10.1016/j.fishres.2004.02.005](https://doi.org/10.1016/j.fishres.2004.02.005)
- Vinagre C, Silva A, Lara M, Cabral HN (2011) Diet and niche overlap of southern populations of brill *Scophthalmus rhombus* and turbot *Scophthalmus maximus*. J Fish Biol 79:1383–1391 [doi:10.1111/j.1095-8649.2011.03116.x](https://doi.org/10.1111/j.1095-8649.2011.03116.x)
- Wanless S, Harris MP, Morris JA (1985) Radio-monitoring as a method for estimating time budgets of guillemots *Uria aalge*. Bird Study 32:170–175  
[doi:10.1080/00063658509476875](https://doi.org/10.1080/00063658509476875)
- Wiebe PH (1988) Functional regression equations for zooplankton displacement volume, wet weight, dry weight, and carbon: a correction. Fish Bull 86:833–835

- Wiebe PH, Boyd S, Cox JL (1975) Relationships between zooplankton displacement volume, wet weight, dry weight, and carbon. *Fish Bull* 73:777–786
- Xavier JC, Chérel Y, Assis CA, Sendão J, Borges TC (2010) Feeding ecology of conger eels (*Conger conger*) in north-east Atlantic waters. *J Mar Biol Assoc UK* 90:493–501  
[doi:10.1017/S0025315409990555](https://doi.org/10.1017/S0025315409990555)
- Zaboukas N, Megalofonou P (2007) Age estimation of the Atlantic bonito in the eastern Mediterranean Sea using dorsal spines and validation of the method. *Sci Mar* 71:691–698  
[doi:10.3989/scimar.2007.71n4691](https://doi.org/10.3989/scimar.2007.71n4691)

Table S2. Diet composition matrix adjusted for the Portuguese continental shelf food web of the Ecopath model.

Preys	Predators														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1 Seabirds	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2 Minkie whale	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3 Common dolphin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4 Striped dolphin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5 Bottlenose dolphin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6 Harbor porpoise	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
7 Tunas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8 Rays	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
9 Hake	0.1	0	0.042	0.1	0.084	0.152	0	0.012	0.02	0.01	0	0	0	0	0
10 Squids	0.01	0	0.047	0.139	0.003	0.003	0.01	0.093	0.01	0.05	0	0.005	0.01	0.02	0.01
11 Benthic cephalopods	0	0	0.008	0.037	0.011	0.001	0	0.021	0	0.004	0.03	0	0	0.03	0
12 Horse mackerel	0.014	0.2552	0.13	0.03	0.009	0.021	0.01	0.03	0.02	0	0	0	0	0	0
13 Blue jack mackerel	0	0.0888	0.025	0.005	0	0	0.01	0	0.01	0	0	0	0	0	0
14 Chub mackerel	0.126	0.108	0.233	0.104	0.007	0.023	0.09	0.002	0.059	0.163	0.083	0	0	0	0
15 Mackerel	0.25	0.002	0.047	0.038	0.004	0	0	0	0.001	0	0	0	0	0	0
16 Demersal piscivorous fish	0	0	0	0	0.673	0.073	0	0	0	0	0	0	0	0	0
17 Demersal invertivorous fish	0.122	0	0.002	0.013	0.004	0.106	0	0.227	0	0.116	0.212	0	0.01	0.01	0
18 Benthopelagic piscivorous fish	0	0	0.001	0	0	0	0	0	0.03	0	0	0	0	0	0
19 Benthopelagic invertivorous fish	0.078	0.021	0.057	0.202	0.013	0.051	0	0.172	0.51	0.183	0.015	0	0	0.02	0.1
20 Flatfish	0	0	0.003	0	0.015	0.026	0	0.025	0	0.031	0.035	0	0	0	0
21 Sparids	0	0.062	0	0	0	0	0	0	0	0.005	0.01	0	0	0	0
22 Anchovy	0.019	0.04	0.007	0.064	0	0	0.01	0	0.025	0.01	0	0	0	0	0.01
23 Sardine	0.253	0.357	0.372	0.226	0.002	0.012	0.37	0.095	0.255	0.261	0.275	0.03	0.02	0.03	0.03
24 Bogue	0	0	0.026	0.002	0.002	0.001	0	0	0.03	0.015	0.09	0	0	0	0
25 Henslow's crab	0	0	0	0	0	0	0	0.074	0	0.002	0.015	0	0	0	0
26 Shrimps	0	0	0	0	0	0	0	0.056	0.005	0.022	0.026	0	0	0	0
27 Macrozoobenthos	0	0	0	0	0	0	0	0.124	0	0.025	0.065	0.1	0.1	0.02	0
28 Suprabenthic invertebrates	0	0	0	0	0	0	0	0.014	0.005	0.034	0.066	0.44	0.32	0.097	0.16
29 Macrozooplankton	0	0	0	0	0	0	0	0.055	0.02	0.069	0.078	0.385	0.24	0.325	0.141
30 Meso and Microzooplankton	0	0	0	0	0	0	0	0	0	0	0	0.04	0.3	0.446	0.549
31 Phytoplankton	0	0	0	0	0	0	0	0	0	0	0	0	0	0.002	0
32 Discards	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0
33 Detritus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
34 Import	0.018	0.066	0	0.04	0.173	0.531	0.5	0	0	0	0	0	0	0	0

Table S2. (Continued)

Preys	Predators														
	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
1 Seabirds	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2 Minkie whale	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3 Common dolphin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4 Striped dolphin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5 Bottlenose dolphin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6 Harbor porpoise	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
7 Tunas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8 Rays	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
9 Hake	0.016	0	0.017	0	0.001	0	0	0	0	0	0	0	0	0	0
10 Squids	0.008	0	0.019	0.002	0.081	0.051	0	0	0	0.03	0	0	0	0	0
11 Benthic cephalopods	0.04	0.002	0.02	0	0.042	0.022	0	0	0	0	0	0	0	0	0
12 Horse mackerel	0.033	0	0	0	0	0	0	0	0	0	0	0	0	0	0
13 Blue jack mackerel	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
14 Chub mackerel	0.076	0	0.032	0	0	0	0	0	0	0	0	0	0	0	0
15 Mackerel	0.012	0	0	0	0	0	0	0	0	0	0	0	0	0	0
16 Demersal piscivorous fish	0.02	0	0.019	0	0	0	0	0	0	0	0	0	0	0	0
17 Demersal invertivorous fish	0.166	0.01	0.086	0.029	0.215	0.05	0	0	0	0	0	0	0	0	0
18 Benthopelagic piscivorous fish	0.078	0	0.004	0	0	0	0	0	0	0	0	0	0	0	0
19 Benthopelagic invertivorous fish	0.249	0	0.241	0.01	0.021	0	0	0	0	0	0	0	0	0	0
20 Flatfish	0.026	0	0.016	0	0.01	0.033	0	0	0	0	0	0	0	0	0
21 Sparids	0	0	0.001	0	0	0	0	0	0	0	0	0	0	0	0
22 Anchovy	0.004	0	0.011	0	0	0	0	0	0	0	0	0	0	0	0
23 Sardine	0.035	0	0.142	0.014	0.009	0	0	0	0	0	0	0	0	0	0
24 Bogue	0	0	0.014	0	0	0	0	0	0	0	0	0	0	0	0
25 Henslow's crab	0.01	0.015	0	0.001	0.003	0	0	0	0	0.038	0	0	0	0	0
26 Shrimps	0.115	0.115	0.211	0.079	0.139	0.022	0	0	0.4	0	0.005	0.017	0	0	0
27 Macrozoobenthos	0.037	0.368	0.085	0.069	0.366	0.588	0	0	0.15	0.41	0.28	0.02	0	0	0
28 Suprabenthic invertebrates	0.009	0.197	0.082	0.37	0.063	0.062	0.007	0	0.15	0.01	0.12	0.044	0	0	0
29 Macrozooplankton	0.066	0.174	0	0.265	0.05	0.152	0.021	0	0.25	0.255	0.05	0.091	0.04	0	0
30 Meso and Microzooplankton	0	0.094	0	0.161	0	0.02	0.97	0.833	0.05	0.163	0.09	0.021	0.2	0.7	0
31 Phytoplankton	0	0.01	0	0	0	0	0.002	0.167	0	0	0.03	0.079	0.08	0.3	0.95
32 Discards	0	0	0	0	0	0	0	0	0	0.01	0.004	0.005	0	0	0
33 Detritus	0	0.015	0	0	0	0	0	0	0	0.084	0.421	0.723	0.68	0	0.05
34 Import	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0